

# The smallest minimal blocking sets of $Q(2n, q)$ , for small odd $q$

Jan De Beule

Ghent University, Department of Pure Mathematics and Computer Algebra,  
Krijgslaan 281, 9000 Gent, Belgium.

*Joint work with: Leo Storme*

In [2] we used results on the size of the smallest minimal blocking sets of  $Q(4, q)$ ,  $q$  even (from [1]) and projection arguments to find the following characterization of the smallest minimal blocking sets of  $Q(6, q)$ ,  $q$  even,  $q \geq 32$ :

**Theorem 1** *Let  $\mathcal{K}$  be a minimal blocking set of  $Q(6, q)$ , different from an ovoid of  $Q(6, q)$ ,  $|\mathcal{K}| \leq q^3 + q$ . Then there is a point  $p \in Q(6, q) \setminus \mathcal{K}$  with the following property:  $T_p(Q(6, q)) \cap Q(6, q) = pQ(4, q)$  and  $\mathcal{K}$  consists of all the points of the lines  $L$  on  $p$  meeting  $Q(4, q)$  in an ovoid  $\mathcal{O}$ , minus the point  $p$  itself, and  $|\mathcal{K}| = q^3 + q$ .*

The results of [1] could be proven for  $q = 3$  and replaced by computer results for  $q = 5, 7$ . Using then the same projection arguments we found the above characterization for  $q = 3, 5, 7$ .

Using inductive arguments we can find results for  $Q(2n, q)$ ,  $q = 3, 5, 7$ . The situation is now very dependent of  $q$ , since for example  $Q(6, 3)$  has an ovoid, but  $Q(6, q)$ ,  $q = 5, 7$ , not. For  $q = 5, 7$ , we found the following characterization.

**Theorem 2** *Let  $\mathcal{K}$  be a minimal blocking set of  $Q(2n + 2, q)$ ,  $n \geq 2$ ,  $|\mathcal{K}| \leq q^{n+1} + q^{n-1}$ . Then there is an  $(n - 2)$ -dimensional space  $\pi$ ,  $\pi \subset Q(2n + 2, q)$ ,  $\pi \cap \mathcal{K} = \emptyset$ , with the following property:  $T_\pi(Q(2n + 2, q)) \cap Q(2n + 2, q) = \pi Q(4, q)$  and  $\mathcal{K}$  consists of all the points of the lines  $M$  on  $p_i$ ,  $p_i \in \pi$ , meeting  $Q(4, q)$  in an ovoid  $\mathcal{O}$ , minus the points  $p_i$  themselves, and  $|\mathcal{K}| = q^{n+1} + q^{n-1}$ .*

For  $q = 3$  we found a characterization using ovoids of  $Q(6, 3)$ .

**Theorem 3** *Let  $\mathcal{K}$  be a minimal blocking set of  $Q(2n + 2, q)$ ,  $n \geq 3$ ,  $|\mathcal{K}| \leq q^{n+1} + q^{n-2}$ . Then there is an  $(n - 3)$ -dimensional space  $\pi$ ,  $\pi \subset Q(2n + 2, 3)$ ,  $\pi \cap \mathcal{K} = \emptyset$ , with the following property:  $T_\pi(Q(2n + 2, 3)) \cap Q(2n + 2, 3) = \pi Q(6, 3)$  and  $\mathcal{K}$  consists of all the points of the lines  $M$  on  $p_i$ ,  $p_i \in \pi$ , meeting  $Q(6, 3)$  in an ovoid  $\mathcal{O}$ , minus the points  $p_i$  themselves, and  $|\mathcal{K}| = q^{n+1} + q^{n-2}$ .*

## References

1. J. Eisfeld, L. Storme, T. Szőnyi, and P. Sziklai, *Covers and blocking sets of classical generalized quadrangles*, Discrete Math., 238(1-3):35–51, 2001.
2. J. De Beule and L. Storme, *The smallest minimal blocking sets of  $Q(6, q)$ ,  $q$  even*, J. Combin. Des., to appear.