Lie algebroid structures on a class of affine bundles

W. Sarlet*, T. Mestdag
Department of Mathematical Physics and Astronomy
Ghent University, Krijgslaan 281, B-9000 Ghent, Belgium
E. Martínez
Departamento de Matemática Aplicada
Universidad de Zaragoza, María de Luna 3, E-50015 Zaragoza, Spain

Keywords: Lie algebroids, affine bundles, exterior calculus.
AMS classification: 37J99, 17B60, 53C15

Abstract. We introduce the notion of a Lie algebroid structure on an affine bundle whose base manifold is fibred over $\mathbb{R}$. It is argued that this is the framework which one needs for coming to a time-dependent generalization of the theory of Lagrangian systems on Lie algebroids. An extensive discussion is given of a way one can think of forms acting on sections of the affine bundle. It is further shown that the affine Lie algebroid structure gives rise to a coboundary operator on such forms. The concept of admissible curves and dynamical systems whose integral curves are admissible, brings an associated affine bundle into the picture, on which one can define in a natural way a prolongation of the original affine Lie algebroid structure.

*email: willy.sarlet@rug.ac.be

1 Introduction

There has been a lot of interest recently in the study of dynamical systems which have a Lie algebroid as carrying space (see e.g. [1, 2, 3, 4, 5, 6]). A Lie algebroid is a vector bundle $\pi : V \to M$, which comes equipped with two operators. To begin with, there is a bracket operation on $Sec(\pi)$, the set of sections of π, which provides it with a real Lie algebra structure . Secondly, there is a linear bundle map $\rho : V \to TM$, called the anchor map, which establishes a Lie algebra homomorphism between $Sec(\pi)$ and the real Lie algebra of vector fields on M and does this in such a way that there is a certain compatibility also with the module structure over $C^\infty(M)$. To be precise, we have

$$[\rho(\sigma), \rho(\eta)] = \rho([\sigma, \eta]) \qquad \text{and} \qquad [\sigma, f\,\eta] = f[\sigma, \eta] + \rho(\sigma)(f)\,\eta,$$

for all $\sigma, \eta \in Sec(\pi)$ and $f \in C^\infty(M)$.

Weinstein's paper on Lagrangian mechanics and groupoids [6] roused new interest into the field of algebroids and groupoids. Weinstein introduces 'Lagrangian systems' on a Lie algebroid by means of a Legendre-type map from V to V^*, associated to a given function L on V. The local coordinate expression of such equations reads:

$$\begin{aligned} \dot{x}^i &= \rho^i_\alpha(x) y^\alpha, \\ \frac{d}{dt}\left(\frac{\partial L}{\partial y^\alpha}\right) &= \rho^i_\alpha \frac{\partial L}{\partial x^i} - C^\gamma_{\alpha\beta} y^\beta \frac{\partial L}{\partial y^\gamma}, \end{aligned} \tag{1}$$

where the x^i are coordinates on M, y^α are fibre coordinates on V and the $C^\gamma_{\alpha\beta}$ are structure functions coming from the Lie algebroid structure. Applications for such model equations can be found e.g. in the theory of systems with symmetries on principal fibre bundles and in rigid body dynamics. Note that, more generally, equations of the form

$$\begin{aligned} \dot{x}^i &= \rho^i_\alpha(x) y^\alpha, \\ \dot{y}^\alpha &= f^\alpha(x, y), \end{aligned} \tag{2}$$

were called "second-order equations on a Lie algebroid" by Weinstein. They are indeed, to some extent, the analogues of second-order dynamics on a tangent bundle. It is clear, however, that these equations truly are second-order differential equations only when the base manifold and the fibres have the same dimension and ρ is injective. We will therefore rather call them 'pseudo-second-order ordinary differential equations', pseudo-SODEs for short. Weinstein also raised the question whether there would be a geometrical way of defining equations of the form (1), much in the line of the geometrical construction of classical Lagrange equations, which makes use of the intrinsic structures living on a tangent bundle.

One of us has recently resolved this issue [5] by introducing a kind of lifted Lie algebroid, where suitable analogues can be introduced of the dilation vector field and the vertical endomorphism on a tangent bundle.

In the present paper, we wish to set the stage for an appropriate generalization of this theory to non-autonomous systems of differential equations. We believe that, for example at the level of pseudo-second-order equations, the right generalization is not just a

matter of allowing the functions ρ^i_α and f^α to depend on time, but rather should produce equations of the form:

$$\begin{aligned} \dot{x}^i &= \rho^i_\alpha(t,x)y^\alpha + \lambda^i(t,x) \\ \dot{y}^\alpha &= f^\alpha(t,x,y) \end{aligned} \tag{3}$$

The reason for this simply is that we wish the structure of the equations to be invariant under time-dependent coordinate transformations. As for Lagrange-type equations, our only concern at the moment is to have an idea of what a time-dependent generalization of (1) should look like. Now, there is a way of developing a kind of formal calculus of variations approach which leads to equations of the form (1), and in which the first set of equations are treated as constraints. We have shown in [7] that if such an approach is adopted when the Lagrangian is allowed to depend on time and the constraints are as in (3), one obtains equations of the form

$$\begin{aligned} \dot{x}^i &= \rho^i_\alpha(t,x)y^\alpha + \lambda^i(t,x), \\ \frac{d}{dt}\left(\frac{\partial L}{\partial y^\alpha}\right) &= \rho^i_\alpha \frac{\partial L}{\partial x^i} - (C^\gamma_{\alpha\beta}y^\beta - C^\gamma_\alpha)\frac{\partial L}{\partial y^\gamma} \end{aligned} \tag{4}$$

where the functions ρ^i_α, λ^i, $C^\gamma_{\alpha\beta}$, C^γ_α satisfy the relations

$$\rho^i_\alpha \frac{\partial \rho^j_\beta}{\partial x^i} - \rho^i_\beta \frac{\partial \rho^j_\alpha}{\partial x^i} = \rho^j_\gamma C^\gamma_{\alpha\beta}, \tag{5}$$

$$\frac{\partial \rho^j_\beta}{\partial t} + \lambda^i \frac{\partial \rho^j_\beta}{\partial x^i} - \rho^i_\beta \frac{\partial \lambda^j}{\partial x^i} = \rho^j_\alpha C^\alpha_\beta. \tag{6}$$

Thus, we want to address the question of explaining the nature of the conditions (5,6), which presumably should again have something to do with a Lie algebroid structure.

Inspired by these analytical considerations, we will introduce the notion of a Lie algebroid structure on an affine bundle $\pi : E \to M$, where the base manifold M in addition is assumed to be fibred over $\mathbb{R}$. The assumption about the additional fibration of M is a very natural one to start the development of this new theory, because the canonical example of an affine Lie algebroid (when the anchor map reduces to the identity) should be the model of time-dependent mechanics, which traditionally is described on the first-jet bundle of a manifold fibred over $\mathbb{R}$ (see e.g. [8]). For the present paper, we will limit ourselves to a number of basic features of such a theory. In particular, we shall show in Section 3 that our defining relations for an affine Lie algebroid are fully consistent with the expectation of being able to develop an exterior differential calculus of sections of the extended dual of this bundle (and its exterior products). We shall further show in Section 4 that vector fields on E, whose integral curves are 'admissible curves' and which in fact model differential equations of the pseudo-SODE type, can be identified in a natural way with special sections of a kind of prolongation of the original affine bundle. This then brings us to a final test, for this paper, of the internal coherence of the newly defined structures: we will verify in Section 5 whether the prolongation of a Lie algebroid, as constructed in [5] for the vector bundle situation, carries over to the present more general situation.

The final section lists a number of other topics of interest, which will be the subject of forthcoming publications. One of our objectives is to arrive at an intrinsic geometrical construction of the time-dependent Lagrangian equations of type (4). For the time being, however, the conditions (5,6) merely serve as benchmarks, to be met by our model of an affine Lie algebroid.

The basic ingredients for our theory are an affine bundle $\pi : E \to M$, where the base manifold M is further fibred over $\mathbb{R}$. In Section 2, we define the concept of a Lie algebroid on π. In Section 3, we show that the axioms for such a Lie algebroid structure give rise to a consistent development of an exterior calculus on sections of the extended dual of E. In Section 4, we discuss a special class of curves on E, which are said to be admissible by the anchor map and we look into the concept of dynamical systems whose integral curves all belong to this special class. In Section 5, we define the prolongation $\pi_1 : J^1_\lambda E \to E$ of $\pi : E \to M$ and show that it inherits the Lie algebroid structure from π.

2 Affine Lie algebroids

Let M be an $(n+1)$-dimensional smooth manifold, which is fibred over $\mathbb{R}$, $\tau : M \to \mathbb{R}$. We denote the first jet bundle of τ by $\tau^0_1 : J^1M \to M$. It is an affine bundle modelled on the bundle of tangent vectors to M which are vertical with respect to τ; this vector bundle will be denoted by $\overline{\tau}^0_1 : VM \to M$. To fix notations further, if $\pi : E \to M$ is an affine bundle and $\overline{\pi} : V \to M$ its associated vector bundle, sections of π will be denoted by ordinary Greek characters, whereas boldface Greek type will be used for sections of $\overline{\pi}$.

Definition 1. *An affine Lie algebroid over M is an affine bundle $\pi : E \to M$, with the following properties:*

1. *$Sec(\overline{\pi})$, the set of sections of the vector bundle $\overline{\pi} : V \to M$ on which E is modelled, is equipped with a skew-symmetric and bilinear (over $\mathbb{R}$) bracket $[\cdot,\cdot]$;*

2. *the affine space $Sec(\pi)$ acts by derivations on the real algebra $Sec(\overline{\pi})$, that is to say, if the same bracket notation $[\zeta, \boldsymbol{\sigma}]$ is used to denote the way $\zeta \in Sec(\pi)$ acts on $\boldsymbol{\sigma} \in Sec(\overline{\pi})$, we have $[\zeta, \boldsymbol{\sigma}] \in Sec(\overline{\pi})$ and*

$$[\zeta, \boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2] = [\zeta, \boldsymbol{\sigma}_1] + [\zeta, \boldsymbol{\sigma}_2], \qquad [\zeta + \boldsymbol{\sigma}, \boldsymbol{\eta}] = [\zeta, \boldsymbol{\eta}] + [\boldsymbol{\sigma}, \boldsymbol{\eta}], \tag{7}$$

$$[\zeta, [\boldsymbol{\sigma}, \boldsymbol{\eta}]] = [[\zeta, \boldsymbol{\sigma}], \boldsymbol{\eta}] + [\boldsymbol{\sigma}, [\zeta, \boldsymbol{\eta}]]; \tag{8}$$

3. *there exists an affine bundle map $\lambda : E \to J^1M$ (over the identity on M), with corresponding vector bundle homomorphism $\rho : V \to VM$, such that the following compatibility condition holds for all $f \in C^\infty(M)$,*

$$[\zeta, f\boldsymbol{\sigma}] = f\,[\zeta, \boldsymbol{\sigma}] + \lambda(\zeta)(f)\boldsymbol{\sigma}. \tag{9}$$

Both the affine map $\lambda : E \to J^1M$ and its linear part $\rho : V \to VM$ will be called anchor maps.

Note that we make no notational distinction between, on the one hand, the affine and linear anchor maps, regarded as maps between total spaces of bundles, and their action on sections of bundles on the other hand. Needless to say, for the interpretation of the bracket in the left-hand side of (12), both $\lambda(\zeta)$ and $\rho(\boldsymbol{\sigma})$ are regarded as vector fields on M.

Let us derive some further properties which follow from this definition. First of all, if we replace ζ in (8) by $\zeta + \boldsymbol{\xi}$, for an arbitrary $\boldsymbol{\xi} \in Sec(\overline{\pi})$, it follows that $\forall \boldsymbol{\xi}, \boldsymbol{\sigma}, \boldsymbol{\eta} \in Sec(\overline{\pi})$:

$$[\boldsymbol{\xi}, [\boldsymbol{\sigma}, \boldsymbol{\eta}]] = [[\boldsymbol{\xi}, \boldsymbol{\sigma}], \boldsymbol{\eta}] + [\boldsymbol{\sigma}, [\boldsymbol{\xi}, \boldsymbol{\eta}]]. \tag{10}$$

This means, in view of the first hypothesis, that the bracket on $Sec(\overline{\pi})$ actually provides $Sec(\overline{\pi})$ with a real Lie algebra structure. Secondly, making the same substitution for ζ in (9), recalling that $\lambda(\zeta + \boldsymbol{\xi}) = \lambda(\zeta) + \rho(\boldsymbol{\xi})$, it follows that

$$[\boldsymbol{\xi}, f\boldsymbol{\sigma}] = f\,[\boldsymbol{\xi}, \boldsymbol{\sigma}] + \rho(\boldsymbol{\xi})(f)\boldsymbol{\sigma}. \tag{11}$$

This means that the linear anchor map $\rho : V \to VM$ defines a Lie algebra homomorphism from $Sec(\overline{\pi})$ into the real Lie algebra of vertical vector fields on M, and that we have a classical Lie algebroid structure on the vector bundle $\overline{\pi} : V \to M$ (although its image cannot reach the whole of TM). Thirdly, replacing $\boldsymbol{\eta}$ by $f\boldsymbol{\eta}$ in (8) and making use of (9) and (11), one obtains the additional compatibility property

$$[\lambda(\zeta), \rho(\boldsymbol{\sigma})] = \rho([\zeta, \boldsymbol{\sigma}]), \tag{12}$$

from which it further follows that

$$[\rho(\boldsymbol{\xi}), \rho(\boldsymbol{\sigma})] = \rho\left([\boldsymbol{\xi}, \boldsymbol{\sigma}]\right). \tag{13}$$

Remark: for an alternative and equivalent definition of an affine Lie algebroid, we could impose first the Lie algebra structure (10) of the bracket on $Sec(\overline{\pi})$, together with the compatibility condition (11) for the anchor map ρ, and subsequently require that the properties (7-9) hold true for at least one $\zeta \in Sec(\pi)$ and for an affine map $\lambda : E \to J^1M$ whose linear part is ρ. It then follows that such properties hold for all ζ.

We can now further extend the bracket operation to $Sec(\pi)$, as follows.

Definition 2. (i) *For $\boldsymbol{\sigma} \in Sec(\overline{\pi})$ and $\zeta \in Sec(\pi)$, we put $[\boldsymbol{\sigma}, \zeta] = -[\zeta, \boldsymbol{\sigma}]$.*

(ii) *For every two sections $\zeta_1, \zeta_2 \in Sec(\pi)$ with $\boldsymbol{\zeta}_{12} = \zeta_2 - \zeta_1$: $[\zeta_1, \zeta_2] = [\zeta_1, \boldsymbol{\zeta}_{12}]$.*

Observe that the extended bracket in (ii) is a map from $Sec(\pi) \times Sec(\pi)$ to $Sec(\overline{\pi})$. As we will show below, it has Lie algebra type properties, which could justify talking about an "affine Lie algebra structure", were it not that this term is in use already in the literature, with an entirely different meaning. The extended bracket also has Lie algebroid type properties with respect to the anchor maps λ and ρ.

Proposition 1. *The bracket* $[\cdot,\cdot] : Sec(\pi)\times Sec(\pi) \to Sec(\overline{\pi})$, *has the following properties:*

$$[\zeta_1, \zeta_2 + \boldsymbol{\sigma}] = [\zeta_1, \zeta_2] + [\zeta_1, \boldsymbol{\sigma}], \tag{14}$$
$$[\zeta_1, \zeta_2] = -[\zeta_2, \zeta_1], \tag{15}$$
$$[[\zeta_1, \zeta_2], \zeta_3] + [[\zeta_2, \zeta_3], \zeta_1] + [[\zeta_3, \zeta_1], \zeta_2] = \mathbf{0}, \tag{16}$$
$$\rho([\zeta_1, \zeta_2]) = [\lambda(\zeta_1), \lambda(\zeta_2)]. \tag{17}$$

Proof: The first property follows immediately from the definition and (7). Next, we have $[\zeta_2, \zeta_1] = [\zeta_2, \boldsymbol{\zeta}_{21}] = -[\zeta_1 + \boldsymbol{\zeta}_{12}, \boldsymbol{\zeta}_{12}] = -[\zeta_1, \zeta_2]$. For the Jacobi identity, using a simple summation sign to indicate the cyclic sum over the three sections in each summand, we have

$$\sum [[\zeta_1, \zeta_2], \zeta_3] = \sum [[\zeta_1, \zeta_2], \zeta_2 + \boldsymbol{\zeta}_{23}] = \sum [[\zeta_1, \zeta_2], \boldsymbol{\zeta}_{23}],$$

in view of the linearity properties and the skew-symmetry of the bracket. Substituting subsequently $\zeta_1 + \boldsymbol{\zeta}_{12}$ for ζ_2, we obtain

$$\sum [[\zeta_1, \zeta_2], \zeta_3] = \sum [[\zeta_1, \boldsymbol{\zeta}_{12}], \boldsymbol{\zeta}_{23}],$$

which is zero in view of (8). Finally, the compatibility property (17) easily follows in the same way from the definition of the extended bracket and (12). □

To understand what an affine Lie algebroid structure means in coordinates, let us coordinatize E in the usual way, as follows: t denotes the coordinate on $\mathbb{R}$; $(x^i)_{1\le i\le n}$ are fibre coordinates on M; we further choose a local section e_0 of π to play the role of zero section and a local basis $(\boldsymbol{e}_\alpha)_{1\le\alpha\le k}$ for $Sec(\overline{\pi})$. Then, if e is a point in the fibre E_m over $m \in M$, it can be written in the form: $e = e_0(m) + y^\alpha \boldsymbol{e}_\alpha(m)$; (t, x^i, y^α) are coordinates of e ((t, x^i) being the coordinates of m).

We have

$$[\boldsymbol{e}_\alpha, \boldsymbol{e}_\beta] = C^\gamma_{\alpha\beta}(t,x)\boldsymbol{e}_\gamma, \qquad [e_0, \boldsymbol{e}_\alpha] = C^\beta_\alpha(t,x)\boldsymbol{e}_\beta, \tag{18}$$

for some *structure functions* $C^\gamma_{\alpha\beta} = -C^\gamma_{\beta\alpha}$ and C^β_α on M. The affine map λ and its linear part ρ are fully determined by

$$\lambda(e_0) = \frac{\partial}{\partial t} + \lambda^i(t,x)\frac{\partial}{\partial x^i}, \qquad \rho(\boldsymbol{e}_\alpha) = \rho^i_\alpha(t,x)\frac{\partial}{\partial x^i}. \tag{19}$$

The further characterization of the Lie algebroid structure now has the following coordinate translation. The derivation property (8) and the resulting Jacobi identity (10) mean that we have:

$$\frac{\partial C^\mu_{\alpha\beta}}{\partial t} + \lambda^i\frac{\partial C^\mu_{\alpha\beta}}{\partial x^i} + C^\gamma_{\alpha\beta}C^\mu_\gamma = C^\mu_{\alpha\gamma}C^\gamma_\beta - C^\mu_{\beta\gamma}C^\gamma_\alpha + \rho^i_\alpha\frac{\partial C^\mu_\beta}{\partial x^i} - \rho^i_\beta\frac{\partial C^\mu_\alpha}{\partial x^i}, \tag{20}$$
$$\sum_{\alpha,\beta,\gamma}\left(\rho^i_\alpha\frac{\partial C^\mu_{\beta\gamma}}{\partial x^i} + C^\mu_{\alpha\nu}C^\nu_{\beta\gamma}\right) = 0, \tag{21}$$

where the summation this time refers to a cyclic sum over α, β, γ and also the compatibility conditions (9),(11) have been invoked. Finally, the properties (12) and (13), for which

it is sufficient to express that $[\lambda(e_0), \rho(\boldsymbol{e}_\alpha)] = \rho([e_0, \boldsymbol{e}_\alpha])$ and $[\rho(\boldsymbol{e}_\alpha), \rho(\boldsymbol{e}_\beta)] = \rho([\boldsymbol{e}_\alpha, \boldsymbol{e}_\beta])$, require that

$$\frac{\partial \rho^j_\beta}{\partial t} + \lambda^i \frac{\partial \rho^j_\beta}{\partial x^i} - \rho^i_\beta \frac{\partial \lambda^j}{\partial x^i} = C^\alpha_\beta \rho^j_\alpha, \tag{22}$$

$$\rho^i_\alpha \frac{\partial \rho^j_\beta}{\partial x^i} - \rho^i_\beta \frac{\partial \rho^j_\alpha}{\partial x^i} = C^\gamma_{\alpha\beta} \rho^j_\gamma. \tag{23}$$

These are precisely the relations (5) and (6) we encountered in the Introduction, in the context of Lagrangian equations of type (4).

It is of some interest to look at the way the various structure and anchor map functions transform under coordinate transformations. There are two distinct levels in making a change of coordinates on E, which we will describe separately. Firstly, we could choose a different (local) zero section $\overline{e}_0$ and a different local basis $\overline{\boldsymbol{e}}_\beta$ for $Sec(\overline{\pi})$: say that $\boldsymbol{e}_\alpha = A^\beta_\alpha \overline{\boldsymbol{e}}_\beta$ and $e_0 = \overline{e}_0 + B^\alpha \overline{\boldsymbol{e}}_\alpha$. This amounts to making an affine change of coordinates in the fibres of the form: $\overline{y}^\alpha = A^\alpha_\beta(t,x) y^\beta + B^\alpha(t,x)$. Putting $[\overline{\boldsymbol{e}}_\alpha, \overline{\boldsymbol{e}}_\beta] = \overline{C}^\gamma_{\alpha\beta} \overline{\boldsymbol{e}}_\gamma$, $[\overline{e}_0, \overline{\boldsymbol{e}}_\alpha] = \overline{C}^\beta_\alpha \overline{\boldsymbol{e}}_\beta$, and also $\lambda(\overline{e}_0) = \frac{\partial}{\partial t} + \overline{\lambda}^i \frac{\partial}{\partial x^i}$, $\rho(\overline{\boldsymbol{e}}_\alpha) = \overline{\rho}^j_\alpha \frac{\partial}{\partial x^j}$, one can verify that the following transformation rules apply:

$$\rho^i_\alpha = A^\beta_\alpha \overline{\rho}^i_\beta, \qquad \overline{\lambda}^i = \lambda^i - B^\alpha \overline{\rho}^i_\alpha,$$

and further

$$\begin{aligned} C^\gamma_{\alpha\beta} A^\mu_\gamma &= \overline{C}^\mu_{\gamma\nu} A^\gamma_\alpha A^\nu_\beta + \rho^i_\alpha \frac{\partial A^\mu_\beta}{\partial x^i} - \rho^i_\beta \frac{\partial A^\mu_\alpha}{\partial x^i}, \\ C^\gamma_\beta A^\alpha_\gamma &= \overline{C}^\alpha_\mu A^\mu_\beta + \overline{C}^\alpha_{\gamma\mu} B^\gamma A^\mu_\beta + \frac{\partial A^\alpha_\beta}{\partial t} + \lambda^i \frac{\partial A^\alpha_\beta}{\partial x^i} - \rho^i_\beta \frac{\partial B^\alpha}{\partial x^i}. \end{aligned}$$

At a different level, one can make a change of coordinates on M, of the form: $t' = t$, $x'^i = x'^i(t,x)$. This has an effect on the anchor map functions of the form:

$$\rho'^j_\alpha = \rho^i_\alpha \frac{\partial x'^j}{\partial x^i}, \qquad \lambda'^j = \frac{\partial x'^j}{\partial t} + \lambda^i \frac{\partial x'^j}{\partial x^i}.$$

A general change of adapted coordinates is of course a composition of the two steps described above.

3 Exterior calculus on an affine Lie algebroid

We first recall some features of the by now standard theory of Lie algebroids on a vector bundle (see [9]). Considering sections of exterior powers of the dual bundle, one gets a notion of forms on sections of the vector bundle, on which an exterior derivative can be defined which involves the Lie algebroid bracket and the anchor map. It then turns out that the Jacobi identity of the Lie algebroid bracket and the compatibility with the bracket of vector fields via the anchor map are exactly the conditions for this exterior derivative to have the co-boundary property $d^2 = 0$ (see also [10, 11, 5]). In our opinion,

such a feature in itself gives a strong indication that the generalization from Lie algebra to Lie algebroid is indeed a meaningful step. We shall therefore investigate in this section whether a similar support can be detected for our extension to Lie algebroids on affine bundles.

The extended dual of the affine space E_m is the space of real valued affine functions on E_m and will be denoted by $E_m^\dagger$. The union of these spaces over all points $m \in M$ gives us a bundle $\pi^\dagger : E^\dagger \to M$ say. Although this is in fact a vector bundle, we are interested in the action of its sections (and sections of its exterior powers) on sections of the affine bundle π. This brings some subtleties into the picture which need to be investigated in sufficient detail. We will write $\overline{\pi}^* : V^* \to M$ for the dual bundle of $\overline{\pi}$ and also use boldface type for its sections (and the sections of its exterior powers). Now, to begin with, if θ is a section of $\pi^\dagger$ and $\zeta \in Sec(\pi)$, $\theta(\zeta)$ is a function on M defined by $\theta(\zeta)(m) = \theta_m(\zeta_m)$. θ_m being an affine function on E_m, there exists an associated element $\boldsymbol{\theta}_m \in V_m^*$ such that $\forall e_m \in E_m,\ \boldsymbol{\sigma}_m \in V_m$, we have $\theta_m(e_m + \boldsymbol{\sigma}_m) = \theta_m(e_m) + \boldsymbol{\theta}_m(\boldsymbol{\sigma}_m)$. Expressed in slightly different terms and now at the level of sections again, a $\theta \in Sec(\pi^\dagger)$ is such that there exist a $\theta_0 \in Sec(\pi^\dagger)$ and a $\boldsymbol{\theta} \in Sec(\overline{\pi}^*)$, such that for all $\zeta \in Sec(\pi)$:

$$\theta(\zeta) = \theta_0(\zeta_0) + \boldsymbol{\theta}(\boldsymbol{\zeta}), \tag{24}$$

where ζ_0 is any section and then $\zeta = \zeta_0 + \boldsymbol{\zeta}$. The two composing elements θ_0 (which in fact is simply θ itself here) and $\boldsymbol{\theta}$ do not depend on the choice of ζ_0. With sections of $\pi^\dagger$ as our notion of 1-forms on $Sec(\pi)$, there is of course no linearity with respect to multiplication by functions on M. We can now come in a similar way to the following concept of k-forms on $Sec(\pi)$ (thereby taking for granted that the meaning of a k-form on a vector bundle such as $Sec(\overline{\pi})$ is known).

Definition 3. *A k-form on the affine bundle $Sec(\pi)$ ($k \geq 1$) is a map $\omega : Sec(\pi) \times \cdots \times Sec(\pi) \to C^\infty(M)$, for which there exists a k-form $\boldsymbol{\omega}$ on the associated vector bundle $Sec(\overline{\pi})$ and a map $\omega_0 : Sec(\pi) \times Sec(\overline{\pi}) \times \cdots \times Sec(\overline{\pi}) \to C^\infty(M)$ with the following properties:*

1. *ω_0 is skew-symmetric and $C^\infty(M)$-linear in its $k-1$ vector arguments;*
2. *$\forall \zeta \in Sec(\pi)$ and $\forall \boldsymbol{\sigma}, \boldsymbol{\zeta}_j \in Sec(\overline{\pi})$, we have*

$$\omega_0(\zeta + \boldsymbol{\sigma}, \boldsymbol{\zeta}_1, \ldots, \boldsymbol{\zeta}_{k-1}) = \omega_0(\zeta, \boldsymbol{\zeta}_1, \ldots, \boldsymbol{\zeta}_{k-1}) + \boldsymbol{\omega}(\boldsymbol{\sigma}, \boldsymbol{\zeta}_1, \ldots, \boldsymbol{\zeta}_{k-1}); \tag{25}$$

3. *$\forall \zeta_i \in Sec(\pi)$, if we choose an arbitrary $\zeta_0 \in Sec(\pi)$ and put $\zeta_i = \zeta_0 + \boldsymbol{\zeta}_i$, we have*

$$\omega(\zeta_1, \ldots, \zeta_k) = \sum_{i=1}^{k} (-1)^{i-1} \omega_0(\zeta_0, \boldsymbol{\zeta}_1, \ldots, \hat{\boldsymbol{\zeta}}_i, \ldots, \boldsymbol{\zeta}_k) + \boldsymbol{\omega}(\boldsymbol{\zeta}_1, \ldots, \boldsymbol{\zeta}_k). \tag{26}$$

There are a number of properties to be checked to make sure that this definition makes sense. First of all, one can verify that with two different choices of a reference section, ζ_0

and ζ_0' for example, related through $\zeta_0 = \zeta_0' + \boldsymbol{\sigma}$, it follows from the second requirement that

$$\sum_{i=1}^{k}(-1)^{i-1}\omega_0(\zeta_0, \boldsymbol{\zeta}_1, \ldots, \hat{\boldsymbol{\zeta}}_i, \ldots, \boldsymbol{\zeta}_k) + \boldsymbol{\omega}(\boldsymbol{\zeta}_1, \ldots, \boldsymbol{\zeta}_k)$$
$$= \sum_{i=1}^{k}(-1)^{i-1}\omega_0(\zeta_0', \boldsymbol{\zeta}_1', \ldots, \hat{\boldsymbol{\zeta}}_i', \ldots, \boldsymbol{\zeta}_k') + \boldsymbol{\omega}(\boldsymbol{\zeta}_1', \ldots, \boldsymbol{\zeta}_k').$$

Secondly, the two elements ω_0 and $\boldsymbol{\omega}$ which make up ω are unique. Indeed, assuming there would be a second couple ω_0' and $\boldsymbol{\omega}'$ making up the same ω, it follows by choosing $\zeta_0 = \zeta_1$ (such that $\boldsymbol{\zeta}_1 = 0$) that $\omega_0 = \omega_0'$, after which it is clear that also $\boldsymbol{\omega} = \boldsymbol{\omega}'$. Note finally that the definition implies that ω itself is skew-symmetric in all its arguments.

The set of forms on $Sec(\pi)$, which we will denote by $\bigwedge(\pi^\dagger)$, is a module over the ring $C^\infty(M)$, which also constitutes the set of 0-forms. The wedge product of two forms is defined in the usual way. By way of example, if α and β are 1-forms, we have

$$(\alpha \wedge \beta)(\zeta_1, \zeta_2) = \alpha(\zeta_1)\beta(\zeta_2) - \alpha(\zeta_2)\beta(\zeta_1),$$

which for every choice of a reference section ζ_0 gives rise to:

$$\begin{aligned}(\alpha \wedge \beta)(\zeta_1, \zeta_2) &= \alpha(\zeta_0)\boldsymbol{\beta}(\boldsymbol{\zeta}_2 - \boldsymbol{\zeta}_1) - \beta(\zeta_0)\boldsymbol{\alpha}(\boldsymbol{\zeta}_2 - \boldsymbol{\zeta}_1) \\ &\quad + (\boldsymbol{\alpha} \wedge \boldsymbol{\beta})(\boldsymbol{\zeta}_1, \boldsymbol{\zeta}_2). \end{aligned} \tag{27}$$

It follows that $\boldsymbol{\alpha} \wedge \boldsymbol{\beta}$ is the 2-form on $Sec(\overline{\pi})$ corresponding to $\alpha \wedge \beta$, and

$$(\alpha \wedge \beta)_0(\zeta, \boldsymbol{\sigma}) = \alpha(\zeta)\boldsymbol{\beta}(\boldsymbol{\sigma}) - \beta(\zeta)\boldsymbol{\alpha}(\boldsymbol{\sigma}). \tag{28}$$

Similarly, for the wedge product of three 1-forms, we have

$$(\alpha \wedge \beta \wedge \gamma)_0(\zeta, \boldsymbol{\zeta}_1, \boldsymbol{\zeta}_2) = \Big(\alpha(\zeta)(\boldsymbol{\beta} \wedge \boldsymbol{\gamma}) + \beta(\zeta)(\boldsymbol{\gamma} \wedge \boldsymbol{\alpha}) + \gamma(\zeta)(\boldsymbol{\alpha} \wedge \boldsymbol{\beta})\Big)(\boldsymbol{\zeta}_1, \boldsymbol{\zeta}_2). \tag{29}$$

These examples suggest to formalize the representation of k-forms a bit further. As a preliminary remark, it may sometimes be of interest to extend the interpretation of the operator ω_0 in such a way that its single affine section argument need not necessarily be the first. This can simply be achieved by declaring ω_0 to be skew-symmetric in all its arguments (but still $C^\infty(M)$-linear in its vector arguments only). More importantly, we shall take the sum of ω_0-terms in (26) to define another operator, denoted by ω^0, as follows:

$$\omega^0(\zeta_1, \ldots, \zeta_k) = \sum_{i=1}^{k}(-1)^{i-1}\omega_0(\zeta_0, \boldsymbol{\zeta}_1, \ldots, \hat{\boldsymbol{\zeta}}_i, \ldots, \boldsymbol{\zeta}_k) \;=\; \sum_{i=1}^{k}\omega_0(\boldsymbol{\zeta}_1, \ldots, \underset{i}{\check{\zeta}_0}, \ldots, \boldsymbol{\zeta}_k), \tag{30}$$

where the second expression takes the above remark into account and the symbol $\underset{i}{\check{\zeta}_0}$ then indicates that ζ_0 has been inserted in the i-th argument. The other important new convention we will adopt is to regard $\boldsymbol{\omega}$ also as acting on affine sections:

$$\boldsymbol{\omega}(\zeta_1, \ldots, \zeta_k) = \boldsymbol{\omega}(\boldsymbol{\zeta}_1, \ldots, \boldsymbol{\zeta}_k). \tag{31}$$

This way, we can formally write

$$\omega = \omega^0 + \boldsymbol{\omega}, \tag{32}$$

whereby it is to be understood that the two composing terms ω^0 and $\boldsymbol{\omega}$ are not k-forms on $Sec(\pi)$ by themselves. In fact, to compute their value when acting on k sections ζ_i, a reference section ζ_0 has to be chosen, but as argued above, the value of the sum $\omega^0 + \boldsymbol{\omega}$ in the end does not depend on that choice.

The rather formal looking decomposition (32) now greatly facilitates the representation of wedge products and will make the general coordinate representation of a form more transparant. For example, the result (29) means that

$$(\alpha \wedge \beta \wedge \gamma)_0 = \alpha^0 \otimes (\boldsymbol{\beta} \wedge \boldsymbol{\gamma}) + \beta^0 \otimes (\boldsymbol{\gamma} \wedge \boldsymbol{\alpha}) + \gamma^0 \otimes (\boldsymbol{\alpha} \wedge \boldsymbol{\beta}), \tag{33}$$

which then implies from (30) that

$$(\alpha \wedge \beta \wedge \gamma)^0 = \alpha^0 \wedge \boldsymbol{\beta} \wedge \boldsymbol{\gamma} + \boldsymbol{\alpha} \wedge \beta^0 \wedge \boldsymbol{\gamma} + \boldsymbol{\alpha} \wedge \boldsymbol{\beta} \wedge \gamma^0, \tag{34}$$

as expected. More generally, it follows directly from the defining formula for wedge products that for $\omega = \omega^0 + \boldsymbol{\omega}$ and $\rho = \rho^0 + \boldsymbol{\rho}$:

$$\omega \wedge \rho = \omega^0 \wedge \rho^0 + \omega^0 \wedge \boldsymbol{\rho} + \boldsymbol{\omega} \wedge \rho^0 + \boldsymbol{\omega} \wedge \boldsymbol{\rho}, \tag{35}$$

where the sum of the first three terms is $(\omega \wedge \rho)^0$.

Suppose that, for a coordinatization of E, we have chosen a zero section e_0 and a local basis of vector sections $\boldsymbol{e}_\alpha$. Denote by $\{\boldsymbol{e}^\beta\}$ the dual basis for $Sec(\overline{\pi}^*)$. There exists a global section of $\pi^\dagger$ which for each m selects in $E^\dagger_m$ the constant function 1. We will call it e^0. The local zero section e_0 of $Sec(\pi)$ can now play the role of the reference section ζ_0 in our general considerations. Writing $\zeta = e_0 + \zeta^\alpha \boldsymbol{e}_\alpha$ for an arbitrary section ζ, we have for each 1-form θ: $\theta(\zeta) = \theta(e_0) + \zeta^\alpha \boldsymbol{\theta}(\boldsymbol{e}_\alpha)$. Putting $\boldsymbol{\theta}(\boldsymbol{e}_\alpha) = \theta_\alpha$ and $\theta(e_0) = \theta_0$, we see that θ has the local representation

$$\theta = \theta_0 e^0 + \theta_\alpha \boldsymbol{e}^\alpha, \tag{36}$$

where, in agreement with the general decomposition (32), $\boldsymbol{e}^\alpha$ has to be regarded now as acting on $Sec(\pi)$ and $\theta^0 = \theta_0 e^0$. To be precise, putting

$$e_\alpha = e_0 + \boldsymbol{e}_\alpha, \tag{37}$$

the action of $\boldsymbol{e}^\beta$ on affine sections, which can be given a meaning only after introducing a reference section, is determined by:

$$\boldsymbol{e}^\beta(e_0) = 0, \qquad \boldsymbol{e}^\beta(e_\alpha) = \delta^\beta_\alpha. \tag{38}$$

There is another slight abuse of notation in (36) since θ_0 could have a double meaning: in (36) it represents a local function on M, whereas it also could refer to the operator introduced in (24) and more generally in Definition 3. We will, however, seldom use the notation θ_0 in the latter sense when dealing with coordinate calculations, so that the meaning will always be clear from the context.

Let us now see how all these notations fit together when we start wedging 1-forms. For two 1-forms $\alpha = \alpha^0 + \boldsymbol{\alpha}$ and $\beta = \beta^0 + \boldsymbol{\beta}$ we find, for example from (28) and (30), that

$$(\alpha \wedge \beta)^0 = \alpha^0 \wedge \boldsymbol{\beta} - \beta^0 \wedge \boldsymbol{\alpha}. \tag{39}$$

This is in agreement with the general formula (35) since obviously $\alpha^0 \wedge \beta^0 = 0$. Expressing α and β with respect to the basis $(e^0, \boldsymbol{e}^\alpha)$, we find

$$\alpha \wedge \beta = (\alpha_0\beta_\gamma - \beta_0\alpha_\gamma)\, e^0 \wedge \boldsymbol{e}^\gamma + \tfrac{1}{2}(\alpha_\gamma\beta_\delta - \alpha_\delta\beta_\gamma)\, \boldsymbol{e}^\gamma \wedge \boldsymbol{e}^\delta. \tag{40}$$

Similarly, for the wedge product of three 1-forms with local representations of the form (36), we obtain

$$\begin{aligned} \alpha \wedge \beta \wedge \gamma \;\; &= \;\; \tfrac{1}{2}\Big(\alpha_0(\beta_\mu\gamma_\nu - \beta_\nu\gamma_\mu) + \beta_0(\gamma_\mu\alpha_\nu - \gamma_\nu\alpha_\mu) \\ &\quad + \gamma_0(\alpha_\mu\beta_\nu - \alpha_\nu\beta_\mu)\Big) e^0 \wedge \boldsymbol{e}^\mu \wedge \boldsymbol{e}^\nu + \boldsymbol{\alpha} \wedge \boldsymbol{\beta} \wedge \boldsymbol{\gamma}. \end{aligned} \tag{41}$$

It should now be clear without going into any further detail that a general k-form on $Sec(\pi)$ locally has the following representation,

$$\omega = \frac{1}{(k-1)!}\, \omega_{0\mu_1\cdots\mu_{k-1}}\, e^0 \wedge \boldsymbol{e}^{\mu_1} \wedge \cdots \wedge \boldsymbol{e}^{\mu_{k-1}} + \frac{1}{k!}\, \omega_{\mu_1\cdots\mu_k}\, \boldsymbol{e}^{\mu_1} \wedge \cdots \wedge \boldsymbol{e}^{\mu_k}, \tag{42}$$

where the coefficients are functions on M, which are skew-symmetric in all their indices (including the zero for the first term); we have

$$\omega_{0\mu_1\cdots\mu_{k-1}} \;\; = \;\; \omega(e_0, e_{\mu_1}, \ldots, e_{\mu_{k-1}}), \tag{43}$$

$$\omega_{\mu_1\cdots\mu_k} \;\; = \;\; \omega(e_{\mu_1}, \ldots, e_{\mu_k}) - \sum_{i=1}^{k} \omega_{\mu_1\cdots\overset{\check{0}}{i}\cdots\mu_k}, \tag{44}$$

where $\overset{\check{0}}{i}$ again means that the index μ_i has been replaced by 0.

Before arriving at our main goal, the development of an exterior calculus on forms, we will recall a few generalities about derivations. Derivations on $\bigwedge(\pi^\dagger)$ are defined in the usual way. Following the standard work of Frölicher and Nijenhuis [12], one easily shows that derivations are local operators and that they are completely determined by their action on functions and 1-forms. The commutator of two derivations D_i, of degree r_i say, is again a derivation, of degree $r_1 + r_2$, defined by

$$[D_1, D_2] = D_1 \circ D_2 - (-1)^{r_1 r_2} D_2 \circ D_1. \tag{45}$$

Perhaps the simplest type of derivation is contraction with a section.

Definition 4. *For $\omega \in \bigwedge^k(\pi^\dagger)$ and $\zeta \in Sec(\pi)$, $i_\zeta\omega \in \bigwedge^{k-1}(\pi^\dagger)$ is defined by*

$$i_\zeta\omega(\zeta_1, \ldots, \zeta_{k-1}) = \omega(\zeta, \zeta_1, \ldots \zeta_{k-1}). \tag{46}$$

The proof that this is a derivation of degree -1 is standard and does not depend on the peculiarities of our present theory. But perhaps we have to convince ourselves in the first place that $i_\zeta\omega$ is indeed a form in the sense of Definition 3.

Proposition 2. *$i_\zeta\omega$ is a $(k-1)$-form which, in the sense of the general defining relation (26), is determined by an operator $(i_\zeta\omega)_0$ and a k-form $\mathbf{i}_{\boldsymbol{\zeta}}\boldsymbol{\omega}$ on $Sec(\overline{\pi})$, defined as follows: for all $\boldsymbol{\zeta}_i \in Sec(\overline{\pi})$, $\zeta_0 \in Sec(\pi)$,*

$$(i_\zeta\omega)_0(\zeta_0, \boldsymbol{\zeta}_2, \ldots \boldsymbol{\zeta}_{k-1}) = -\omega_0(\zeta_0, \boldsymbol{\zeta}, \boldsymbol{\zeta}_2, \ldots \boldsymbol{\zeta}_{k-1}), \quad \textit{where} \quad \boldsymbol{\zeta} = \zeta - \zeta_0, \tag{47}$$

$$\mathbf{i}_{\boldsymbol{\zeta}}\boldsymbol{\omega}(\boldsymbol{\zeta}_2, \ldots, \boldsymbol{\zeta}_k) = \omega_0(\zeta, \boldsymbol{\zeta}_2, \ldots, \boldsymbol{\zeta}_k). \tag{48}$$

We further have the property (with $\boldsymbol{\zeta}_1 = \boldsymbol{\zeta}$):

$$i_\zeta\omega(\zeta_2, \ldots, \zeta_k) = \sum_{i=1}^{k} (-1)^{i-1} \, (i_\zeta\omega_0) \, (\boldsymbol{\zeta}_1, \ldots, \hat{\boldsymbol{\zeta}}_i, \ldots, \boldsymbol{\zeta}_k). \tag{49}$$

Proof: A direct computation, using (46) and (26), gives

$$\begin{aligned} i_\zeta\omega(\zeta_2, \ldots, \zeta_k) &= \omega(\zeta, \zeta_2, \ldots, \zeta_k) \\ &= \omega_0(\zeta_0, \boldsymbol{\zeta}_2, \ldots, \boldsymbol{\zeta}_k) + \sum_{j=1}^{k-1} (-1)^j \omega_0(\zeta_0, \boldsymbol{\zeta}, \boldsymbol{\zeta}_2, \ldots, \hat{\boldsymbol{\zeta}}_j, \ldots, \boldsymbol{\zeta}_k) + \boldsymbol{\omega}(\boldsymbol{\zeta}, \boldsymbol{\zeta}_2, \ldots, \boldsymbol{\zeta}_k) \\ &= \sum_{j=1}^{k-1} (-1)^j \omega_0(\zeta_0, \boldsymbol{\zeta}, \boldsymbol{\zeta}_2, \ldots, \hat{\boldsymbol{\zeta}}_j, \ldots, \boldsymbol{\zeta}_k) + \omega_0(\zeta, \boldsymbol{\zeta}_2, \ldots, \boldsymbol{\zeta}_k), \end{aligned}$$

from which we are led to introduce $(i_\zeta\omega)_0$ and $\mathbf{i}_{\boldsymbol{\zeta}}\boldsymbol{\omega}$ as in (47) and (48). It is then straightforward to verify that these two operators are linked by a property of type (25), so the first statement follows. Observe that, with an obvious meaning for contraction of the operator ω_0 with ζ, we can write: $\mathbf{i}_{\boldsymbol{\zeta}}\boldsymbol{\omega} = i_\zeta\omega_0$. The somewhat peculiar feature of the additional property is that $i_\zeta\omega$ can be completely computed from $i_\zeta\omega_0$. To prove this we again start from (26) to write (with $\zeta_1 = \zeta$)

$$i_\zeta\omega(\zeta_2, \ldots, \zeta_k) = \sum_{i=1}^{k} (-1)^{i-1} \omega_0(\zeta_0, \boldsymbol{\zeta}_1, \ldots, \hat{\boldsymbol{\zeta}}_i, \ldots, \boldsymbol{\zeta}_k) + \boldsymbol{\omega}(\boldsymbol{\zeta}_1, \ldots, \boldsymbol{\zeta}_k).$$

This time, we substitute $\zeta_1 - \boldsymbol{\zeta}_1$ for ζ_0 and observe that the second part of the sum involving ω_0, in view of (25) then precisely cancels the last term. □

Maintaining the same line of approach requires that also for the exterior derivative $d\omega$, we identify the associated $(d\omega)_0$ and the form $\mathbf{d}\boldsymbol{\omega}$ on $Sec(\overline{\pi})$ in the sense of Definition 3. Before we can introduce $d\omega$, we need to give a meaning also to the value of a k-form ω, when say its first argument is taken to be a vector section.

Definition 5. *If ω is a k-form on $Sec(\pi)$, then for $\boldsymbol{\sigma} \in Sec(\overline{\pi})$ and $\zeta_i \in Sec(\pi)$, we put*

$$\omega(\boldsymbol{\sigma}, \zeta_2, \ldots, \zeta_k) = \omega(\zeta_1 + \boldsymbol{\sigma}, \zeta_2, \ldots, \zeta_k) - \omega(\zeta_1, \zeta_2, \ldots, \zeta_k), \tag{50}$$

where ζ_1 is chosen arbitrarily.

For this to make sense of course, we need to be sure that the result does not depend on the choice of ζ_1. Now, if we evaluate the right-hand side of the defining relation by using (26), we obtain

$$\omega(\boldsymbol{\sigma},\zeta_2,\ldots,\zeta_k)=\boldsymbol{\omega}(\boldsymbol{\sigma},\boldsymbol{\zeta}_2,\ldots,\boldsymbol{\zeta}_k)+\sum_{i=2}^{k}(-1)^{i-1}\omega_0(\zeta_0,\boldsymbol{\sigma},\boldsymbol{\zeta}_2,\ldots,\hat{\boldsymbol{\zeta}}_i,\ldots,\boldsymbol{\zeta}_k). \tag{51}$$

The right-hand side of this explicit expression makes no mentioning of ζ_1 anymore. It might seem at first sight that we have shifted the problem, because it does depend on the reference section ζ_0. However, we have argued before that (26) does not depend on the choice of such a reference section, whence our newly defined concept makes sense.

The explicit formula (51) further shows that $i_{\boldsymbol{\sigma}}\omega$ is well defined as a $(k-1)$-form, in the sense of Definition 3. The first term on the right identifies its associated form on $Sec(\overline{\pi})$, whereas the second term, upon swapping the first two arguments, reveals that $(i_{\boldsymbol{\sigma}}\omega)_0=i_{\boldsymbol{\sigma}}\omega_0$. As for local computations, it follows from the definition (50) that $e^0(\boldsymbol{\sigma})=0$, whereas the $\boldsymbol{e}^\alpha$ act on $\boldsymbol{\sigma}$ simply as duals of $Sec(\overline{\pi})$.

Definition 6. *The exterior derivative of ω, denoted by $d\omega$ is defined by*

$$\begin{aligned} d\omega(\zeta_1,\ldots,\zeta_{k+1}) &= \sum_{i=1}^{k+1}(-1)^{i-1}\lambda(\zeta_i)\Big(\omega(\zeta_1,\ldots,\hat{\zeta}_i,\ldots,\zeta_{k+1})\Big) \\ &\quad+\sum_{1\le i<j\le k+1}(-1)^{i+j}\omega([\zeta_i,\zeta_j],\zeta_1,\ldots,\hat{\zeta}_i,\ldots,\hat{\zeta}_j,\ldots,\zeta_{k+1}). \end{aligned} \tag{52}$$

Note first that we are making use of definition 5 in the second term on the right, because the bracket of two affine sections is a vector section. It is fairly obvious that $d\omega$ is skew-symmetric in all its arguments. To further justify this definition in the sense of the defining property (26) of forms, note first that we of course have an exterior derivative at our disposal for the k-form $\boldsymbol{\omega}$ on $Sec(\overline{\pi})$ which we denote by d also. We know that $d\boldsymbol{\omega}$ has a property of type (52) (or can be defined that way), with vector sections replacing affine sections and ρ as anchor map instead of λ. It remains to give a meaning to $d\omega_0$.

Definition 7. *For $\omega_0 : Sec(\pi)\times Sec(\overline{\pi})\times\cdots\times Sec(\overline{\pi})\to C^\infty(M)$, we define $d\omega_0$, an operator of the same type, but depending on one more vector section, by*

$$\begin{aligned} d\omega_0(\zeta,\boldsymbol{\zeta}_2,\ldots,\boldsymbol{\zeta}_{k+1}) &= \lambda(\zeta)\Big(\boldsymbol{\omega}(\boldsymbol{\zeta}_2,\ldots,\boldsymbol{\zeta}_{k+1})\Big) \\ &\quad+\sum_{i=2}^{k+1}(-1)^{i-1}\rho(\boldsymbol{\zeta}_i)\Big(\omega_0(\zeta,\boldsymbol{\zeta}_2,\ldots,\hat{\boldsymbol{\zeta}}_i,\ldots,\boldsymbol{\zeta}_{k+1})\Big) \\ &\quad+\sum_{j=2}^{k+1}(-1)^{j+1}\boldsymbol{\omega}([\zeta,\boldsymbol{\zeta}_j],\boldsymbol{\zeta}_2,\ldots,\hat{\boldsymbol{\zeta}}_j,\ldots,\boldsymbol{\zeta}_{k+1}) \\ &\quad-\sum_{2\le i<j\le k+1}(-1)^{i+j}\omega_0(\zeta,[\boldsymbol{\zeta}_i,\boldsymbol{\zeta}_j],\boldsymbol{\zeta}_2,\ldots,\hat{\boldsymbol{\zeta}}_i,\ldots,\hat{\boldsymbol{\zeta}}_j,\ldots,\boldsymbol{\zeta}_{k+1}). \end{aligned} \tag{53}$$

This expression may look rather exotic at first, but it is obtained by formally copying the definition (52) and writing in that process either λ or ρ, and either ω_0 or $\boldsymbol{\omega}$, in such a way that every term in the right-hand side has a proper meaning. There are two important observations to be made here. First of all, the required linearity of $d\omega_0$ in its vector arguments relies on the properties (9) and (11) of our Lie algebroid bracket. Secondly, replacing the affine section ζ in the definition by $\zeta + \boldsymbol{\sigma}$, we find:

$$d\omega_0(\zeta + \boldsymbol{\sigma}, \boldsymbol{\zeta}_2, \ldots, \boldsymbol{\zeta}_{k+1}) = d\omega_0(\zeta, \boldsymbol{\zeta}_2, \ldots, \boldsymbol{\zeta}_{k+1}) + d\boldsymbol{\omega}(\boldsymbol{\sigma}, \boldsymbol{\zeta}_2, \ldots, \boldsymbol{\zeta}_{k+1}). \tag{54}$$

We now know what to expect for the decomposition (26) of $d\omega$ and this is confirmed by the following result.

Proposition 3. *We have* $(d\omega)_0 = d\omega_0$ *and* $\mathbf{d}\boldsymbol{\omega} = d\boldsymbol{\omega}$.

The proof involves a rather technical but straightforward calculation and therefore is omitted. It is of some interest, however, to work out some simple cases in detail. For a function $f \in C^\infty(M)$, df is defined by

$$df(\zeta) = \lambda(\zeta)(f) = \lambda(\zeta_0)(f) + \rho(\boldsymbol{\zeta})(f), \tag{55}$$

from which we learn that $(df)_0 = df$ (as expected) and $\mathbf{df}(\boldsymbol{\sigma}) = \rho(\boldsymbol{\sigma})(f)$. If θ is a 1-form, the defining relation (52) for its exterior derivative reads

$$d\theta(\zeta_1, \zeta_2) = \lambda(\zeta_1)(\theta(\zeta_2)) - \lambda(\zeta_2)(\theta(\zeta_1)) - \boldsymbol{\theta}([\zeta_1, \zeta_2]). \tag{56}$$

Introducing an arbitrary reference section ζ_0, it is easy to verify that this can be rewritten as

$$d\theta(\zeta_1, \zeta_2) = (d\theta)_0(\zeta_0, \boldsymbol{\zeta}_2) - (d\theta)_0(\zeta_0, \boldsymbol{\zeta}_1) + \mathbf{d}\boldsymbol{\theta}(\boldsymbol{\zeta}_1, \boldsymbol{\zeta}_2), \tag{57}$$

where $\mathbf{d}\boldsymbol{\theta} = d\boldsymbol{\theta}$ and

$$(d\theta)_0(\zeta, \boldsymbol{\sigma}) = \lambda(\zeta)(\boldsymbol{\theta}(\boldsymbol{\sigma})) - \rho(\boldsymbol{\sigma})(\theta(\zeta)) - \boldsymbol{\theta}([\zeta, \boldsymbol{\sigma}]). \tag{58}$$

This is in perfect agreement with the results of Proposition 3 and Definition 7.

Concerning derivation properties, it is trivial to verify that for the product of functions: $d(fg) = f dg + g df$. Also, from (56) applied to $f\theta$ we get:

$$\begin{aligned} d(f\theta)(\zeta_1, \zeta_2) &= \lambda(\zeta_1)(f\theta(\zeta_2)) - \lambda(\zeta_2)(f\theta(\zeta_1)) - f\boldsymbol{\theta}([\zeta_1, \zeta_2]) \\ &= df(\zeta_1)\theta(\zeta_2) - df(\zeta_2)\theta(\zeta_1) + f\Big(\lambda(\zeta_1)(\theta(\zeta_2)) - \lambda(\zeta_2)(\theta(\zeta_1)) - \boldsymbol{\theta}([\zeta_1, \zeta_2])\Big), \end{aligned}$$

from which we conclude that

$$d(f\theta) = f\, d\theta + df \wedge \theta, \tag{59}$$

i.e. d has the right derivation property for the action on functions and 1-forms. Recalling now the general statements about derivations we made before, we can conclude that there is a unique derivation of degree 1 on $\bigwedge(\pi^\dagger)$, which coincides with our d on functions and

1-forms. Not surprisingly, this is the d we have introduced by (52), but we again omit the rather technical proof of this claim.

We now reach the main question which is about the relationship between d^2 and the compatibility requirements in the definition of an affine Lie algebroid. To appreciate the meaning of the following lemma, we take a step back and assume now that the bracket $[\zeta_i, \zeta_j]$ figuring in the definition (52) of d satisfies the 'Leibniz-type property' (9) with respect to the module structure of $Sec(\pi)$ (and the resulting property (11)), but no further compatibility or Lie algebra conditions a priori. Remember that the property (9) of the bracket was necessary to make sure that $d\omega$ is a form in the first place.

Lemma 1. *For all $\omega \in \bigwedge^k(\pi^\dagger)$ and $\zeta_i \in Sec(\pi)$, we have*

$$\begin{aligned} &d^2\omega(\zeta_1, \ldots, \zeta_{k+2}) = \\ &\quad \sum_{1\le i<j\le k+2} (-1)^{i+j} \Big(\rho([\zeta_i,\zeta_j]) - [\lambda(\zeta_i), \lambda(\zeta_j)]\Big) \Big(\omega(\zeta_1, \ldots, \hat{\zeta}_i, \ldots, \hat{\zeta}_j, \ldots, \zeta_{k+2})\Big) \\ &\qquad + \sum_{1\le i<j<l\le k+2} (-1)^{i+j+l} \omega(\textstyle\sum_{i,j,l}[\zeta_i, [\zeta_j, \zeta_l]], \zeta_1, \ldots, \hat{\zeta}_i, \ldots, \hat{\zeta}_j, \ldots, \hat{\zeta}_l, \ldots, \zeta_{k+2}), \end{aligned} \tag{60}$$

(where the smaller summation sign of course refers again to a cyclic sum over the three indices involved).

In fact this lemma, with suitable adaptations, also has a rather universal validity, though we were unable to find similar statements in the literature. For this reason, we give a sketch of the proof here.

If ω is a k-form, then $d^2\omega$ is a $(k+2)$-form with

$$\begin{aligned} d^2\omega(\zeta_1, \ldots, \zeta_{k+2}) \;&=\; \sum_{i=1}^{k+2} (-1)^{i-1} \lambda(\zeta_i) \Big(d\omega(\zeta_1, \ldots, \hat{\zeta}_i, \ldots, \zeta_{k+2})\Big) \\ &\quad + \sum_{1\le i<j\le k+2} (-1)^{i+j} d\omega([\zeta_i,\zeta_j], \zeta_1, \ldots, \hat{\zeta}_i, \ldots, \hat{\zeta}_j, \ldots, \zeta_{k+2}). \end{aligned} \tag{61}$$

If we plug in the definition of $d\omega$, the first term on the right will further decompose into two parts, one involving double and the other involving triple sums. After some elementary multiple sum manipulations, it can be written in the form

$$\begin{aligned} &\sum_{1\le i<j\le k+2} (-1)^{i+j} \Big(\lambda(\zeta_j)\lambda(\zeta_i) - \lambda(\zeta_i)\lambda(\zeta_j)\Big) \Big(\omega(\zeta_1, \ldots, \hat{\zeta}_i, \ldots, \hat{\zeta}_j, \ldots, \zeta_{k+2})\Big) \\ &+ \sum_{1\le i<j<l\le k+2} (-1)^{i+j+l-1} \textstyle\sum_{i,j,l} \Big\{\lambda(\zeta_i)\Big(\omega([\zeta_j,\zeta_l]\Big\}, \zeta_1, \ldots, \hat{\zeta}_i, \ldots, \hat{\zeta}_j, \ldots, \hat{\zeta}_l, \ldots, \zeta_{k+2})\Big), \end{aligned} \tag{62}$$

where the smaller summation sign stands for a cyclic sum, the range of which is delimited by the curly brackets. For the second term on the right in (61), we have to remember that the first argument is a vector section. Using the defining relation (50), applied to

$d\omega$, we obtain:

$$\begin{aligned} d\omega(\boldsymbol{\sigma},\zeta_1,\ldots,\zeta_k) &= \rho(\boldsymbol{\sigma})\Big(\omega(\zeta_1,\ldots,\zeta_k)\Big) \\ &+ \sum_{i=1}^{k}(-1)^i\lambda(\zeta_i)\Big(\omega(\boldsymbol{\sigma},\zeta_1,\ldots,\hat{\zeta_i},\ldots,\zeta_k)\Big) \\ &+ \sum_{j=1}^{k}(-1)^j\omega([\boldsymbol{\sigma},\zeta_j],\zeta_1,\ldots,\hat{\zeta_j},\ldots,\zeta_k) \\ &+ \sum_{1\le i<j\le k}(-1)^{i+j}\omega([\zeta_i,\zeta_j],\boldsymbol{\sigma},\zeta_1,\ldots,\hat{\zeta_i},\ldots,\hat{\zeta_j},\ldots,\zeta_k). \end{aligned} \tag{63}$$

The last line here has two vector arguments, but this is consistent with the application of definition 5 to a form of type $i_{\boldsymbol{\zeta}}\omega$. We look at the effect of each of these four terms, when inserted in the second sum of (61). The first one simply gives:

$$\sum_{1\le<i<j\le k+2}(-1)^{i+j}\rho([\zeta_i,\zeta_j])\Big(\omega(\zeta_1,\ldots,\hat{\zeta_i},\ldots,\hat{\zeta_j},\ldots,\zeta_{k+2})\Big). \tag{64}$$

The second one is easily seen to give rise to terms which cancel exactly the second sum in (62). The third term of (63) gives rise to expressions involving double brackets, which combine to:

$$\sum_{1\le i<j<l\le k+2}(-1)^{i+j+l}\omega(\textstyle\sum_{i,j,l}[[\zeta_i,\zeta_j],\zeta_l],\zeta_1,\ .\ ,\hat{\zeta_i},\ .\ ,\hat{\zeta_j},\ .\ ,\hat{\zeta_l},\ .\ ,\zeta_{k+2}). \tag{65}$$

The fourth term of (63) finally creates terms which involve two double sums, and in each of the summands the first two arguments of ω are brackets. One can show by a suitable re-shuffling of summations, that all terms in this part cancel out. What we are left with in the end is the first term of (62), (64) and (65): they precisely combine to the statement in the lemma.

Proposition 4. *The exterior derivative has the property $d^2 = 0$ if and only if the bracket further satisfies the Jacobi identity (16) (or equivalently (8)).*

Proof: If (9) and (16) hold true, we also have (8) and know from previous considerations that (12) then holds as well. The above lemma this way trivially implies $d^2 = 0$. For the converse, we observe that $d^2 f = 0$, for $f \in C^\infty(M)$, implies (17), from which it subsequently follows that $d^2\theta = 0$, with $\theta \in \bigwedge^1(\pi^\dagger)$, implies (16). □

It remains now to list coordinate expressions for the basic exterior derivatives. Let (t, x^i) as before be coordinates on M. For their exterior derivatives we obtain the following: for all $\zeta \in Sec(\pi)$,

$$dt(\zeta) = \lambda(\zeta)(t) = 1, \qquad dx^i(\zeta) = \lambda(\zeta)(x^i),$$

from which it follows that $\mathbf{dt} = \mathbf{0}$ and $\mathbf{dx^i}(\boldsymbol{\sigma}) = \rho(\boldsymbol{\sigma})(x^i)$ (and of course $(dt)_0 = dt$, $(dx^i)_0 = dx^i$). In terms of the general representation (36) of a 1-form, we thus have:

$$dt = e^0, \tag{66}$$

$$dx^i = \lambda^i e^0 + \rho^i_\alpha \boldsymbol{e}^\alpha. \tag{67}$$

Obviously, we have $de^0 = 0$. We further calculate, making use, for example, of (43,44), the general formula (56) and the coordinate expressions (18), that

$$d\boldsymbol{e}^\alpha = -C^\alpha_\beta e^0 \wedge \boldsymbol{e}^\beta - \tfrac{1}{2} C^\alpha_{\beta\gamma} \boldsymbol{e}^\beta \wedge \boldsymbol{e}^\gamma. \tag{68}$$

It is instructive to verify that expressing the properties $d^2\boldsymbol{e}^\alpha = 0$ and $d^2 x^i = 0$ is indeed equivalent to the requirements (20,21) and (22,23), respectively.

To complete the picture of basic derivations on $\bigwedge(\pi^\dagger)$, we have a closer look at the analogue of the classical Lie derivative.

Definition 8. *For every $\zeta \in Sec(\pi)$, the derivation d_ζ of degree zero is defined as*

$$d_\zeta = [i_\zeta, d] = i_\zeta \circ d + d \circ i_\zeta. \tag{69}$$

So, since d_ζ is defined as a commutator of derivations, we know that it will itself be a derivation of degree zero:

$$d_\zeta(\omega \wedge \mu) = d_\zeta \omega \wedge \mu + \omega \wedge d_\zeta \mu. \tag{70}$$

Likewise, we can rely on proofs similar to those in the standard theory to conclude that the following commutator properties will hold true:

$$[d_\zeta, i_\eta] = i_{[\zeta,\eta]}, \qquad [d_\zeta, d] = 0, \qquad [d_\zeta, d_\eta] = d_{[\zeta,\eta]}. \tag{71}$$

Note, however, that a Lie-type derivation with respect to a vector section turns up in the last property, and this is indeed well defined also as: $d_{\boldsymbol{\sigma}} = [i_{\boldsymbol{\sigma}}, d]$. It is further natural to extend the action of d_ζ to $Sec(\pi)$ by duality. It then follows, as expected, that for $\eta, \zeta \in Sec(\pi)$,

$$d_\zeta \eta = [\zeta, \eta]. \tag{72}$$

As a result of such an extension, d_ζ has Leibniz-type properties also with respect to the evaluation of forms on the appropriate number of affine (or vector) sections; the following property, which could be verified by a direct computation from the definition of d_ζ, thus becomes self-evident:

$$d_\eta \omega(\zeta_1, \ldots, \zeta_k) = \lambda(\eta)\Big(\omega(\zeta_1, \ldots, \zeta_k)\Big) + \sum_{j=1}^{k} (-1)^j \omega([\eta, \zeta_j], \zeta_1, \ldots, \hat{\zeta_j}, \ldots, \zeta_k). \tag{73}$$

In the interest of doing computations, we list the Lie-type derivatives of functions $f \in C^\infty(M)$ and the local basis of 1-forms. For $\zeta = e_0 + \zeta^\alpha \boldsymbol{e}_\alpha$,

$$d_\zeta f = \lambda(\zeta)(f), \qquad d_\zeta e^0 = 0, \qquad d_\zeta \boldsymbol{e}^\alpha = C^\alpha_\beta \zeta^\beta e^0 - C^\alpha_\beta \boldsymbol{e}^\beta + C^\alpha_{\beta\gamma} \zeta^\gamma \boldsymbol{e}^\beta + d\zeta^\alpha.$$

4 λ-admissible curves and dynamics

As we expressed in the introduction, the model of affine Lie algebroids we are developing should in the first place offer an environment in which one can accommodate the time-dependent Lagrange-type equations (4). At present, we wish to look in more detail at the geometric nature of the more general dynamical systems, which we call pseudo-second-order equations, and are those described by differential equations of the form (3). For this purpose in fact, we do not need the full machinery of algebroids: it suffices to assume that E is an affine bundle over M and $\lambda : E \to J^1M$ an affine bundle map over the identity.

Definition 9. *A curve ψ in E, which is a section of $\tau \circ \pi$, is said to be λ-admissible, if $\lambda \circ \psi = j^1(\pi \circ \psi)$.*

One could say that ψ is the λ-prolongation of a curve in M. In coordinates, we have

$$\psi : t \mapsto (t, x^i(t), y^\alpha(t)), \qquad \text{with} \quad \dot{x}^i(t) = \lambda^i(t, x(t)) + \rho^i_\alpha(t, x(t))\, y^\alpha(t).$$

Note in passing that, not unexpectedly, one can characterize λ-admissibility via a concept of contact forms: putting $\Theta^i = \lambda^*\theta^i$, where the θ^i are the contact forms on J^1M, we have that ψ is a λ-admissible curve in E if and only if $\psi^*\Theta^i = 0$.

Pseudo-second-order equation fields on E are vector fields whose integral curves all are λ-admissible curves. As in the standard theory of SODEs on a tangent bundle or first jet bundle, however, there is a simple direct characterization of such vector fields.

Definition 10. *$\Gamma \in \mathcal{X}(E)$ is a pseudo-second-order equation field if*

$$T\pi \circ \Gamma = i \circ \lambda,$$

where i is the injection of J^1M into TM.

Clearly, in coordinates, a pseudo-SODE is of the form

$$\Gamma = \frac{\partial}{\partial t} + (\lambda^i(t,x) + \rho^i_\alpha(t,x)y^\alpha)\frac{\partial}{\partial x^i} + f^\alpha(t,x,y)\frac{\partial}{\partial y^\alpha}, \tag{74}$$

for some functions f^α, and it is obvious that all its integral curves will be λ-admissible.

The following diagram visualizes the notions of λ-admissible curves and pseudo-SODEs.



An important point now, however, is that there is a natural way of interpreting the vector field Γ as section of a different bundle.

From the above definition, it is clear that a pseudo-SODE is actually a section of $(\tau \circ \pi)^0_1 : J^1E \to E$, with the additional property that for all $p \in E$, $T\pi|_{J^1E}(\Gamma(p)) = \lambda(p)$. An equivalent way of saying the same thing, by definition of the concept of a pullback bundle, is that $(p, \Gamma(p))$ is a point of $\lambda^* J^1E$, with J^1E regarded as fibred over J^1M via $T\pi|_{J^1E}$. From now on, we will write $J^1_\lambda E$ for $\lambda^* J^1E$, and denote its two projections as indicated in the following diagram:

$$\begin{array}{ccc} J^1_\lambda E & \xrightarrow{\lambda^1} & J^1E \\ {\scriptstyle \pi_2}\downarrow & & \downarrow{\scriptstyle T\pi} \\ E & \xrightarrow[\lambda]{} & J^1M \end{array}$$

If we finally put $\pi_1 = (\tau \circ \pi)^0_1 \circ \lambda^1$, there is yet another way of expressing the characterization of a pseudo-SODE. Indeed, from the trivial observation that $(\tau \circ \pi)^0_1(\Gamma(p)) = p = \pi_2((p, \Gamma(p)))$, it follows that a pseudo-SODE Γ can be regarded also as a section of the bundle $\pi_1 : J^1_\lambda E \to E$, with the property that $\pi_2 \circ \Gamma = \pi_1 \circ \Gamma$.

The various spaces and projections, described in this discussion, are depicted in the diagram of the next section. This diagram immediately suggests the following question: if we put more structure into the scheme by assuming now again that $\pi : E \to M$ carries an affine Lie algebroid structure, is it possible to prolong this structure to the bundle $\pi_1 : J^1_\lambda E \to E$, in such a way of course that λ^1 becomes the anchor map of the induced affine Lie algebroid?

5 Prolongation of affine Lie algebroids

We shall now look in more detail at the bundle $\pi_1 : J^1_\lambda E \to E$. Its total space is the manifold

$$J^1_\lambda E = \lambda^* J^1E = \{(q, Z) \in E \times J^1E \mid \lambda(q) = T\pi|_{J^1E}(Z)\},$$

but the fibration we want to focus on is not one of the projections which define $J^1_\lambda E$, but rather the map $\pi_1 = (\tau \circ \pi)^0_1 \circ \lambda^1$. As such, we are looking at an affine bundle, modeled on the vector bundle $\overline{\pi}_1 : V_\rho E \to E$, with total space

$$V_\rho E = \{(\boldsymbol{v}, \boldsymbol{V}) \in V \times VE \mid \rho(\boldsymbol{v}) = T\pi|_{VE}(\boldsymbol{V})\}.$$

The affine bundles involved, and their underlying vector bundles are illustrated below.



A section Z of π_1 is completely determined once we know the maps $\pi_2 \circ Z : E \to E$ and $\lambda^1 \circ Z : E \to J^1E$. Likewise, vector sections $\boldsymbol{Z}$ of $\overline{\pi}_1$ are determined by $\overline{\pi}_2 \circ \boldsymbol{Z}$ and $\rho^1 \circ \boldsymbol{Z}$. For example, let $e \in E$ be a point with coordinates (t, x^i, y^α), so that (t, x^i) are the coordinates of $\pi(e) \in M$ and e has the representation $e = e_0 + y^\alpha \boldsymbol{e}_\alpha$. If then Z is a section of π_1, we will have:

$$\begin{aligned} \pi_2 \circ Z : &\qquad (t,x,y) \longmapsto (t,x,z^\alpha(t,x,y)), \\ \lambda^1 \circ Z : &\qquad (t,x,y) \longmapsto \left(\frac{\partial}{\partial t} + (\lambda^i + \rho^i_\alpha z^\alpha)\frac{\partial}{\partial x^i} + Z^\alpha \frac{\partial}{\partial y^\alpha}\right)\Big|_e , \end{aligned}$$

and determining Z in coordinates of course amounts to assigning the functions (z^α, Z^α) on E.

It is worthwhile looking at the representation of such a Z with respect to suitably selected local sections of π_1 and $\overline{\pi}_1$, which will exhibit the affine structure of π_1 and are adapted to the basis which was selected to coordinatize E. To this end, we introduce two sets of local sections $\boldsymbol{\mathcal{X}}_\alpha$ and $\boldsymbol{\mathcal{V}}_\alpha$ of $\overline{\pi}_1$ which will span $Sec(\overline{\pi}_1)$, and select a zero section $\mathcal{E}_0$ as follows. The $\boldsymbol{\mathcal{V}}_\alpha$ span 'vertical sections' and are determined by: $\overline{\pi}_2 \circ \boldsymbol{\mathcal{V}}_\alpha = \mathbf{0}$, while for $e \in E_m$ we let $\rho^1 \circ \boldsymbol{\mathcal{V}}_\alpha(e)$ be the tangent vector to the curve $s \mapsto e + s\,\boldsymbol{e}_\alpha(m)$ in E_m. Verticality is an intrinsic property whereas, as usual, there is no intrinsic notion of horizontality. The determination of the $\boldsymbol{\mathcal{X}}_\alpha$ and $\mathcal{E}_0$ will therefore rely on pure coordinate arguments. For the projection onto V we put $\overline{\pi}_2 \circ \boldsymbol{\mathcal{X}}_\alpha = \boldsymbol{e}_\alpha \circ \pi$ and then, fixing $\rho^1 \circ \boldsymbol{\mathcal{X}}_\alpha$ (as a vector field on E) further requires making a prescription for the vertical components, which we simply take to be zero. Similarly, for the choice of a zero section, we could take any vector field on E which projects under $T\pi$ onto $\lambda(e_0) \in \mathcal{X}(M)$ (and as such defines also a section of π_1), but we will fix it also by taking the vertical components to be zero. Thus we have:

$$\boldsymbol{\mathcal{X}}_\alpha(e) = \left(\boldsymbol{e}_\alpha(\pi(e)),\, \rho^i_\alpha(t,x)\frac{\partial}{\partial x^i}\Big|_e\right) \qquad \boldsymbol{\mathcal{V}}_\alpha(e) = \left(\mathbf{0}(\pi(e)),\, \frac{\partial}{\partial y^\alpha}\Big|_e\right), \tag{75}$$

and

$$\mathcal{E}_0(e) = \left(e_0(\pi(e)),\, \left(\frac{\partial}{\partial t} + \lambda^i(t,x)\frac{\partial}{\partial x^i}\right)\Big|_e\right). \tag{76}$$

The general section Z of π_1 then has the local representation:

$$Z = \mathcal{E}_0 + z^\alpha(t,x,y)\boldsymbol{\mathcal{X}}_\alpha + Z^\alpha(t,x,y)\boldsymbol{\mathcal{V}}_\alpha. \tag{77}$$

Note that pseudo-SODEs, as discussed in the previous section, are precisely those sections Γ of π_1, for which $z^\alpha(t,x,y) = y^\alpha$.

Let now E be equipped with an affine Lie algebroid structure. To be in line with the notations we used in Definition 1, we will from now on also write $\lambda^1(Z)$ instead of $\lambda^1 \circ Z$, and likewise for the π_2-projection and the corresponding projections of vector sections.

We wish to establish that there is an induced Lie algebroid structure on the affine bundle π_1. To this end, following the scheme of Definition 1, we have to identify a bracket on $\overline{\pi}_1$ and an action of affine sections on vector sections, such that all the necessary requirements are met. The idea is to define such brackets by requiring roughly that its two projections are determined by the known brackets of the projected sections. But there are some technical complications which we will address now.

For $\boldsymbol{Z}_1, \boldsymbol{Z}_2 \in Sec(\overline{\pi}_1)$, a preliminary observation is that the Lie bracket of their image under ρ^1 (which gives rise to vector fields on E), belongs to the image of ρ^1. A coordinate calculation can confirm this. Putting

$$\rho^1(\boldsymbol{Z}_i) = z_i^\alpha \rho_\alpha^j \frac{\partial}{\partial x^j} + Z_i^\alpha \frac{\partial}{\partial y^\alpha},$$

we have

$$\begin{aligned}[\rho^1(\boldsymbol{Z}_1), \rho^1(\boldsymbol{Z}_2)] \;=\; & \Big(\rho^1(\boldsymbol{Z}_1)(z_2^\alpha) - \rho^1(\boldsymbol{Z}_2)(z_1^\alpha)\Big)\rho_\alpha^j \frac{\partial}{\partial x^j} \\ & + \Big(z_2^\alpha \rho^1(\boldsymbol{Z}_1)(\rho_\alpha^j) - z_1^\alpha \rho^1(\boldsymbol{Z}_2)(\rho_\alpha^j)\Big)\frac{\partial}{\partial x^j} + \cdots \frac{\partial}{\partial y^\alpha}.\end{aligned}$$

The first term on the right manifestly belongs to the image of ρ^1, whereas the last term is irrelevant for that purpose. The middle term can be rewritten as

$$z_2^\alpha z_1^\beta \left(\rho_\beta^j \frac{\partial \rho_\alpha^i}{\partial x^j} - \rho_\alpha^j \frac{\partial \rho_\beta^i}{\partial x^j}\right) \frac{\partial}{\partial x^i},$$

which is seen to belong to the image of ρ^1 in view of the property (23). It is therefore natural to impose right away that the bracket $[\cdot,\cdot]^1$ under construction, which of course is required to be skew-symmetric and $\mathbb{R}$-bilinear, should satisfy

$$\rho^1\left([\boldsymbol{Z}_1, \boldsymbol{Z}_2]^1\right) = [\rho^1(\boldsymbol{Z}_1), \rho^1(\boldsymbol{Z}_2)]. \tag{78}$$

This will have for consequence that for $F_i \in C^\infty(E)$,

$$\begin{aligned}\rho^1\left([F_1\,\boldsymbol{Z}_1, F_2\,\boldsymbol{Z}_2]^1\right) \;=\; & \\ & F_1F_2\,[\rho^1(\boldsymbol{Z}_1), \rho^1(\boldsymbol{Z}_2)] + F_1\,\rho^1(\boldsymbol{Z}_1)(F_2)\,\rho^1(\boldsymbol{Z}_2) - F_2\,\rho^1(\boldsymbol{Z}_2)(F_1)\,\rho^1(\boldsymbol{Z}_1).\end{aligned}$$

It remains then to make sure that the projection under $\overline{\pi}_2$ can be specified in a compatible way. The above coordinate calculation to some extent illustrates how one should proceed. If we apply $T\pi$ to the preceding equality, we get (pointwise)

$$\begin{aligned}T\pi\left(\rho^1\left([F_1\,\boldsymbol{Z}_1, F_2\,\boldsymbol{Z}_2]^1\right)\right) = (F_1F_2)\,T\pi\left([\rho^1(\boldsymbol{Z}_1), \rho^1(\boldsymbol{Z}_2)]\right) \\ + F_1\,\rho^1(\boldsymbol{Z}_1)(F_2)\,\rho(\overline{\pi}_2(\boldsymbol{Z}_2)) - F_2\,\rho^1(\boldsymbol{Z}_2)(F_1)\,\rho(\overline{\pi}_2(\boldsymbol{Z}_1)).\end{aligned}$$

In general, the $\rho(\overline{\pi}_2(\boldsymbol{Z}_i))$ are vector fields along π for which there is no standard Lie bracket available. If the $\boldsymbol{Z}_i$ are projectable, however, meaning that there exist $\boldsymbol{\zeta}_i \in Sec(\overline{\pi})$ such that $\overline{\pi}_2 \circ \boldsymbol{Z}_i = \boldsymbol{\zeta}_i \circ \pi$, the vector fields $\rho^1(\boldsymbol{Z}_i)$ on E are π-related to the vector fields $\rho(\boldsymbol{\zeta}_i)$ on M. Hence, the corresponding brackets are also π-related, meaning that for projectable $\boldsymbol{Z}_i$, we can put

$$\overline{\pi}_2\left([\boldsymbol{Z}_1, \boldsymbol{Z}_2]^1\right) = [\overline{\pi}_2(\boldsymbol{Z}_1), \overline{\pi}_2(\boldsymbol{Z}_2)], \tag{79}$$

and then the property (13) (which in coordinates gives (23)) ensures that

$$T\pi\left(\rho^1\left([\boldsymbol{Z}_1, \boldsymbol{Z}_2]^1\right)\right) = \rho \circ \overline{\pi}_2\left([\boldsymbol{Z}_1, \boldsymbol{Z}_2]^1\right)$$

as it should. The expression for $T\pi\left(\rho^1\left([F_1\,\boldsymbol{Z}_1, F_2\,\boldsymbol{Z}_2]^1\right)\right)$ further shows that the $\overline{\pi}_2$ and ρ^1 projections of the bracket under construction will still match up if for projectable $\boldsymbol{Z}_i$ and for any $F_i \in C^\infty(E)$, we define

$$\begin{aligned}\overline{\pi}_2\left([F_1\,\boldsymbol{Z}_1, F_2\,\boldsymbol{Z}_2]^1\right) &= F_1 F_2\,[\overline{\pi}_2(\boldsymbol{Z}_1), \overline{\pi}_2(\boldsymbol{Z}_2)] \\ &\quad + F_1\,\rho^1(\boldsymbol{Z}_1)(F_2)\,\overline{\pi}_2(\boldsymbol{Z}_2) - F_2\,\rho^1(\boldsymbol{Z}_2)(F_1)\,\overline{\pi}_2(\boldsymbol{Z}_1).\end{aligned} \tag{80}$$

It then follows that

$$[F_1\,\boldsymbol{Z}_1, F_2\,\boldsymbol{Z}_2]^1 = F_1 F_2\,[\boldsymbol{Z}_1, \boldsymbol{Z}_2]^1 + F_1\,\rho^1(\boldsymbol{Z}_1)(F_2)\,\boldsymbol{Z}_2 - F_2\,\rho^1(\boldsymbol{Z}_2)(F_1)\,\boldsymbol{Z}_1, \tag{81}$$

since both sides have the same $\overline{\pi}_2$ and ρ^1 projections.

The final point to observe now is that sections of $\overline{\pi}_1$ (locally) are finitely generated, over the ring $C^\infty(E)$, by projectable sections. Hence, the defining relations (78) and (80) are sufficient to define the bracket $[\cdot,\cdot]^1$ on vector sections. The property (81) will hold by extension for all vector sections and the bracket will satisfy the Jacobi identity as a result of the Jacobi identity of the Lie algebroid bracket we start from and the same identity for vector fields on E.

To define the action of $Z \in Sec(\pi_1)$ on $\boldsymbol{V} \in Sec(\overline{\pi}_1)$, we proceed in exactly the same manner. First, one easily verifies that the Lie bracket of $\lambda^1(Z)$ and $\rho^1(\boldsymbol{V})$ belongs to the image of ρ^1, this time in view of the properties (22) and (23). Hence, it makes sense to put

$$\rho^1\left([Z, \boldsymbol{V}]^1\right) = [\lambda^1(Z), \rho^1(\boldsymbol{V})], \tag{82}$$

and we of course require the bracket $[\cdot,\cdot]^1$ to have linearity properties of the kind of (7). For projectable sections, we can put

$$\overline{\pi}_2\left([Z, \boldsymbol{V}]^1\right) = [\pi_2(Z), \overline{\pi}_2(\boldsymbol{V})], \tag{83}$$

and be assured of consistency with the projection (82). Next, still for projectable Z and $\boldsymbol{V}$ and for any $F \in C^\infty(E)$, we define

$$\overline{\pi}_2\left([Z, F\,\boldsymbol{V}]^1\right) = F\,[\pi_2(Z), \overline{\pi}_2(\boldsymbol{V})] + \lambda^1(Z)(F)\,\overline{\pi}_2(\boldsymbol{V}). \tag{84}$$

Sections of π_1 can be written as a projectable zero section, plus a linear combination of projectable vector sections with coefficients in $C^\infty(E)$. In combination with the earlier arguments for vector sections, we are again led to the conclusion that the requirements

(82) and (84) are sufficient to define $[Z, \boldsymbol{V}]^1$ for arbitrary $Z \in Sec(\pi_1)$ and $\boldsymbol{V} \in Sec(\overline{\pi}_1)$ and that we will have the property

$$[Z, F\,\boldsymbol{V}]^1 = F\,[Z, \boldsymbol{V}]^1 + \lambda^1(Z)(F)\,\boldsymbol{V}. \tag{85}$$

The final requirement of type (8) then also easily follows, which concludes the construction of the prolonged affine Lie algebroid.

For computational purposes, it remains to list the brackets of the local sections which are used in the general representation of a section of π_1 as in (77). We have

$$\begin{array}{lll} [\mathcal{E}_0, \boldsymbol{\mathcal{X}}_\alpha]^1 = C^\beta_\alpha \boldsymbol{\mathcal{X}}_\beta, & & [\mathcal{E}_0, \boldsymbol{\mathcal{V}}_\alpha]^1 = \mathbf{0}, \\ [\boldsymbol{\mathcal{X}}_\alpha, \boldsymbol{\mathcal{X}}_\beta]^1 = C^\gamma_{\alpha\beta} \boldsymbol{\mathcal{X}}_\gamma, & [\boldsymbol{\mathcal{X}}_\alpha, \boldsymbol{\mathcal{V}}_\beta]^1 = \mathbf{0}, & [\boldsymbol{\mathcal{V}}_\alpha, \boldsymbol{\mathcal{V}}_\beta]^1 = \mathbf{0}. \end{array}$$

It is perhaps worthwhile to repeat hereby that the two projections have to be looked at to verify these statements, although of course they are bound to match up if our new bracket has been defined consistently. Thus we have, for example:

$$\begin{aligned} \overline{\pi}_2\left([\mathcal{E}_0, \boldsymbol{\mathcal{X}}_\alpha]^1\right) &= [\pi_2(\mathcal{E}_0), \overline{\pi}_2(\boldsymbol{\mathcal{X}}_\alpha)] = [e_0, \boldsymbol{e}_\alpha] = C^\beta_\alpha \boldsymbol{e}_\beta, \\ \rho^1\left([\mathcal{E}_0, \boldsymbol{\mathcal{X}}_\alpha]^1\right) &= [\lambda^1(\mathcal{E}_0), \rho^1(\boldsymbol{\mathcal{X}}_\alpha)] = \left[\frac{\partial}{\partial t} + \lambda^i \frac{\partial}{\partial x^i}, \rho^j_\alpha \frac{\partial}{\partial x^j}\right] = C^\beta_\alpha\, \rho^1(\boldsymbol{\mathcal{X}}_\beta), \end{aligned}$$

where (22) has been used again in the last line.

6 Discussion and outlook for future work

The form of equations (4), which we claim to be the appropriate generalization of Lagrangian systems on Lie algebroids to a situation where explicit time-dependence is involved, has brought us to the introduction of the new concept of Lie algebroids on affine bundles which are fibred over $\mathbb{R}$. More precisely, the first guidance for developing this concept was provided by the conditions (5) and (6) which the various functions appearing in (4) have to satisfy. Ultimately, of course, we want to arrive at an intrinsic geometrical construction of such Lagrangian systems. There are many aspects to be explored yet, but we have sufficiently paved the way already to be able to predict what the outcome of subsequent studies will bring.

One of us has shown [5] that the prolongation of a Lie algebroid (in the standard situation of vector bundles) provides a platform where there exist analogues of the intrinsic structures living on a tangent bundle and these in turn give rise to an intrinsic definition of Lagrangian systems via Poincaré-Cartan type forms. This is the reason why we were keen to verify immediately that the same notion of prolongation exists in our affine set-up. There is little doubt now that we will find intrinsic objects on such a prolonged affine Lie algebroid, which are analogues of what is known to give rise to an intrinsic definition of time-dependent Lagrangian systems on the first jet bundle of a manifold fibred over $\mathbb{R}$. But there is more to it. Even when there is no Lagrangian for the dynamics under consideration and we are, in other words, talking about pseudo-SODEs on a Lie algebroid, we

expect to be able to develop a machinery of associated non-linear and linear connections, which again is analogous to the standard theory of connections associated to SODEs on a tangent bundle or first jet bundle. A paper on these issues is in preparation.

One of the features we examined in this paper as a kind of test for the relevance and internal consistency of the generalized notion of Lie algebroids, was the existence of an associated coboundary operator d. But of course, there are still other interesting properties which standard Lie algebroids are known to exhibit. Let us briefly highlight another one here and show that it also survives our generalization, namely the existence of an associated Poisson structure. Specifically, we want to establish that there exists a canonically defined Poisson structure on the extended dual $E^\dagger$.

Sections of π (respectively $\overline{\pi}$) can be identified with linear functions on $E^\dagger$ (respectively V^*). Explicitly, if $\zeta \in Sec(\pi)$, we consider the function $\hat{\zeta} \in C^\infty(E^\dagger)$ defined by: for each $p \in E^\dagger$, $p \in E^\dagger_m$ say, $\hat{\zeta}(p) = p(\zeta_m)$. Likewise, if $\boldsymbol{\sigma} \in Sec(\overline{\pi})$, we denote by $\hat{\boldsymbol{\sigma}} \in C^\infty(V^*)$ the function defined by $\hat{\boldsymbol{\sigma}}(\boldsymbol{p}) = \boldsymbol{p}(\boldsymbol{\sigma}_m)$, where $\boldsymbol{p} \in V^*_m$. In coordinates, if $\zeta = e_0 + \zeta^\alpha \boldsymbol{e}_\alpha$ and $p \in E^\dagger$ has coordinates (t, x^i, p_0, p_α), then $\hat{\zeta}(p) = p_0 + p_\alpha \zeta^\alpha(t, x)$ and similarly for $\hat{\boldsymbol{\sigma}}$. Now, for any two sections $\zeta, \eta \in Sec(\pi)$, we define the function $\{\hat{\zeta}, \hat{\eta}\}$ on $E^\dagger$ by

$$\{\hat{\zeta}, \hat{\eta}\}(p) = \widehat{[\zeta, \eta]}(\boldsymbol{p}), \tag{86}$$

whereby we recall that $[\zeta, \eta]$ is a section of $\overline{\pi}$ and for $p \in E^\dagger_m$, $\boldsymbol{p}$ is the associated element of V^*_m. If further f, g are functions on M and we make no notational distinction for their pullback to $E^\dagger$, internal consistency of (86) for the action of $Sec(\overline{\pi})$ on $Sec(\pi)$ requires that we further put

$$\{\hat{\zeta}, f\} = -\{f, \hat{\zeta}\} = \lambda(\zeta)(f), \qquad \{f, g\} = 0. \tag{87}$$

The construction then uniquely extends to a skew-symmetric, $\mathbb{R}$-bilinear bracket operation on $E^\dagger$ with the required derivation property. This bracket satisfies the Jacobi identity as a result of the Lie algebroid Jacobi identity (16).

The brackets for the coordinate functions on $E^\dagger$ are found to be:

$$\begin{array}{lll}
\{t, t\} = 0 & \{t, x^i\} = 0 & \{x^i, x^j\} = 0 \\
\{p_0, t\} = 1 & \{p_0, x^i\} = \lambda^i & \{p_0, p_\beta\} = C^\gamma_\beta p_\gamma \\
\{p_\alpha, t\} = 0 & \{p_\alpha, x^i\} = \rho^i_\alpha & \{p_\alpha, p_\beta\} = C^\gamma_{\alpha\beta} p_\gamma .
\end{array}$$

There is an interesting observation to be made here. Recall that $E^\dagger$ is actually a vector bundle and note now that the bracket we have constructed preserves the subset of functions on $E^\dagger$ which are linear in the fibre coordinates. As a result, we know that there is an induced Lie algebroid structure on the bundle $(E^\dagger)^* \to M$. There are many new insights to be gained from approaching the subject of a Lie algebroid structure on the affine bundle $E \to M$ from this angle; that is to say, by regarding $E \to M$ as an affine subbundle of $(E^\dagger)^* \to M$ and taking an appropriate Lie algebroid structure on $(E^\dagger)^* \to M$ as the starting point. Also this is the subject of a forthcoming paper [13] and is the point of view also in the preprint [14] which came out during the revision process of the present

work. One of the main advantages which such an approach offers is that it enables us to get beyond the assumption about the extra fibration of M and thus gives rise to a more general concept of affine Lie algebroids. It further leads to considerable simplifications in proving results about the calculus of forms, as discussed here in Section 3. We believe, nevertheless, that the present approach preserves its merits mainly for two reasons: one is the repeatedly mentioned motivation coming from time-dependent mechanics; the second one is that the theory of differential forms on sections of an affine bundle $E \to M$, as investigated in Section 3 from a kind of 'internal point of view', i.e. without recourse to the larger vector bundle in which E can be imbedded, is something which is being developed here (to the best of our knowledge) for the first time.

Acknowledgements. E. Martínez acknowledges partial financial support from CICYT grant BFM2000-1066-C03-01.

References

[1] J.F. Cariñena, In: *Symmetries in quantum mechanics and quantum optics* (University of Burgos (Spain), 1999), 67.

[2] J.F. Cariñena and E. Martínez, In: *Lie algebroids and related topics in differential geometry* (Banach Center Publications **54**, 2001), 201.

[3] J. Clemente-Gallardo, In: *Nonlinear control in the new millenium*, F. Lamnabhi-Lagarrigur, W. Respondek and A. Isidori, eds., (Springer Verlag, 2000).

[4] P. Libermann, Arch. Math. (Brno) **32**, 147 (1996).

[5] E. Martínez, Acta. Appl. Math. **67**, 295 (2001).

[6] A. Weinstein, In: *Mechanics day (Waterloo, ON, 1992)*, (Fields Institute Communications **7**, American Mathematical Society, 1996), 207.

[7] W. Sarlet, T. Mestdag and E. Martínez, In: *Differential Geometry and its Applications, Proc. 8th Int. Conf. (Opava 2001)*, D. Krupka et al (Eds.), to appear.

[8] M. Crampin, *Jet bundle techniques in analytical mechanics*, Quaderni del consiglio nazionale delle ricerche, gruppo nazionale di fisica matematica **47** (1995).

[9] K. Mackenzie, *Lie groupoids and Lie algebroids in differential geometry*, London Math. Soc. Lect. Note Series **124** (Cambridge Univ. Press, 1987).

[10] J. Grabowski and P. Urbanski, Ann. Global An. Geom. **15**, 447 (1997).

[11] J. Grabowski and P. Urbanski, Rep. Math. Phys. **40**, 195 (1997).

[12] A. Frölicher and A. Nijenhuis, Proc. Ned. Acad. Wetensch. Sér. A **59**, 338 (1956).

[13] E. Martínez, T. Mestdag and W. Sarlet, math.DG/0203178 (2002), J. Geom. Phys., to appear.

[14] J. Grabowski, K. Grabowska and P. Urbanski, math.DG/0203112 (2002).