Note on generalised connections and affine bundles

T. Mestdag, W. Sarlet
Department of Mathematical Physics and Astronomy
Ghent University, Krijgslaan 281, B-9000 Ghent, Belgium
E. Martínez
Departamento de Matemática Aplicada
Facultad de Ciencias, Universidad de Zaragoza
Pedro Cerbuna 12, E-50009 Zaragoza, Spain

October 4, 2002

Abstract. We develop an alternative view on the concept of connections over a vector bundle map, which consists of a horizontal lift procedure to a prolonged bundle. We further focus on prolongations to an affine bundle and introduce the concept of affineness of a generalised connection.

1 Introduction

There has been a lot of interest, recently, in potential applications of Lie algebroids in physics, control theory and other fields of applied mathematics. Among papers which study, in particular, aspects of Lagrangian systems on Lie algebroids, we mention [24, 15, 18, 3, 4, 5, 22, 20]. There is of course an enormous literature on more purely mathematical aspects of Lie algebroids, of which we cite only the standard work [16], and [8] for its particular relevance to this paper.

Our recent joint work in the field finds its roots in searching for the right geometrical model for a kind of time-dependent generalisation of 'Lagrangian mechanics' on Lie algebroids. Since ordinary time-dependent mechanics is usually described on the first-jet space J^1M of a manifold M fibred over $\mathbb{R}$ (see for example [6, 17]), the direct model for the generalisation we had in mind was a kind of Lie algebroid structure, whose anchor map takes values in J^1M rather than TM. This was explored in detail in [22], which in turn rose interest in the more general features of having a Lie algebroid structure on an affine bundle, without the requirement that the base manifold be fibred over $\mathbb{R}$. Those ideas were developed in [20] and to some extent (that is without reference to dynamical systems) also in [10].

A continuation of this work is in preparation, in particular with the purpose of bringing a suitable theory of connections into the picture of dynamical systems on affine algebroids.

But the path to these further developments has led us to discover some general features on connections and affine spaces, which do not require a Lie algebroid structure and seem well worth being brought under the attention separately. This brings us to the subject matter of the present paper.

In Section 2, the main objective is to discuss two interesting constructions from the recent literature on generalised connections and algebroids which, when brought together in a unifying picture, will open the way to explain in detail how they are related. Both constructions may have their roots in the theory of Lie algebroids, but have been formulated recently in the more general framework where a vector bundle has a kind of anchor map, but need not be equipped with a Lie algebra structure on the real vector space of its sections. The first topic we are referring to is the notion of generalised connection on a vector bundle map, as introduced by Cantrijn and Langerock [1], inspired by a similar construction on Lie algebroids by Fernandes [8]. The second is the idea of prolongation, which has been discussed in the context of Lie algebroids, for example in [18, 22, 20], but, as shown in [20], can also be defined without the need of a Lie algebra structure. Also relevant is work by Popescu, who in fact already developed the same ideas in the case that all bundles involved are vector bundles; for this we refer to [21] and references therein. We will arrive, in Section 2, at an alternative view on the generalised connections of [1]. But let us mention here already that this alternative view can be developed without needing the generalised connection idea of [1]. This is in fact one of the main discoveries of [18] and [21] and it is being explored to full extent in [19]. The purpose of the present note, however, is to explain in detail the interrelationship between the two ideas.

In Section 3, we focus on the case of the prolongation of an affine bundle $E \to M$ over a vector bundle $V \to M$. We show that bringing the bidual of E into the picture enables us to give a clear and concise definition of the concept of an affine connection over a vector bundle map and prove a result about the equivalent characterisation of such a connection via a kind of covariant derivative operator. The relevance of these results for the future developments we have in mind is briefly indicated in the final section.

2 Connections over a vector bundle map and the horizontal subbundle of a prolonged bundle

We start by recalling the prolongation idea, as developed in [20], following ideas first introduced in [11].

Let $\mu : P \to M$ be an arbitrary fibre bundle and $\tau : V \to M$ a vector bundle. Assume there exists an *anchor map* $\rho : V \to TM$, which for the time being is just a vector bundle morphism.

Definition 1. *The ρ-prolongation of $\mu : P \to M$ is the bundle $\mu^1 : T^\rho P \to P$, constructed as follows: (i) the total space $T^\rho P$ is the total space of the pullback bundle $\rho^* TP$*

$$T^\rho P = \{(v, X_p) \in V \times TP \mid \rho(v) = T\mu(X_p)\}; \tag{1}$$

(ii) if ρ^1 denotes the projection of ρ^*TP into TP and τ_P is the tangent bundle projection, then $\mu^1 = \tau_P \circ \rho^1$.

The situation is summarised in the following diagram, whereby the projection on the first element of a pair $(v, X_p) \in T^\rho P$ is denoted by μ^2.



Notice that, if we have two bundles $\mu_i : P_i \to M$ $(i = 1, 2)$, and a bundle map f (over the identity on M) between them, then the tangent map $Tf : TP_1 \to TP_2$ extends to a map $T^\rho P_1 \to T^\rho P_2 : (v, X_p) \mapsto (v, Tf(X_p))$. Indeed, we have $T\mu_2(Tf(X_p)) = T\mu_1(X_p) = \rho(v)$. There is more to say about the tangent-bundle-like behaviour of $T^\rho P$, but we will not elaborate on that here.

Coming back to the diagram above, an element of $T^\rho P$ is called *vertical* if it is in the kernel of the projection μ^2. The set of all vertical elements in $T^\rho P$ is a vector subbundle of μ^1 and will be denoted by $\mathcal{V}^\rho P$. If $(0, Q) \in \mathcal{V}^\rho P$, then $Q = \rho^1(0, Q)$ will also be vertical in TP, since $T\mu(Q) = \rho(0) = 0$. The idea of arriving at a notion of horizontality on TP, adapted to the presence of the anchor map in the picture, lies at the basis of the following concept, introduced in [1].

Definition 2. *A ρ-connection on μ is a linear bundle map $h : \mu^*V \to TP$ (over the identity on P), such that $\rho \circ p_V = T\mu \circ h$, where p_V is the projection of μ^*V onto V.*

There is a quite striking similarity between our first diagram and the one we can draw here for the illustration of all spaces involved in the definition of a ρ-connection:



Note that points in the image $\rho^1(T^\rho P)$ can be vertical in TP when the corresponding point in the domain is not vertical in $T^\rho P$ (because ρ need not be injective). This is related to the observation that $\operatorname{Im} h$ can have a non-empty intersection with the vertical vectors on P. As discussed in detail in [1], $\operatorname{Im} h$ will in general also fail to determine a full complement to the vertical vectors on P. That is why one refers to a ρ-connection on μ also as a 'generalised connection'.

The point we would like to emphasise, however, is that it is perhaps not such a good idea to concentrate on horizontality on TP. Instead, as one may conjecture from an inspection of the two diagrams above, the better fibration to look for horizontality in this framework is the prolonged bundle $\mu^1 : T^\rho P \to P$. In other words, we think it is important to bring the pictures of *ρ-prolongation* and *ρ-connection* together into the following scheme.



What we propose to discuss in detail now is that, given a ρ-connection on μ, there is an associated, genuine decomposition of the bundle μ^1, i.e. a 'horizontal subspace', at each point $p \in P$, of the fibre of $T^\rho P$, which is complementary to the vertical subspace at p. In other words, instead of considering a horizontal lift operation from sections of τ to sections of τ_P, as is done in [1], it is more appropriate to focus on a horizontal lift from sections of τ, and by extension sections of the pullback bundle p, to sections of the bundle μ^1.

The fibre linear map $j : T^\rho P \to \mu^* V : (v, Q) \mapsto (\tau_P(Q), v)$ is surjective and its kernel is $\mathcal{V}^\rho P$. Therefore we have the following short exact sequence of vector bundles:

$$0 \to \mathcal{V}^\rho P \to T^\rho P \xrightarrow{j} \mu^* V \to 0, \tag{2}$$

where the second arrow is the natural injection.

Theorem 1. *The existence of a ρ-connection on μ is equivalent to the existence of a splitting ^{H} of the short exact sequence (2); we have $\rho^1 \circ {}^H = h$.*

Proof Let $h : \mu^* V \to TP$ be given and satisfy the requirements of a ρ-connection on μ. To define the 'horizontal lift' of a point $(p, v) \in \mu^* V$, as a point in $T^\rho P$, it suffices to fix the projections of $(p, v)^H$ under ρ^1 and μ^2 in a consistent way. We put:

$$\rho^1\big((p, v)^H\big) := h(p, v) \qquad \text{and} \qquad \mu^2\big((p, v)^H\big) := v. \tag{3}$$

This determines effectively an element of $T^\rho P$ since $\rho\circ\mu^2((p,v)^{^H}) = \rho(v) = \rho\circ p_{_V}((p,v)) = T\mu\circ h((p,v)) = T\mu\circ\rho^1((p,v)^{^H})$. The horizontal lift is obviously a splitting of (2), since by construction $j\left((p,v)^{^H}\right) = (\tau_P(h(p,v)),v) = (p,v)$.

Conversely, if a splitting ${}^{^H}$ of (2) is given, we define $h : \mu^*V \to TP$ by $h(p,v) = \rho^1((p,v)^{^H})$. It satisfies the required properties, i.e. h is a linear bundle map and we have

$$T\mu\circ h = T\mu\circ\rho^1\circ{}^{^H} = \rho\circ\mu^2\circ{}^{^H} = \rho\circ p_{_V}\circ j\circ{}^{^H} = \rho\circ p_{_V},$$

which concludes the proof. □

Denoting the subbundle of $T^\rho P$ which is complementary to $\mathcal{V}^\rho P$ by $\mathcal{H}^\rho P$, it follows that

$$T^\rho P = \mathcal{H}^\rho P \oplus \mathcal{V}^\rho P. \tag{4}$$

An equivalent way of expressing this decomposition (analogous to what is familiar for the case of a classical Ehresmann connection) is the following: there exist two complementary projection operators $P_{_H}$ and $P_{_V}$ on $T^\rho P$, i.e. we have $P_{_H} + P_{_V} = id$, and

$${P_{_H}}^2 = P_{_H}, \qquad {P_{_V}}^2 = P_{_V}, \qquad P_{_H}\circ P_{_V} = P_{_V}\circ P_{_H} = 0.$$

As usual, (2) leads to an associated short exact sequence for the set of sections of these spaces, regarded as bundles over P:

$$0 \to Ver(\mu^1) \to Sec(\mu^1) \xrightarrow{j} Sec(p) \to 0, \tag{5}$$

where $Ver(\mu^1)$ denotes the set of vertical sections of μ^1. The same symbol j is used for this second interpretation, so that for $\mathcal{Z} \in Sec(\mu^1)$ and $p \in P$: $j(\mathcal{Z})(p) = j(\mathcal{Z}(p))$. Via the composition with μ, sections of τ can be regarded as maps from P to V and, as such, are (*basic*) sections of $p : \mu^*V \to P$. We will use the notations $P_{_V}$ and $P_{_H}$ also when we regard these projectors as acting on sections of μ^1, rather than points in $T^\rho P$.

Apart from the already mentioned applications to Lie algebroids [8, 21], it has recently been shown that ρ-connections can be an important tool in, for example, nonholonomic mechanics [12], sub-Riemannian geometry [13], Poisson geometry [9] and in control theory [14].

The case of linear ρ-connections

Assume now that $\mu : P \to M$ now is a *vector* bundle. Linearity of a connection is characterised in [1] by an invariance property of the map h under the flow of the dilation field on P. A more direct characterisation of linearity is the following. Let $\Sigma_\lambda : P\times_M P \to P$ denote the linear combination map: $\Sigma_\lambda(p_1,p_2) = p_1 + \lambda p_2$. A ρ-connection on μ is said to be *linear* if the map $h : \mu^*V \to TP$ has the property

$$h(p_1 + \lambda p_2, v) = T_{(p_1,p_2)}\Sigma_\lambda\left(h(p_1,v), h(p_2,v)\right), \tag{6}$$

for all $(p_1,p_2) \in P\times_M P$, $\lambda \in \mathbb{R}$ and $v \in V$.

As is shown in [1], any operator $\nabla : Sec(\tau) \times Sec(\mu) \to Sec(\mu)$ which is $\mathbb{R}$-bilinear and has the properties

$$\nabla_{f\zeta}\sigma = f\nabla_\zeta\sigma, \qquad \nabla_\zeta(f\sigma) = f\nabla_\zeta\sigma + \rho(\zeta)(f)\sigma, \tag{7}$$

for all $\zeta \in Sec(\tau)$, $\sigma \in Sec(\mu)$ and $f \in C^\infty(M)$, defines a unique *linear* ρ-connection on μ. As usual, the linearity of the covariant derivative operator ∇ in its first argument, implies that the value of $\nabla_\zeta\sigma$ at a point $m \in M$, only depends on the value ζ at m and thus gives rise to an operator $\nabla_v : Sec(\mu) \to P_{\tau(v)}$, for each $v \in V$, determined by

$$\nabla_v\eta := \nabla_\zeta\eta(m), \qquad \text{with} \quad \zeta(m) = v.$$

In order to come to a covariant derivative along curves and a rule of parallel transport, we make the following preliminary observation. Going back to the overall diagram, we see two ways to go from $T^\rho P$ to TP, namely the direct map ρ^1 and $h \circ j$. By definition, the image for both maps projects under $T\mu$ onto the same $\rho(v)$, so that the difference is a vertical vector at some point $p \in P$ which, when P is a vector bundle, can be identified with an element of $P_{\mu(p)}$. With these identifications understood, we eventually get a map from $T^\rho P$ to P which is called the connection map in [1] (by analogy with the connection map in [23]). Let us summarise this by writing simply

$$K := \rho^1 - h \circ j : T^\rho P \to P \tag{8}$$

(read: K is $\rho^1 - h \circ j$, when regarded as map from $T^\rho P$ into P). The following side observation is worth being made here. In the alternative concept of ρ-connections, as established by Theorem 1, it is clear that the connection map K is nothing but the vertical projector P_V, with a similar identification being understood (to be precise: the isomorphism between $\mathcal{V}^\rho_p P$ and $V_p P$, followed by the identification with $P_{\mu(p)}$ again). In fact this illustrates that the alternative view is superior to the one expressed by Definition 2, in the following sense. Once the importance of the space $T^\rho P$ is recognised, one can (in the present case that P is a vector bundle) define a vertical lift operation from $P_{\mu(p)}$ to $\mathcal{V}^\rho_p P$ in the usual way (see the next section for more details); it extends to sections of bundles over P, i.e. yields a vertical lift from sections of $\mu^* P \to P$ to $Sec(\mu^1)$. So, it is a matter of developing first these tangent bundle like features of the ρ-prolongation, after which all tools are available to discuss ρ-connections without ever needing the map h. This is the main merit of the approach taken in [21] and [19]. For the sake of further unifying both pictures, however, we will continue here to take advantage of the insight which is being offered by our overall diagram.

Let now $c : I \to V$ be a ρ-admissible curve, which means that $\dot{c}_M = \rho \circ c$, where $c_M = \tau \circ c$ is the projected curve in M. Consider further a curve $\psi : I \to P$ in P which projects on c_M, i.e. such that $\psi_M := \mu \circ \psi = c_M$. It follows that $T\mu \circ \dot{\psi} = \rho \circ c$, so that such a ψ actually gives rise to a curve in $T^\rho P$: $t \mapsto (c(t), \dot{\psi}(t))$. As a result, making use of the map K, we can obtain a new curve in P, which is denoted by $\nabla_c\psi$:

$$\nabla_c\psi(t) := K((c(t), \dot{\psi}(t))) = \dot{\psi}(t) - h((\psi(t), c(t))), \tag{9}$$

(the identification of P with VP being understood). If η is a section of μ and c is an admissible curve, then denoting by ψ the restriction of η to that curve, $\psi(t) = \eta(c_M(t))$, one can show that

$$\nabla_c \psi(t) = \nabla_{c(t)} \eta. \tag{10}$$

As can be readily seen from (9), given an admissible curve c and a point $p \in P$, finding a curve ψ in P which starts at p and makes $\nabla_c \psi = 0$ is a well-posed initial value problem for a first-order ordinary differential equation, and hence gives rise to a unique solution. The solution is called the *horizontal lift* of c through p, denoted by c^h. Hence, we have

$$\nabla_c c^h = 0, \tag{11}$$

and points in the image of c^h are said to be obtained from p by *parallel transport along* c.

It is of some interest to rephrase what we have said at the beginning of the discussion on ρ-admissible curves: if $c : I \to V$ is ρ-admissible, then for every $\psi : I \to P$ which projects onto c_M, the curve $t \mapsto (c(t), \dot{\psi}(t))$ in fact is a ρ^1-admissible curve in $T^\rho P$. This idea can be pushed a bit further. Indeed, when thinking of curves in the context of our alternative view on ρ-connections, it is rather the following construction which looks like the natural thing to do.

Consider a curve γ in $\mu^* V$, i.e. γ is of the form $\gamma : t \mapsto (\psi(t), c(t))$, with $c : I \to V$ and $\psi : I \to P$, whereby the only assumption at the start is that $\psi_M = c_M$. Take its horizontal lift $\gamma^H : I \to T^\rho P$ which is defined, according to (3), by

$$t \mapsto \gamma^H(t) = \big(c(t), h(\psi(t), c(t))\big). \tag{12}$$

Then, we could define ψ to be c^h, the horizontal lift of c, if γ^H is a ρ^1-admissible curve in $T^\rho P$. Indeed, it is clear by construction that $\mu^1 \circ \gamma^H = \psi$, so that ρ^1-admissibility requires that $\dot{\psi} = \rho^1 \circ \gamma^H = h(\psi, c)$. Since $\psi_M = c_M$, this implies in particular that $\dot{c}_M = T\mu \circ \dot{\psi} = T\mu(h(\psi, c)) = \rho \circ c$. So, this alternative definition implies that c will necessarily have to be ρ-admissible. Furthermore, from comparing what ρ^1-admissibility means with (9) and (11), it is clear that we are talking then about the same concept of horizontal lift c^h.

Note, by the way, that this other way of defining c^h by no means relies on the assumption of linearity of the ρ-connection. So, it is perfectly possible to talk about parallel transport also in the context of non-linear connections. The difference then is, of course, that if we look at points of P in the image of curves c^h with different initial values in P_m, and this as a map between fibres of P, there need not be any special feature to talk about (compared to the fibre-wise linear action of this map we have in the case of a linear connection); also, if c has a given interval as domain, c^h need not be defined over the same domain. Needless to say, one can introduce such a generalisation also within the more traditional approach described first. Indeed, the map K makes sense for arbitrary ρ-connections and as a result one can introduce an operation $\nabla_\zeta \sigma$ also in this more general situation. This then still depends on the section ζ of V in a $C^\infty(M)$-linear way, but the fact that such a ∇ is not very commonly used comes from the failure of having a derivation property with respect to the module structure of $Sec(\mu)$.

3 The case of an affine bundle and its bidual

Suppose that $\pi : E \to M$ is an affine bundle, modelled on a vector bundle $\overline{\pi} : \overline{E} \to M$. For any $m \in M$, $E^\dagger_m := \mathrm{Aff}(E_m, \mathbb{R})$ is the set of all affine functions on E_m and $E^\dagger = \bigcup_{m\in M} E^\dagger_m$ is a vector bundle over M, called the extended dual of E. In turn, the dual of $\pi^\dagger : E^\dagger \to M$, denoted by $\tilde{\pi} : \tilde{E} := (E^\dagger)^* \to M$, is a vector bundle into which both E and $\overline{E}$ can be mapped via canonical injections, denoted respectively by ι and $\boldsymbol{\iota}$. The map ι is affine and has $\boldsymbol{\iota}$ as its associated linear map. With reference to the previous section, the situation we will focus on now is the case where $\mu : P \to M$ is the affine bundle $\pi : E \to M$, whereas $\tau : V \to M$ still is an arbitrary vector bundle. Our main objective is to define and characterise ρ-connections on π which are *affine*. For that purpose, we will need the overall diagram of the previous section also with the vector bundle $\tilde{\pi} : \tilde{E} \to M$ in the role of $\mu : P \to M$.

Definition 3. *A ρ-connection h on the affine bundle $\pi : E \to M$ is said to be affine, if there exists a linear ρ-connection $\tilde{h} : \tilde{\pi}^*V \to T\tilde{E}$ on $\tilde{\pi} : \tilde{E} \to M$ such that,*

$$\tilde{h} \circ \iota = T\iota \circ h.$$

Both sides in the above commutative scheme of course are regarded as maps from π^*V to $T\tilde{E}$, which means that the ι on the left stands for the obvious extension $\iota : \pi^*V \to \tilde{\pi}^*V, (e, v) \mapsto (\iota(e), v)$.

Probably the best way to see what this concept means is to look at a coordinate representation. Let x^i denote coordinates on M and y^α fibre coordinates on E with respect to some local frame $(e_0; \{\boldsymbol{e}_\alpha\})$ for $Sec(\pi)$. The induced basis for $Sec(\pi^\dagger)$ is denoted by (e^0, e^α) and defined as follows: for each $a \in Sec(\pi)$ with local representation $a(x) = e_0(x) + a^\alpha(x)\boldsymbol{e}_\alpha(x)$,

$$e^0(a)(x) = 1, \ \forall x, \qquad e^\alpha(a)(x) = a^\alpha(x).$$

In turn, we denote the dual basis for $Sec(\tilde{\pi})$ by (e_0, e_α) (so that in fact $\iota(e_0) = e_0$ and $\boldsymbol{\iota}(\boldsymbol{e}_\alpha) = e_\alpha$). Induced coordinates on $\tilde{E}$ are denoted by $(x^i, y^A) = (x^i, y^0, y^\alpha)$. For the coordinate representation of a point $v \in V$, we will typically write (x^i, v^a). The anchor map $\rho : V \to TM$ then takes the form $\rho : (x^i, v^a) \mapsto \rho^i_a(x)v^a \frac{\partial}{\partial x^i}$.

Following [1], we know that the map $h : \pi^*V \to TE$ locally is of the form:

$$h(x^i, y^\alpha, v^a) = (x^i, y^\alpha, \rho^i_a(x)v^a, -\Gamma^\alpha_a(x, y)v^a), \tag{13}$$

whereby we have adopted a different sign convention concerning the connection coefficients Γ^α_a. Similarly, $\tilde{h} : \tilde{\pi}^*V \to T\tilde{E}$, which is further assumed to be linear, takes the form

$$\tilde{h}(x^i, y^A, v^a) = (x^i, y^A, \rho^i_a(x)v^a, -\tilde{\Gamma}^A_{aB}(x)y^B v^a). \tag{14}$$

We have

$$\tilde{h}(\iota(e), v) = \Big(x^i, 1, y^\alpha, \rho^i_a(x)v^a, -(\tilde{\Gamma}^A_{a0}(x) + \tilde{\Gamma}^A_{a\beta}(x)y^\beta)v^a)\Big),$$

whereas

$$Ti \circ h(e, v) = \Big(x^i, 1, y^\alpha, \rho^i_a(x)v^a, 0, -\Gamma^\alpha_a(x, y)v^a\Big).$$

It follows that $\tilde{\Gamma}^0_{aB} = 0$ and, more importantly, that the connection coefficients of the affine ρ-connection h are of the form (omitting tildes)

$$\Gamma^\alpha_a(x,y) = \Gamma^\alpha_{a0}(x) + \Gamma^\alpha_{a\beta}(x)y^\beta. \tag{15}$$

Notice that $\overline{\pi} : \overline{E} \to M$ is a (proper) vector subbundle of $\tilde{\pi}$. With respect to the given anchor map, it of course also has its ρ-prolongation $T^\rho\overline{E}$. Taking the restriction of the linear ρ-connection $\tilde{h}$ to $\overline{\pi}^*V$, we get a linear ρ-connection $\overline{h}$ on $\overline{\pi}$, meaning that $\tilde{h} \circ \boldsymbol{\iota} = T\boldsymbol{\iota} \circ \bar{h}$. The above coordinate expressions make this very obvious. Indeed, if (x^i, w^α) are the coordinates of an element $\boldsymbol{w} \in \overline{E}$, we have

$$\begin{aligned}
\overline{h}(x^i, w^\alpha, v^a) &= \tilde{h}(x^i, 0, w^\alpha, v^a) \\
&= (x^i, 0, w^\alpha, \rho^i_a v^a, 0, -\Gamma^\alpha_{a\beta} w^\beta v^a) \quad \text{as element of } T\tilde{E} \\
&= (x^i, w^\alpha, \rho^i_a v^a, -\Gamma^\alpha_{a\beta} w^\beta v^a) \quad \text{as element of } T\overline{E}.
\end{aligned}$$

Note further that we can formally write for the coordinate expression of $h(e + \boldsymbol{w}, v)$:

$$\begin{aligned}
h(x^i, y^\alpha + w^\alpha, v^a) &= \left(x^i, y^\alpha + w^\alpha, \rho^i_a v^a, -(\Gamma^\alpha_{a0} + \Gamma^\alpha_{a\beta} y^\beta)v^a - \Gamma^\alpha_{a\beta} w^\beta v^a\right) \\
&= h(x^i, y^\alpha, v^a) + \overline{h}(x^i, w^\alpha, v^a).
\end{aligned}$$

But this is more than just a formal way of writing: the following intrinsic construction which generalises (6) is backing it. Let Σ denote the action of $\overline{E}$ on E which defines the affine structure, i.e. $\Sigma(e, \boldsymbol{w}) = e + \boldsymbol{w}$ for $(e, \boldsymbol{w}) \in E \times_M \overline{E}$. Then the above formal relation expresses that we have:

$$h(e + \boldsymbol{w}, v) = T_{(e,\boldsymbol{w})}\Sigma\left(h(e, v), \overline{h}(\boldsymbol{w}, v)\right) \tag{16}$$

In fact, by reading the above coordinate considerations backwards, roughly speaking, one can see that (16), for a given linear $\overline{h}$, will imply that the connection coefficients of the ρ-connection h have to be of the form (15). In other words, the following is an equivalent definition of affineness of h.

Definition 4. *A ρ-connection h on the affine bundle $\pi : E \to M$ is affine, if there exists a linear ρ-connection $\overline{h} : \overline{\pi}^*V \to T\overline{E}$ on $\overline{\pi} : \overline{E} \to M$, such that (16) holds for all $(e, \boldsymbol{w}) \in E \times_M \overline{E}$.*

One can then construct an extension $\tilde{h} : \tilde{\pi}^*V \to T\tilde{E}$, which coincides with $\overline{h}$ when restricted to $\overline{\pi}^*V$, by requiring that $\tilde{h}$ be linear and satisfy $\tilde{h} \circ \iota = T\iota \circ h$.

As shown in Theorem 1, a ρ-connection on π is equivalent to a decomposition of the bundle $T^\rho E$, originating from a horizontal lift operation from π^*V to $T^\rho E$ (or sections thereof). In the representation (1) of points of $T^\rho E$ as couples of an element of V and a suitable tangent vector of E, the horizontal lift is given by

$$(x^i, y^\alpha, v^a)^H = \left((x^i, v^a), v^a\left(\rho^i_a \frac{\partial}{\partial x^i} - \Gamma^\alpha_a \frac{\partial}{\partial y^\alpha}\right)\right).$$

At this stage, it is of interest to introduce a local basis for sections of the ρ-prolongation $\pi^1 : T^\rho E \to E$. A natural choice, adapted to the choice of a local frame in $Sec(\pi)$, the natural basis of $\mathcal{X}(E)$ and the choice of a local basis of sections $\boldsymbol{v}_a$ of τ, is determined as follows: for each $e \in E$, if x are the coordinates of $\pi(e) \in M$,

$$\mathcal{X}_a(e) = \left(\boldsymbol{v}_a(x),\, \rho_a^i(x)\frac{\partial}{\partial x^i}\bigg|_e\right), \quad \mathcal{V}_\alpha(e) = \left(0,\, \frac{\partial}{\partial y^\alpha}\bigg|_e\right). \tag{17}$$

Coordinates of a point $(v, X_e) \in T^\rho E$ are of the form: $(x^i, y^\alpha, v^a, X^\alpha)$. A general section of the ρ-prolongation can be represented locally in the form:

$$\mathcal{Z} = \zeta^a(x,y)\mathcal{X}_a + Z^\alpha(x,y)\mathcal{V}_\alpha. \tag{18}$$

Its projection onto $Sec(p)$ ($p : \pi^* V \to E$) is $\zeta = \zeta^a \boldsymbol{v}_a$. Now, once we have a given ρ-connection on π (affine or not), we are led to introduce a local basis for the horizontal sections of π^1, which is given by

$$\mathcal{H}_a = P_{_H}(\mathcal{X}_a) = \mathcal{X}_a - \Gamma_a^\alpha(x,y)\mathcal{V}_\alpha. \tag{19}$$

A better representation of the section (18), adapted to the given connection, then becomes:

$$\mathcal{Z} = \zeta^\alpha \mathcal{H}_a + (Z^\alpha + \Gamma_b^\alpha \zeta^b)\mathcal{V}_\alpha. \tag{20}$$

Let us repeat that, as a result of Theorem 1 and Definition 4, the existence of an affine ρ-connection on π is equivalent to the existence of a horizontal lift from $Sec(p)$ to $Sec(\pi^1)$, giving rise to a direct sum decomposition (4), and which is such that, in coordinates, the connection coefficients (19) are of the form (15).

We next turn our attention to the concept of connection map, and want to see for the particular case of an affine ρ-connection, to what extent it gives rise also to a covariant derivative operator and a notion of parallel transport.

When considering the ρ-prolongation of different bundles P, it is convenient to indicate the dependence on P also in the map ρ^1. Given a ρ-connection h on the affine bundle $\pi : E \to M$, the map $\rho_E^1 - h \circ j : T^\rho E \to TE$ gives rise (as before) to a vertical tangent vector to E, at the point e say. As such, this vector can be identified with an element of $\overline{E}$, the vector bundle on which E is modelled, at the point $\pi(e)$. With the same notational simplification as before, we thus get a connection map

$$K := \rho_E^1 - h \circ j : T^\rho E \to \overline{E}. \tag{21}$$

K of course also extends to a map from $Sec(\pi^1)$ to $Sec(\overline{\pi})$. It follows directly from the definition that we have

$$K(\mathcal{H}_a) = 0, \qquad K(\mathcal{V}_\alpha) = \boldsymbol{e}_\alpha. \tag{22}$$

We wish to come back here in some more detail to the relation between the map K and the vertical projector $P_{_V} = id - P_{_H}$, coming from the direct sum decomposition of $T^\rho E$. In the present case of an affine bundle $\pi : E \to M$ over a vector bundle $\overline{\pi} : \overline{E} \to M$, there

is a natural vertical lift operation from $\overline{E}_m$ to T_eE for each $e \in E_m$. It is determined by: $\boldsymbol{w} \mapsto w_e^V$, where for each $f \in C^\infty(E)$,

$$w_e^V(f) = \frac{d}{dt} f(e + t\boldsymbol{w})\bigg|_{t=0}.$$

This in turn extends to an operator ${}^V : \pi^*\overline{E} \to T^\rho E$, determined by $(e, \boldsymbol{w})^V = (0, w_e^V)$, which defines an isomorphism between $\pi^*\overline{E}$ and $\operatorname{Im} P_V$. The short exact sequence (2) of which a ρ-connection is a splitting, can thus be replaced by

$$0 \to \pi^*\overline{E} \xrightarrow{V} T^\rho E \xrightarrow{j} \mu^* V \to 0. \tag{23}$$

Within this picture of ρ-connections, the connection map K thus is essentially the co-splitting of the splitting ^{H}, that is to say, we have $K \circ {}^V = id_{\pi^*\overline{E}}$ and ${}^V \circ K + {}^H \circ j = id_{T^\rho E}$.

The map K becomes more interesting when the connection is affine. Indeed, denoting the projection of $T^\rho\tilde{E}$ onto $\tilde{\pi}^*V$ by $\tilde{j}$, it then follows from Definition 4 that we also have a connection map

$$\tilde{K} := \rho^1_{\tilde{E}} - \tilde{h} \circ \tilde{j} : T^\rho\tilde{E} \to \tilde{E}. \tag{24}$$

The map $T\iota : TE \to T\tilde{E}$ extends to a map from $T^\rho E$ to $T^\rho\tilde{E}$ in the following obvious way: $T\iota : (v, X_e) \mapsto (v, T\iota(X_e))$. Indeed, we have $T\tilde{\pi}(T\iota(X_e)) = T(\tilde{\pi} \circ i)(X_e) = T\pi(X_e) = \rho(v)$, as required.

Proposition 1. *For an affine ρ-connection on π we have*

$$\boldsymbol{\iota} \circ K = \tilde{K} \circ T\iota. \tag{25}$$

Proof In coordinates, K and $\tilde{K}$ are given by

$$\begin{aligned} K : &\quad (x^i, v^a, y^\alpha, Z^\alpha) \mapsto (Z^\alpha + \Gamma^\alpha_a v^a)\, \boldsymbol{e}_\alpha(x) \\ \tilde{K} : &\quad (x^i, v^a, y^A, Z^A) \mapsto Z^0 e_0(x) + (Z^\alpha + \Gamma^\alpha_{aB} y^B v^a)\, e_\alpha(x). \end{aligned}$$

Hence,

$$\begin{aligned} \tilde{K} \circ T\iota(x^i, v^a, y^\alpha, Z^\alpha) &= \tilde{K}(x^i, v^a, 1, y^\alpha, 0, Z^\alpha) \\ &= \left(Z^\alpha + (\Gamma^\alpha_{a0} + \Gamma^\alpha_{a\beta} y^\beta) v^a\right) e_\alpha(x), \end{aligned}$$

from which the result follows in view of (15). □

Notice that $\overline{h}$ also has a corresponding connection map $\overline{K} : T^\rho\overline{E} \to \overline{E}$, which obviously coincides with $\tilde{K}|_{T^\rho\overline{E}}$, so that we also have

$$\boldsymbol{\iota} \circ \overline{K} = \tilde{K} \circ T\boldsymbol{\iota}. \tag{26}$$

Let now ζ be a section of τ and σ a section of π. If we apply the tangent map $T\sigma : TM \to TE$ to $\rho(\zeta(m))$, it is obvious by construction that $\big(\zeta(m), T\sigma(\rho(\zeta(m)))\big)$ will be an element of $T^\rho E$. The connection map K maps this into a point of $\overline{E}|_m$. Hence, the covariant

derivative operator of interest in this context is the map $\nabla : Sec(\tau) \times Sec(\pi) \to Sec(\overline{\pi})$, defined by

$$\nabla_\zeta \sigma(m) = K\big(\zeta(m), T\sigma(\rho(\zeta(m)))\big). \tag{27}$$

To discover the properties which uniquely characterise the covariant derivative associated to an affine ρ-connection, we merely have to exploit the results of Proposition 2. In doing so, we will of course rely on the known properties (see [1]) of the covariant derivative $\tilde{\nabla}$, associated to the linear ρ-connection $\tilde{h}$. We observe that ∇ is manifestly $\mathbb{R}$-linear in its first argument and now further look at its behaviour with respect to the $C^\infty(M)$-module structure on $Sec(\tau)$. From (25), it follows that for $f \in C^\infty(M)$,

$$\begin{aligned}
\boldsymbol{\iota}\big((\nabla_{f\zeta}\sigma)(m)\big) &= \boldsymbol{\iota}\Big(K\big(f\zeta(m), T\sigma(\rho(f\zeta(m)))\big)\Big) \\
&= \tilde{K}\big(f\zeta(m), T(\iota\sigma)(\rho(f\zeta(m)))\big) \\
&= \tilde{\nabla}_{f\zeta}(\iota\sigma)(m) = f(m)\,\tilde{\nabla}_\zeta(\iota\sigma)(m) \\
&= f(m)\tilde{K}\big(\zeta(m), T\iota \circ T\sigma(\rho(\zeta(m)))\big) \\
&= f(m)\,\boldsymbol{\iota}\Big(K\big(\zeta(m), T\sigma(\rho(\zeta(m)))\big)\Big) \\
&= \boldsymbol{\iota}\big(f(m)\,\nabla_\zeta\sigma(m)\big),
\end{aligned}$$

from which it follows that

$$\nabla_{f\zeta}\sigma = f\,\nabla_\zeta\sigma. \tag{28}$$

For the behaviour in the second argument, we replace σ by $\sigma + f\boldsymbol{\eta}$, with $f \in C^\infty(M)$ and $\boldsymbol{\eta} \in Sec(\overline{\pi})$. Denoting the linear covariant derivative coming from the restriction $\overline{K}$ by $\overline{\nabla}$, we compute in the same way, using (25) and (26):

$$\begin{aligned}
\boldsymbol{\iota}\big(\nabla_\zeta(\sigma + f\boldsymbol{\eta})(m)\big) &= \boldsymbol{\iota}\Big(K\big(\zeta(m), T(\sigma + f\boldsymbol{\eta})(\rho(\zeta(m)))\big)\Big) \\
&= \tilde{K}\big(\zeta(m), T(\iota\sigma + f\boldsymbol{\iota\eta})(\rho(\zeta(m)))\big) = \tilde{\nabla}_\zeta(\iota\sigma + f\boldsymbol{\iota\eta})(m) \\
&= \tilde{\nabla}_\zeta\iota\sigma(m) + f(m)\big(\tilde{\nabla}_\zeta\boldsymbol{\iota\eta}\big)(m) + \rho(\zeta)(f)(m)\,\boldsymbol{\iota\eta}(m) \\
&= \tilde{K}\big(\zeta(m), T\iota \circ T\sigma(\rho(\zeta(m)))\big) + f(m)\,\tilde{K}\big(\zeta(m), T\boldsymbol{\iota} \circ T\boldsymbol{\eta}(\rho(\zeta(m)))\big) \\
&\quad + \rho(\zeta)(f)(m)\,\boldsymbol{\iota\eta}(m) \;=\; \boldsymbol{\iota}\Big(K\big(\zeta(m), T\sigma(\rho(\zeta(m)))\big)\Big) \\
&\quad + f(m)\,\boldsymbol{\iota}\Big(\overline{K}\big(\zeta(m), T\boldsymbol{\eta}(\rho(\zeta(m)))\big)\Big) + \rho(\zeta)(f)(m)\,\boldsymbol{\iota\eta}(m) \\
&= \boldsymbol{\iota}\Big(\nabla_\zeta\sigma(m) + f(m)\,\overline{\nabla}_\zeta\boldsymbol{\eta}(m) + \rho(\zeta)(f)(m)\boldsymbol{\eta}(m)\Big).
\end{aligned}$$

This expresses that we have the property:

$$\nabla_\zeta(\sigma + f\boldsymbol{\eta}) = \nabla_\zeta\sigma + f\,\overline{\nabla}_\zeta\boldsymbol{\eta} + \rho(\zeta)(f)\,\boldsymbol{\eta}. \tag{29}$$

In coordinates we have, for $\zeta = \zeta^a(x)\boldsymbol{v}_a$ and $\sigma = e_0 + \sigma^\alpha(x)\boldsymbol{e}_\alpha$:

$$\nabla_\zeta\sigma = \left(\frac{\partial\sigma^\alpha}{\partial x^i}\rho^i_a(x) + \Gamma^\alpha_{a0}(x) + \Gamma^\alpha_{a\beta}(x)\sigma^\beta(x)\right)\zeta^a(x)\,\boldsymbol{e}_\alpha. \tag{30}$$

As one can see, the linearity in ζ makes that the value of $\nabla_\zeta\sigma$ at a point m only depends of the value of ζ at m, so that the usual extension works, whereby for any fixed $v \in V$, ∇_v is a map from $Sec(\mu)$ to $\overline{E}_m$, defined by $\nabla_v\sigma = \nabla_\zeta\sigma(m)$, for any ζ such that $\zeta(m) = v$.

Theorem 2. *An affine ρ-connection h on π is uniquely characterised by the existence of an operator $\nabla : Sec(\tau) \times Sec(\pi) \to Sec(\overline{\pi})$ and an associated $\overline{\nabla} : Sec(\tau) \times Sec(\overline{\pi}) \to Sec(\overline{\pi})$, such that ∇ is $\mathbb{R}$-linear in its first argument, $\overline{\nabla}$ satisfies the requirements for the determination of a linear ρ-connection on $\overline{\pi}$, and the properties (28) and (29) hold true.*

Proof Given an affine ρ-connection h on π, the existence of operators ∇ and $\overline{\nabla}$ with the required properties has been demonstrated above. Assume conversely that such operators are given. Then, there exists an extension $\tilde{\nabla} : Sec(\tau) \times Sec(\tilde{\pi}) \to Sec(\tilde{\pi})$, which is defined as follows. Every $\tilde{\sigma} \in Sec(\tilde{\pi})$ locally is either of the form $\tilde{\sigma} = f\,\iota(\sigma)$ for some $\sigma \in Sec(\pi)$ or of the form $\tilde{\sigma} = \boldsymbol{\iota}(\boldsymbol{\eta})$ for some $\boldsymbol{\eta} \in Sec(\overline{\pi})$. In the first case, we put

$$\tilde{\nabla}_\zeta\tilde{\sigma} = f\,\boldsymbol{\iota}(\nabla_\zeta\sigma) + \rho(\zeta)(f)\iota(\sigma);$$

in the second case, we put

$$\tilde{\nabla}_\zeta\tilde{\sigma} = \boldsymbol{\iota}(\overline{\nabla}_\zeta\boldsymbol{\eta}).$$

We further impose $\tilde{\nabla}$ to be $\mathbb{R}$-linear in its second argument. $\mathbb{R}$-linearity as well as $C^\infty(M)$-linearity in the first argument trivially follows from the construction. It is further easy to verify that for $g \in C^\infty(M)$: $\tilde{\nabla}_\zeta(g\tilde{\sigma}) = g\,\tilde{\nabla}_\zeta\tilde{\sigma} + \rho(\zeta)(g)\,\tilde{\sigma}$. Indeed, in the case that $\tilde{\sigma} = f\,\iota(\sigma)$, for example, we have

$$\begin{aligned}\tilde{\nabla}_\zeta(g\tilde{\sigma}) &= gf\,\boldsymbol{\iota}(\nabla_\zeta\sigma) + \big(f\,\rho(\zeta)(g) + g\,\rho(\zeta)(f)\big)\iota(\sigma)\\ &= g\,\tilde{\nabla}_\zeta\tilde{\sigma} + \rho(\zeta)(g)\,\tilde{\sigma},\end{aligned}$$

and likewise for the other case. Following [1] we thus conclude that $\tilde{\nabla}$ uniquely determines a linear ρ-connection on $\tilde{\pi}$ by the following construction: for each $(\tilde{e}, v) \in \tilde{\pi}^*V$, take any $\tilde{\psi} \in Sec(\tilde{\pi})$ for which $\tilde{\psi}(\tau(v)) = \tilde{e}$, and put

$$\tilde{h}(\tilde{e}, v) = T\tilde{\psi}(\rho(v)) - (\tilde{\nabla}_v\tilde{\psi})^V_{\tilde{e}},$$

where the last term stands for the element $\tilde{\nabla}_v\tilde{\psi}(\tau(v)) \in \tilde{E}_{\tau(v)}$, vertically lifted to a vector tangent to the fibre of $\tilde{E}$ at $\tilde{e}$.

Likewise, we define a fibre linear map $h : \pi^*V \to TE$ by

$$h(e, v) = T\psi(\rho(v)) - (\nabla_v\psi)^V_e,$$

which can be seen to be independent of the choice of a section ψ for which $\psi(\tau(v)) = e$. It is obvious that h satisfies the requirements of a ρ-connection on π. It remains to show that $\tilde{h} \circ \iota = T\iota \circ h$. We have

$$\begin{aligned}\tilde{h}(\iota(e), v) &= T(\iota\psi)(\rho(v)) - \big(\tilde{\nabla}_v(\iota\psi)\big)^V_{\iota(e)}\\ &= T(\iota\psi)(\rho(v)) - (\boldsymbol{\iota}\nabla_v\psi)^V_{\iota(e)}\\ &= T\iota \circ T\psi(\rho(v)) - T\iota\big((\nabla_v\psi)^V_e\big)\\ &= T\iota(h(e, v)),\end{aligned}$$

which completes the proof. □

Another interesting question one can raise in this context is about the circumstances under which a linear ρ-connection $\tilde{h}$ on $\tilde{\pi}$ is associated to an affine ρ-connection h on π in the sense of Definition 4. A simple look at coordinate expressions leads to the following result with a global meaning.

Proposition 2. *A linear ρ-connection on $\tilde{\pi}$ is associated to an affine ρ-connection on π if and only if e^0 is parallel.*

PROOF For the covariant derivative operator $\tilde{\nabla}$ associated to a linear $\tilde{h}$, we have for the local basis of $Sec(\tilde{\pi})$:

$$\tilde{\nabla}_\zeta e_A = \zeta^a \tilde{\Gamma}^B_{aA}\, e_B,$$

and by duality, for the basis of $Sec(\pi^\dagger)$:

$$\tilde{\nabla}_\zeta e^A = -\zeta^a \tilde{\Gamma}^A_{aB}\, e^B.$$

It follows that $\tilde{\nabla}_\zeta e^0 = 0 \;\Leftrightarrow\; \tilde{\Gamma}^0_{aB} = 0$. The restriction of $\tilde{h}$ to $\iota(E)$ then defines an affine ρ-connection on π. □

A few words are in order, finally, about the concept of parallel transport in this case. Following the comments about ρ^1-admissibility of a curve γ^H made at the end of the previous section, we know that a curve ψ in E, with coordinate representation $t \mapsto (x^i(t), \psi^\alpha(t))$ will be the horizontal lift c^h of a ρ-admissible curve $c : t \mapsto (x^i(t), c^a(t))$, provided that (cf. the coordinate expressions (13) and (15)) $x^i(t)$ and $\psi^\alpha(t)$ satisfy the differential equations:

$$\dot{x}^i = \rho^i_a(x)c^a(t), \tag{31}$$

$$\dot{\psi}^\alpha = -\Gamma^\alpha_{a0}(x)c^a(t) - \Gamma^\alpha_{a\beta}(x)c^a(t)\,\psi^\beta. \tag{32}$$

In the more standard approach to the definition of c^h, if c is a ρ-admissible curve in V and ψ a curve in E which projects onto c_M, we can define a new curve $\nabla_c\psi$ by a formula which is formally identical to (9). Note, however, that $\nabla_c\psi$ is a curve in $\overline{E}$ now. Nevertheless, it makes perfect sense to say that ψ in E is c^h if the associated curve $\nabla_c\psi$ in $\overline{E}$ is zero for all t. It can be seen from the coordinate expression (32) that for different initial values in a fixed fibre of E, we get an affine action between the affine fibres of E, whose corresponding linear part comes from the parallel transport rule associated to the linear connection $\overline{h}$ on $\overline{E}$. This is in agreement with the property, coming from (29), that

$$\nabla_\zeta(\sigma + \boldsymbol{\eta}) = \nabla_\zeta\sigma + \overline{\nabla}_\zeta\boldsymbol{\eta}. \tag{33}$$

4 Discussion

We have focussed our main efforts on understanding in detail what affineness of a ρ-connection means. This may seem like a purely mathematical issue, but we will argue now why such a concept has a lot of relevance also for physical applications. Perhaps the

simplest example of the natural appearance of an affine ρ-connection (though for a trivial ρ), is the following. Take E to be the first-jet bundle J^1M of a manifold M which is fibred over $\mathbb{R}$, and $V = TM$ with $\rho = id_{TM}$. Then $T^\rho E = TE$ and we are in the situation which has been extensively studied in [7]. It is well-known that every second-order differential equation field (SODE) on J^1M, say

$$\Gamma = \frac{\partial}{\partial t} + v^i \frac{\partial}{\partial x^i} + f^i(t, x, v) \frac{\partial}{\partial v^i}, \tag{34}$$

defines a non-linear connection whose connection coefficients are

$$\Gamma^i_j = -\frac{1}{2} \frac{\partial f^i}{\partial v^j} \qquad \Gamma^i_0 = -f^i + \frac{1}{2} \frac{\partial f^i}{\partial v^j} v^j. \tag{35}$$

To say that the forces f^i are quadratic in the velocities, i.e. are of the form

$$f^i = f^i_0(t, x) + f^i_j(t, x) v^j + f^i_{jk}(t, x) v^j v^k, \tag{36}$$

is an invariant condition and clearly gives rise then to a connection of affine type, as discussed in the previous section. Examples of such systems in classical mechanics are abundant. In fact, every holonomic Lagrangian system, for which the transformation formulas to generalised coordinates are time-dependent and non-linear, falls into this category, with non-constant coefficients in the quadratic expressions (36). But there is more: even when the SODE is more general (and may represent an arbitrary time-dependent Newtonian system for example), there is an affine connection around. Indeed, it is well known (see e.g. [7]) that the SODE-connection (35) gives rise to a linear connection on the pullback bundle $\tilde{\pi} : \pi_1^{0*}TM \to J^1M$, where π_1^0 denotes the projection $J^1M \to M$. That is to say, there is a corresponding covariant derivative operator $\tilde{\nabla} : \mathcal{X}(J^1M) \times Sec(\tilde{\pi}) \to Sec(\tilde{\pi})$. The notations have been chosen here to relate to those of Section 3, which means for example that the tangent bundle $TJ^1M \to J^1M$ here plays the role of the arbitrary vector bundle $V \to M$ in the general set-up of that section (while the anchor map still is the identity). The point now is that the vector bundle $\tilde{\pi} : \pi_1^{0*}TM \to J^1M$ has $\pi_1^{0*}J^1M \to J^1M$ as an affine subbundle and we have $\tilde{\nabla}_X dt = 0$, for all vector fields X on J^1M. In accordance with Proposition 2, therefore, we know that there is an affine connection on $\pi_1^{0*}J^1M \to J^1M$, characterised by a covariant derivative operator ∇ which has the properties required by Theorem 2. This very general occurrence of an affine connection may be seen perhaps as having a rather hidden existence, but one should not forget that $\tilde{\nabla}$ and ∇ have been the essential tools for characterising qualitative features of general SODEs, such as linearisability (see [7]) or separability [2].

Admittedly, this full reservoir of examples of physical systems where an affine ρ-connection makes its appearance, corresponds to the case where ρ is trivial. However, since Weinstein's fundamental article [24], one knows of many interesting physical problems where a non-trivial anchor map occurs, namely in the context where one has the additional structure of a Lie algebroid. For this reason, we intend to study in a forthcoming paper a quite general situation of affine ρ-connections, where the same kind of extra structure is available. To that end we will take $\pi : E \to M$ to be a general affine bundle and let the vector bundle $\tau : V \to M$ be the bidual $\tilde{\pi} : \tilde{E} \to M$. In addition we will assume

that E comes equipped with an affine Lie algebroid structure (as studied for example in [22, 20]). This implies that the anchor map $\rho : V \to TM$ then also becomes the anchor of a (vector) Lie algebroid. As shown in [20], the prolonged bundle $T^\rho E$ inherits a Lie algebroid structure; moreover there is a canonical endomorphism on sections of $T^\rho E$, which is exactly the analogue of the vertical endomorphism on a first-jet bundle. Not surprisingly therefore, it is possible to define dynamical systems of Lagrangian type on such an affine Lie algebroid. Much of this has been explored already in the above cited papers, but the theory of affine connections and so-called pseudo-SODEs in that context still needs to be developed.

Acknowledgements. E. Martínez acknowledges partial financial support from CICYT grant BFM2000-1066-C03-01.

References

[1] F. Cantrijn and B. Langerock, Generalised connections over a bundle map, *Diff. Geom. Appl.*, (2002) to appear (math.DG/0201274).

[2] F. Cantrijn, W. Sarlet, A. Vandecasteele and E. Martínez, Complete separability of time-dependent second-order ordinary differential equations, *Acta Appl. Math.* **42** (1996) 309–334.

[3] J.F. Cariñena, Lie groupoids and algebroids in classical and quantum mechanics, In: *Symmetries in quantum mechanics and quantum optics*, University of Burgos (Spain) (1999) 67–81.

[4] J.F. Cariñena and E. Martínez, Lie algebroid generalization of geometric mechanics, In: *Lie algebroids and related topics in differential geometry*, Banach Center Publications **54** (2001), 201–215.

[5] J. Clemente-Gallardo, Applications of Lie algebroids in mechanics and control theory, In: *Nonlinear control in the new millenium*, F. Lamnabhi-Lagarrigur, W. Respondek and A. Isidori, eds., (Springer Verlag) (2000).

[6] M. Crampin, *Jet bundle techniques in analytical mechanics*, Quaderni del consiglio nazionale delle ricerche, gruppo nazionale di fisica matematica **47** (1995).

[7] M. Crampin, E. Martínez and W. Sarlet, Linear connections for systems of second-order ordinary differential equations, *Ann. Inst. H. Poincaré* **65** (1996) 223–249.

[8] R.L. Fernandes, Lie algebroids, holonomy and characteristic classes, *Adv. Math.*, (2002) to appear (math.DG/0007132).

[9] R.L. Fernandes, Connections in Poisson geometry I: holonomy and invariants, *J. Diff. Geom.* **54** (2000) 303–365.

[10] J. Grabowski, K. Grabowska and P. Urbański, Lie brackets on affine bundles, preprint (2002) (math.DG/0203112).

[11] P.J. Higgins and K. Mackenzie, Algebraic constructions in the category of Lie algebroids, *J. of Algebra* **129** (1990) 194–230.

[12] B. Langerock, Nonholonomic mechanics and connections over a bundle map, *J. Phys. A: Math. Gen.* **34** (2001) L609–L615.

[13] B. Langerock, A connection theoretic approach to sub-Riemannian geometry, *J. Geom. Phys.* (2002) to appear.

[14] B. Langerock, Geometric aspects of the Maximum Principle and lifts over a bundle map, preprint (2002).

[15] P. Libermann, Lie algebroids and Mechanics , *Arch. Math. (Brno)* **32** (1996) 147–162.

[16] K. Mackenzie, *Lie groupoids and Lie algebroids in differential geometry*, London Math. Soc. Lect. Note Series 124 (Cambridge Univ. Press) (1987).

[17] L. Mangiarotti and G. Sardanashvily, *Connections in classical and quantum field theory*, (World Scientific, River Edge, NJ) (2000).

[18] E. Martínez, Lagrangian Mechanics on Lie algebroids, *Acta. Appl. Math.* **67** (2001) 295–320.

[19] E. Martínez, Connection theory on Lie algebroids, preprint (2002).

[20] E. Martínez, T. Mestdag and W. Sarlet, Lie algebroid structures and Lagrangian systems on affine bundles, *J. Geom. Phys.* (2002) to appear (math.DG/0203178).

[21] M. Popescu and P. Popescu, Geometric objects defined by almost Lie structures, In: *Lie algebroids and related topics in differential geometry*, Banach Center Publications **54** (2001) 217–233.

[22] W. Sarlet, T. Mestdag and E. Martínez, Lie algebroid structures on a class of affine bundles, preprint (2002) (math.DG/0201264).

[23] J. Vilms, Connections on tangent bundles, *J. Diff. Geom.* **1** (1967) 235–243.

[24] A. Weinstein, Lagrangian Mechanics and groupoids, In: *Mechanics day (Waterloo, ON, 1992)*, *Fields Institute Communications* **7**, American Mathematical Society (1996) 207–231.