

A gentle (without chopping) approach to the full Kostant Toda–Lattice

Pantelis A. DAMIANOU and Franco MAGRI

Department of Mathematics and Statistics, University of Cyprus, 1678, Nicosia, Cyprus

Email: damianou@ucy.ac.cy

Department of Mathematics, University of Milano Bicocca, Via Corsi 58, I 20126 Milano, Italy

Email: magri@matapp.unimib.it

Abstract

In this paper we propose a new algorithm for obtaining the rational integrals of the full Kostant–Toda lattice. This new approach is based on a reduction of a bi–Hamiltonian system on $gl(n, \mathbf{R})$. This system was obtained by reducing the space of maps from Z_n to $GL(n, \mathbf{R})$ endowed with a structure of a pair of Lie–algebroids

1 Introduction

The Toda lattice is arguably the most fundamental and basic of all finite dimensional integrable systems. It has various intriguing connections with other parts of mathematics and physics.

The Hamiltonian of the Toda lattice is given by

$$H(q_1, \dots, q_N, p_1, \dots, p_N) = \sum_{i=1}^N \frac{1}{2} p_i^2 + \sum_{i=1}^{N-1} e^{q_i - q_{i+1}} . \quad (1)$$

Equation (1) is known as the classical, finite, non–periodic Toda lattice to distinguish the system from the many and various other versions, e.g., the relativistic, quantum, periodic etc. This system was investigated in [9], [10], [14], [18], [21], [22], [24] and numerous of other papers; see [4] for a more extensive bibliography.

Hamilton’s equations become

$$\begin{aligned} \dot{q}_j &= p_j \\ \dot{p}_j &= e^{q_{j-1} - q_j} - e^{q_j - q_{j+1}} . \end{aligned}$$

The system is integrable. One can find a set of independent functions $\{H_1, \dots, H_N\}$ which are constants of motion for Hamilton's equations. To determine the constants of motion, one uses Flaschka's transformation:

$$a_i = \frac{1}{2} e^{\frac{1}{2}(q_i - q_{i+1})} \quad , \quad b_i = -\frac{1}{2} p_i \quad . \quad (2)$$

Then

$$\begin{aligned} \dot{a}_i &= a_i (b_{i+1} - b_i) \\ \dot{b}_i &= 2(a_i^2 - a_{i-1}^2) \quad . \end{aligned} \quad (3)$$

These equations can be written as a Lax pair $\dot{L} = [B, L]$, where L is the Jacobi matrix

$$L = \begin{pmatrix} b_1 & a_1 & 0 & \cdots & \cdots & 0 \\ a_1 & b_2 & a_2 & \cdots & & \vdots \\ 0 & a_2 & b_3 & \ddots & & \\ \vdots & & \ddots & \ddots & & \vdots \\ \vdots & & & \ddots & \ddots & a_{N-1} \\ 0 & \cdots & & \cdots & a_{N-1} & b_N \end{pmatrix} \quad , \quad (4)$$

and B is the skew-symmetric part of L . This is an example of an isospectral deformation; the entries of L vary over time but the eigenvalues remain constant. It follows that the functions $H_i = \frac{1}{i} \text{tr} L^i$ are constants of motion.

Note that the Lax pair (4) has the form

$$\dot{L}(t) = [P L(t), L(t)]$$

where P denotes the projection onto the skew-symmetric part in the decomposition of L into skew-symmetric plus lower triangular.

The Toda lattice was generalized in several directions.

We mention the Bogoyavlensky–Toda lattices which generalize the Toda lattice (which corresponds to a root system of type A_n) to other simple Lie groups. This generalization is due to Bogoyavlensky [1]. These systems were

studied extensively in [15] where the solution of the systems was connected intimately with the representation theory of simple Lie groups.

Another generalization is due to Deift, Li, Nanda and Tomei [5] who showed that the system remains integrable when L is replaced by a full (generic) symmetric $n \times n$ matrix.

Another variation is the full Kostant-Toda lattice (FKT) [6], [11], [23]. We briefly describe the system: In [15] Kostant conjugates the matrix L in (4) by a diagonal matrix to obtain a matrix of the form

$$L = \begin{pmatrix} b_1 & 1 & 0 & \cdots & \cdots & 0 \\ a_1 & b_2 & 1 & \ddots & & \vdots \\ 0 & a_2 & b_3 & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & & 0 \\ \vdots & & & \ddots & \ddots & 1 \\ 0 & \cdots & \cdots & 0 & a_{n-1} & b_n \end{pmatrix}. \quad (5)$$

The equations take the form

$$\dot{X}(t) = [X(t), P X(t)]$$

where P is the projection onto the strictly lower triangular part of $X(t)$. This form is convenient in applying Lie theoretic techniques to describe the system.

To obtain the full Kostant-Toda lattice we fill the lower triangular part of L in (4) with additional variables. (P is again the projection onto the strictly lower part of $X(t)$). So, using the notation from [6], [11] and [23]

$$\dot{X}(t) = [X(t), P X(t)] ,$$

where X is in $\epsilon + B_-$ and $P X$ is in N_- . B_- is the Lie algebra of lower triangular matrices and N_- is the Lie algebra of strictly lower triangular matrices. The fixed matrix ϵ has a general form in terms of root systems:

$$\epsilon = \sum_{\alpha \in \Delta} x_{\alpha} ,$$

where Δ denotes the set of simple roots. In fact the FKT lattice itself can easily be generalized for each simple Lie group; see [2], [11].

In the case of $sl(4, \mathbf{C})$ the matrix X has the form

$$X = \begin{pmatrix} f_1 & 1 & 0 & 0 \\ g_1 & f_2 & 1 & 0 \\ h_1 & g_2 & f_3 & 1 \\ k_1 & h_2 & g_3 & f_4 \end{pmatrix}, \quad (6)$$

with $\sum_i f_i = 0$.

The functions $H_i = \frac{1}{i} \text{Tr } X^i$ are still in involution but they are not enough to ensure integrability. This is a crucial point: the existence of a Lax pair does not guarantee integrability. There are, however, additional integrals which are rational functions of the entries of X . The method used to obtain these additional integrals is called chopping and was used originally in [5] for the full symmetric Toda and later in [6] for the case of the full Kostant–Toda lattice.

In this paper we use a different method of obtaining these rational integrals which does not involve chopping. This method uses a reduction of a bi-Hamiltonian system on $gl(N, \mathbf{R})$, a system which was first defined in [19], [20]. In [20] A. Meucci presents the bi-Hamiltonian structure of Toda_3 , a dynamical system studied by Kupershmidt in [16] as a reduction of the KP hierarchy. Meucci derives this structure by a suitable restriction of the set of maps from Z_d to $GL(3, \mathbf{R})$, in the context of Lie algebroids.

In [19] the bi-Hamiltonian structure of the periodic Toda lattice is investigated by the reduction process described above using maps from Z_d to $GL(2, \mathbf{R})$. This approach parallels the work of [7] where the continuous analog of the Toda lattice is studied, namely the KdV. If instead the target space is $gl(3, \mathbf{R})$ one obtains the Boussineq hierarchy. The work of Meucci is a discrete version of this approach. If one generalizes the cases $N = 2, 3$ i.e. consider maps from Z_d to $GL(N, \mathbf{R})$ the resulting system will be denoted by Toda_N . In the present paper we use the results of [19], [20] as a starting point. We begin with the bi-Hamiltonian system obtained on $gl(N, \mathbf{R})$ in the particular case $d = N$ and use a further reduction to obtain the dynamics

of the full Kostant–Toda lattice. We then propose a new algorithm which produces all the rational integrals for the FKT lattice without using chopping. We present this algorithm by specific examples ($N = 3, 4$) in section 4. Sections 2 and 3 contain a general review of the old methods and results on integrability and bi–Hamiltonian structure of the FKT lattice.

2 Integrability of the FKT lattice

Let $\mathcal{G} = sl(n)$, the Lie algebra of $n \times n$ matrices of trace zero. Using the decomposition $\mathcal{G} = B_+ \oplus N_-$ we can identify B_+^* with the annihilator of N_- with respect to the Killing form. This annihilator is B_- . Thus we can identify B_+^* with B_- and therefore with $\epsilon + B_-$ as well.

The Lie Poisson bracket in the case of $sl(4, \mathbf{C})$ is given by the following defining relations:

$$\begin{aligned} \{g_i, g_{i+1}\} &= h_i, \\ \{g_i, f_i\} &= -g_i, \\ \{g_i, f_{i+1}\} &= g_i, \\ \{h_i, f_i\} &= -h_i, \\ \{h_i, f_{i+2}\} &= h_i, \\ \{g_1, h_2\} &= k_1, \\ \{g_3, h_1\} &= -k_1, \\ \{k_1, f_1\} &= -k_1, \\ \{k_1, f_4\} &= k_1. \end{aligned}$$

All other brackets are zero. Actually, we calculated the brackets on $gl(4, \mathbf{C})$; the trace of X now becomes a Casimir. The Hamiltonian in this bracket is $H_2 = \frac{1}{2} \text{Tr } X^2$.

Remark: If we use a more conventional notation for the matrix X , i.e. x_{ij} for $i \geq j$, $x_{ii+1} = 1$, and all other entries zero, then the bracket is simply

$$\{x_{ij}, x_{kl}\} = \delta_{li}x_{kj} - \delta_{jk}x_{il} . \quad (7)$$

The functions $H_i = \frac{1}{i} \text{Tr } X^i$ are still in involution but they are not enough to ensure integrability. There are, however, additional integrals and the in-

interesting feature of this system is that the additional integrals turn out to be rational functions of the entries of X . We describe the constants of motion following references [6], [10], [23].

For $k = 0, \dots, \lfloor \frac{n-1}{2} \rfloor$, denote by $(X - \lambda \text{Id})_{(k)}$ the result of removing the first k rows and last k columns from $X - \lambda \text{Id}$, and let

$$\det (X - \lambda \text{Id})_{(k)} = E_{0k} \lambda^{n-2k} + \dots + E_{n-2k,k} .$$

Set

$$\frac{\det (X - \lambda \text{Id})_{(k)}}{E_{0k}} = \lambda^{n-2k} + I_{1k} \lambda^{n-2k-1} + \dots + I_{n-2k,k} .$$

The functions I_{rk} , $r = 1, \dots, n - 2k$, are constants of motion for the FKT lattice.

Example We consider in detail the $gl(3, \mathbf{C})$ case:

$$X = \begin{pmatrix} f_1 & 1 & 0 \\ g_1 & f_2 & 1 \\ h_1 & g_2 & f_3 \end{pmatrix} .$$

Taking $H_2 = \frac{1}{2} \text{tr} X^2$ as the Hamiltonian, and the above Poisson bracket

$$\dot{x} = \{H_2, x\}$$

gives the following equations:

$$\begin{aligned} \dot{f}_1 &= -g_1 \\ \dot{f}_2 &= g_1 - g_2 \\ \dot{f}_3 &= g_2 \\ \dot{g}_1 &= g_1(f_1 - f_2) - h_1 \\ \dot{g}_2 &= g_2(f_2 - f_3) + h_1 \\ \dot{h}_1 &= h_1(f_1 - f_3) \end{aligned}$$

Note that $H_1 = f_1 + f_2 + f_3$ while $H_2 = \frac{1}{2}(f_1^2 + f_2^2 + f_3^2) + g_1 + g_2$. These equations can be written in Lax pair form, $\dot{X} = [B, X]$, by taking

$$B = \begin{pmatrix} 0 & 0 & 0 \\ g_1 & 0 & 0 \\ h_1 & g_2 & 0 \end{pmatrix} .$$

The chopped matrix is given by

$$\begin{pmatrix} g_1 & f_2 - \lambda \\ h_1 & g_2 \end{pmatrix} .$$

The determinant of this matrix is $h_1\lambda + g_1g_2 - h_1f_2$ and we obtain the rational integral

$$I_{11} = \frac{g_1g_2 - h_1f_2}{h_1} . \quad (8)$$

Note that the phase space is six dimensional, we have two Casimirs (H_1, I_{11}) and the functions (H_2, H_3) are enough to ensure integrability.

Another Example In the case of $gl(4, \mathbf{C})$ the additional integral is

$$I_{21} = \frac{g_1g_2g_3 - g_1f_3h_2 - f_2g_3h_1 + h_1h_2}{k_1} + f_2f_3 - g_2 .$$

and

$$I_{11} = \frac{g_1h_2 + g_3h_1}{k_1} - f_2 - f_3$$

is a Casimir.

In this example the phase space is ten dimensional, we have two Casimirs (H_1, I_{11}) and the functions (H_2, H_3, H_4, I_{21}) are independent and pairwise in involution.

3 bi-Hamiltonian structure

We recall the definition and basic properties of master symmetries following Fuchssteiner [13]. Consider a differential equation on a manifold M defined by a vector field χ . We are mostly interested in the case where χ is a Hamiltonian vector field. A vector field Z is a symmetry of the equation if

$$[Z, \chi] = 0.$$

A vector field Z is called a master symmetry if

$$[[Z, \chi], \chi] = 0,$$

but

$$[Z, \chi] \neq 0.$$

Master symmetries were first introduced by Fokas and Fuchssteiner in [12] in connection with the Benjamin-Ono Equation.

A bi-Hamiltonian system is defined by specifying two Hamiltonian functions H_1 , H_2 and two Poisson tensors π_1 and π_2 , that give rise to the same Hamiltonian equations. Namely, $\pi_1 \nabla H_2 = \pi_2 \nabla H_1$. The notion of bi-Hamiltonian system was introduced in [17] in 1978.

We now return to the FKT lattice. We want to define a second bracket π_2 so that H_1 is the Hamiltonian and

$$\pi_2 \nabla H_1 = \pi_1 \nabla H_2 .$$

i.e. we want to construct a bi-Hamiltonian pair. We will achieve this by finding a master symmetry X_1 so that

$$X_1(\text{Tr } X^i) = i \text{Tr } X^{i+1} .$$

We construct X_1 by considering the equation

$$\dot{X} = [Y, X] + X^2 . \tag{9}$$

We choose Y in such a way that the equation is consistent. One solution is

$$Y = \sum_{i=1}^n \alpha_i E_{ii} + \sum_{i=1}^{n-1} \beta_i E_{i,i+1} ,$$

where

$$\beta_i = i,$$

$$\alpha_i = i f_i + \sum_{k=1}^{i-1} f_k .$$

The vector field X_1 is defined by the right hand side of (9).

For example, in $gl(4, \mathbf{C})$ the components of X_1 are:

$$\begin{aligned}
X_1(f_1) &= 2g_1 + f_1^2, \\
X_1(f_2) &= 3g_2 + f_2^2, \\
X_1(f_3) &= -g_2 + 4g_3 + f_3^2, \\
X_1(f_4) &= -2g_3 + f_4^2, \\
X_1(g_1) &= 3h_1 + g_1f_1 + 3g_1f_2, \\
X_1(g_2) &= 4h_2 + 4g_2f_3, \\
X_1(g_3) &= -h_2 - g_3f_3 + 5g_3f_4, \\
X_1(h_1) &= g_1g_2 + 4k_1 + h_1f_1 + h_1f_2 + 4h_1f_3, \\
X_1(h_2) &= g_2g_3 + h_2f_3 + 5h_2f_4, \\
X_1(k_1) &= g_3h_1 + g_1h_2 + k_1f_1 + k_1f_2 + k_1f_3 + 5k_1f_4.
\end{aligned}$$

The second bracket π_2 is defined by taking the Lie derivative of π_1 in the direction of X_1 . The bracket π_2 is at most quadratic i.e. in the case $n = 4$

$$\begin{aligned}
\{g_i, g_{i+1}\} &= g_i g_{i+1} + h_i f_{i+1}, \\
\{g_i, h_i\} &= g_i h_i, \\
\{g_{i+1}, h_i\} &= -g_{i+1} h_i, \\
\{g_i, f_i\} &= -g_i f_i, \\
\{g_i, f_{i+1}\} &= g_i f_{i+1}, \\
\{g_i, f_{i+2}\} &= h_i, \\
\{g_{i+1}, f_i\} &= -h_i, \\
\{h_i, f_i\} &= -h_i f_i, \\
\{h_i, f_{i+2}\} &= h_i f_{i+2}, \\
\{f_i, f_{i+1}\} &= g_i.
\end{aligned}$$

This bracket was first obtained in [3] using the method described above (i.e. master symmetries). A closed expression for this bracket was obtained later by Faybusovich and Gekhman in [8]. It was obtained using R-matrices and the expression takes the following simple form:

$$\{x_{ij}, x_{kl}\} = \begin{cases} \text{sign}(k-i)x_{ij}x_{kj} & \text{if } j = l \\ x_{ij}x_{il} & \text{if } k = i \\ x_{ij}x_{kl} + x_{il}x_{kj} & \text{if } i < k \leq j \\ x_{il} & \text{if } k = j + 1. \end{cases} \quad (10)$$

As we will see in the next section, there is a linear and a quadratic bracket on $gl(n, \mathbf{R})$ whose restriction to the FKT lattice coincides with the brackets π_1, π_2 .

4 A new approach

As we mentioned in the introduction, our starting point is the work of A. Meucci [19], [20]. His work is a discrete analogue of a procedure that produces the KdV, Boussineq and Gelfand–Dickey hierarcies [7]. For example, the KdV is bi–Hamiltonian and it can be obtained by reducing the space of C^∞ maps from S^1 to $gl(2, \mathbf{R})$. If one considers the space of maps from S^1 to $gl(3, \mathbf{R})$ the Boussineq hierarchy is obtained. In [19], [20] the discrete version of the procedure is considered. The circle, S^1 , is replaced by the cyclic group \mathbf{Z}_d and one considers maps from \mathbf{Z}_d to $GL(N, \mathbf{R})$. One obtains, after reduction, equations that are bi–Hamiltonian. In the case $N = 2$ the resulting system is the periodic Toda lattice and for $N = 3$ a system studied by Kupershmidt in [16]. We will not get into the details of the procedure but rather we will use the results as our starting point for our own purposes. We will content with a short outline of the constructions of [19] and [20]. The basic object is the space of maps from \mathbf{Z}_d to $GL(N, \mathbf{R})$. This space is endowed with a structure consisting of a Poisson manifold together with a pair of Lie–algebroids suitably related. The next step is a Marsden–Ratiu type of reduction and the result is a bi–Hamiltonian system with a pair of Poisson stuctures on $gl(N, \mathbf{R})$. The Lax pair of the resulting system contains two spectral parameters and the theory of Gelfand–Zakarevich applies. The system turns out to be integrable with the required number of integrals. We give explicit formulas that we have computed from [19] in the case $N = 3$ both for the pair of Poisson brackets, the Lax pair and the integrals of motion. In the case $N = 4$ we display the Lax pair and the polynomial integrals of motion. To obtain the FKT lattice one has to perform a further reduction to the phase space of the FKT lattice and to obtain the rational integrals we propose a new algorithm which is the central new result of our paper. We illustrate with two examples:

Example 1: The $gl(3, \mathbf{C})$ case

The phase space of the system obtained by the procedure of Meucci is nine dimensional, i.e. matrices of the form

$$\begin{pmatrix} f_1 & h_2 & g_3 \\ g_1 & f_2 & h_3 \\ h_1 & g_2 & f_3 \end{pmatrix} .$$

We compute the pair of Poisson brackets on the extended space with variables

$$\{f_1, f_2, f_3, g_1, g_2, g_3, h_1, h_2, h_3\} .$$

The Lie–Poisson bracket is defined by the following structure matrix

$$\begin{pmatrix} 0 & 0 & 0 & -g_1 & 0 & g_3 & -h_1 & h_2 & 0 \\ 0 & 0 & 0 & g_1 & -g_2 & 0 & 0 & -h_2 & h_3 \\ 0 & 0 & 0 & 0 & g_2 & -g_3 & h_1 & 0 & -h_3 \\ g_1 & -g_1 & 0 & 0 & -h_1 & h_3 & 0 & 0 & 0 \\ 0 & g_2 & -g_2 & h_1 & 0 & -h_2 & 0 & 0 & 0 \\ -g_3 & 0 & g_3 & -h_3 & h_2 & 0 & 0 & 0 & 0 \\ h_1 & 0 & -h_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -h_2 & h_2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -h_3 & h_3 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} ,$$

and the quadratic Poisson bracket is defined by $A - A^t$ where A is the matrix

$$\begin{pmatrix} 0 & g_1 & -g_3 & -g_1 f_1 & h_1 - h_2 & g_3 f_1 & -h_1 f_1 & h_2 f_1 & 0 \\ 0 & 0 & g_2 & g_1 f_2 & -g_2 f_2 & h_2 - h_3 & 0 & -h_2 f_2 & h_3 f_2 \\ 0 & 0 & 0 & h_3 - h_1 & g_2 f_3 & -g_3 f_3 & h_1 f_3 & 0 & -h_3 f_3 \\ 0 & 0 & 0 & 0 & -h_1 f_2 - g_1 g_2 & h_3 f_1 + g_1 g_3 & -g_1 h_1 & 0 & g_1 h_3 \\ 0 & 0 & 0 & 0 & 0 & -h_2 f_3 - g_2 g_3 & g_2 h_1 & -g_2 h_2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & g_3 h_2 & -g_3 h_3 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} .$$

The Lax matrix with two spectral parameters is given by

$$L_{\lambda, \mu} = \begin{pmatrix} (f_1 + \lambda)\mu^2 & h_2 - \mu^3 & \mu g_3 \\ \mu g_1 & (f_2 + \lambda)\mu^2 & h_3 - \mu^3 \\ h_1 - \mu^3 & \mu g_2 & (f_3 + \lambda)\mu^2 \end{pmatrix} .$$

Let $p(\lambda, \mu) = \det L_{\mu, \lambda}$. Write

$$p(\lambda, \mu) = -\mu^9 + c_2(\lambda)\mu^6 + c_1(\lambda)\mu^3 + c_0(\lambda) .$$

Then

$$c_2(\lambda) = \lambda^3 + k_2 \lambda^2 + k_1 \lambda + k_0$$

where

$$\begin{aligned} k_2 &= f_1 + f_2 + f_3 \\ k_1 &= f_1 f_2 + f_1 f_3 + f_2 f_3 + g_1 + g_2 + g_3 \\ k_0 &= f_1 f_2 f_3 + f_1 g_2 + f_2 g_3 + f_3 g_1 + h_1 + h_2 + h_3 \end{aligned}$$

and

$$c_1(\lambda) = l_1 \lambda + l_0$$

where

$$\begin{aligned} l_1 &= -h_3 g_2 - g_3 h_1 - h_2 g_1 \\ l_0 &= -f_1 g_2 h_3 + g_1 g_3 g_2 - g_1 h_2 f_3 - h_1 g_3 f_2 - h_2 h_3 - h_3 h_1 - h_1 h_2 . \end{aligned}$$

Finally,

$$c_0(\lambda) = h_1 h_2 h_3 .$$

The functions k_i, l_i, c_0 are all in involution in the Lie–Poisson bracket. The functions k_2, l_1, c_0 are all Casimirs.

In the quadratic bracket k_i, l_i , and c_0 are all in involution. k_0, l_0 and c_0 are Casimirs.

Let

$$l = \frac{l_0}{l_1} = \frac{-f_1 g_2 h_3 + g_1 g_3 g_2 - g_1 h_2 f_3 - h_1 g_3 f_2 - h_2 h_3 - h_3 h_1 - h_1 h_2}{-h_3 g_2 - g_3 h_1 - h_2 g_1} .$$

Setting $h_2 = h_3 = 0$ in l we obtain I_{11} (??).

Example 2: The $gl(4, \mathbf{C})$ case

In order to give a complete comparison of the previous and the present method of obtaining the rational invariants we consider first integrability using chopping.

We consider the matrix L given by

$$L = \begin{pmatrix} f_1 & -1 & 0 & 0 \\ g_1 & f_2 & -1 & 0 \\ h_1 & g_2 & f_3 & -1 \\ k_1 & h_2 & g_3 & f_4 \end{pmatrix} .$$

Note that we are using $-\epsilon$ instead of ϵ in order to get the integrals to match exactly.

In this case the chopped matrix has the form

$$Ch_1(\lambda) = \begin{pmatrix} g_1 & f_2 - \lambda & -1 \\ h_1 & g_2 & f_3 - \lambda \\ k_1 & h_2 & g_3 \end{pmatrix} .$$

The characteristic polynomial has the form

$$k_1 \lambda^2 + (g_1 h_2 + h_1 g_3 - k_1 f_2 - k_1 f_3) \lambda + g_1 g_3 g_2 - g_1 h_2 f_3 - h_1 f_2 g_3 - h_1 h_2 + k_1 f_2 f_3 + k_1 g_2 .$$

We obtain the following two rational invariants

$$i_{11} = \frac{h_2 g_1 + g_3 h_1}{k_1} - (f_2 + f_3) ,$$

and

$$i_{21} = \frac{g_1 g_3 g_2 - g_1 h_2 f_3 - h_1 f_2 g_3 - h_1 h_2}{k_1} + f_2 f_3 + g_2 .$$

We now turn to the gentle approach i.e. integrability without chopping. Consider the following Lax pair with two spectral parameters:

$$L_{\lambda, \mu} = \begin{pmatrix} (f_1 + \lambda)\mu^3 & k_2 - \mu^4 & h_3\mu & g_4\mu^2 \\ g_1\mu^2 & (f_2 + \lambda)\mu^3 & k_3 - \mu^4 & h_4\mu \\ h_1\mu & g_2\mu^2 & (f_3 + \lambda)\mu^3 & k_4 - \mu^4 \\ k_1 - \mu^4 & h_2\mu & g_3\mu^2 & (f_4 + \lambda)\mu^3 \end{pmatrix} . \quad (11)$$

Taking determinant we obtain the polynomial

$$p_{\lambda, \mu} = -\mu^{16} + K_3(\lambda)\mu^{12} + K_2(\lambda)\mu^8 + K_1(\lambda)\mu^4 + K_0(\lambda) .$$

We present the explicit expressions for the polynomials $K_i(\lambda)$.

•

$$K_3(\lambda) = K_{33}\lambda^3 + K_{32}\lambda^2 + K_{31}\lambda + K_{30} ,$$

where

$$K_{33} = f_1 + f_2 + f_3 + f_4 ,$$

$$K_{32} = g_1 + g_2 + g_3 + g_4 + f_1 f_2 + f_1 f_3 + f_1 f_4 + f_2 f_3 + f_2 f_4 + f_3 f_4 ,$$

$$K_{31} = h_1 + h_2 + h_3 + h_4 + f_1 f_2 f_3 + f_1 f_2 f_4 + f_1 f_3 f_4 + f_2 f_3 f_4 + f_2 g_4 \\ + f_3 g_4 + f_3 g_1 + f_4 g_1 + f_4 g_2 + f_2 g_3 + f_1 g_2 + f_1 g_3 ,$$

$$K_{30} = k_1 + k_2 + k_3 + k_4 + f_1 f_2 f_3 f_4 + g_1 g_3 + g_2 g_4 + f_2 h_3 + f_4 h_1 + \\ f_1 h_2 + f_3 h_4 + f_2 f_3 g_4 + f_3 f_4 g_1 + f_1 f_4 g_2 + f_1 f_2 g_3 .$$

•

$$K_2(\lambda) = K_{22}\lambda^2 + K_{21}\lambda + K_{20} ,$$

where

$$K_{22} = -h_2h_4 - g_2k_3 - g_3k_4 - h_1h_3 - g_1k_2 - k_1g_4 ,$$

$$\begin{aligned} K_{21} = & g_2h_4g_3 - g_1k_2f_4 - f_1h_2h_4 - k_1g_4f_3 - k_1f_2g_4 - f_1g_2k_3 - h_2h_4f_3 \\ & - g_1k_2f_3 + g_1g_2h_3 + h_1g_4g_3 + g_1h_2g_4 - f_1g_3k_4 - h_1f_2h_3 - g_2k_3f_4 \\ & - f_2g_3k_4 - h_2k_4 - h_2k_3 - h_1h_3f_4 - k_2h_4 - h_3k_4 - h_1k_2 - h_1k_3 \\ & - k_1h_3 - k_1h_4 , \end{aligned}$$

$$\begin{aligned} K_{20} = & -k_2k_4 - k_2k_3 - k_1k_2 - k_1k_3 - k_1k_4 - k_3k_4 - k_1f_2g_4f_3 \\ & + f_1g_2h_4g_3 - f_1h_2h_4f_3 - h_1k_2f_4 - f_1g_2k_3f_4 - f_1f_2g_3k_4 \\ & - g_1g_2g_4g_3 + g_1h_2g_4f_3 - h_1f_2h_3f_4 + g_1g_2h_3f_4 + h_1f_2g_4g_3 \\ & - g_1k_2f_3f_4 - f_1h_2k_3 - g_1k_2g_3 - f_1h_2k_4 - k_1g_2g_4 \\ & + h_1h_2g_4 - k_2h_4f_3 - f_2h_3k_4 + g_2h_3h_4 - g_2g_4k_3 \\ & - g_1g_3k_4 + g_1h_2h_3 - h_1k_3f_4 + h_1h_4g_3 - k_1f_2h_3 - k_1h_4f_3 . \end{aligned}$$

• $K_1(\lambda) = K_{11}\lambda + K_{10}$ where

$$K_{11} = h_1k_2k_3 + h_2k_3k_4 + k_1k_2h_4 + k_1h_3k_4 ,$$

$$\begin{aligned} K_{10} = & h_1k_2k_3f_4 + f_1h_2k_3k_4 - k_1g_2h_3h_4 + k_1g_2g_4k_3 + k_1k_2h_4f_3 \\ & + k_1f_2h_3k_4 - h_1h_2g_4k_3 + h_1h_2h_3h_4 - h_1k_2h_4g_3 - g_1h_2h_3k_4 \\ & + g_1k_2g_3k_4 + k_1k_2k_4 + k_1k_2k_3 + k_2k_3k_4 + k_1k_3k_4 . \end{aligned}$$

$$K_0(\lambda) = -k_1 k_2 k_3 k_4 .$$

Remark We note that $K_3(\lambda)$ gives the polynomial invariants. Clearly $H_1 = K_{33} = \text{trace } L$.

We also have $H_2 = \frac{1}{2}\text{trace}L^2 = \frac{1}{2}K_{33}^2 - K_{32}$ and $H_3 = \frac{1}{3}\text{trace}L^3 = \frac{1}{3}K_{33}^3 - K_{33}K_{32} + K_{31}$.

Finally, $H_4 = \frac{1}{4}K_{33}^4 + K_{33}K_{31} + \frac{1}{2}K_{32} - K_{30} - K_{33}K_{32}$.

These last relations hold provided that $k_2 = k_3 = k_4 = 0$, $h_3 = h_4 = 0$ and $g_4 = 0$.

Remark The next coefficient $K_2(\lambda)$ gives the rational invariants:

We form the quotient $\frac{K_{21}}{K_{22}}$ and set $k_2 = k_3 = k_4 = 0$ and $h_3 = h_4 = 0$. We obtain

$$\frac{K_{21}}{K_{22}} = \frac{-g_4(k_1 f_3 + k_1 f_2 - h_1 g_3 - g_1 h_2)}{-k_1 g_4} = f_2 + f_3 - \frac{(h_1 g_3 + g_1 h_2)}{k_1} .$$

This is precisely $-i_{11}$.

Similarly, we form $\frac{K_{20}}{K_{22}}$ to obtain precisely i_{21} .

Remark The last two terms, namely $K_1(\lambda)$ and $K_0(\lambda)$ become identically zero once we set $k_2 = k_3 = k_4 = 0$, $h_3 = h_4 = 0$.

In the general case, the polynomial $p_{\lambda,\mu}$ involves polynomials $A(\lambda)$, $B_1(\lambda), \dots, B_k(\lambda)$ and $C_1(\lambda), \dots, C_s(\lambda)$ with $k = \lfloor \frac{(n-1)}{2} \rfloor$ and $s = n - k - 1$. The polynomial $A(\lambda)$ can be used to obtain the polynomial integrals. The polynomials $B_i(\lambda)$ give the rational integrals using the procedure described above and the $C_j(\lambda)$ vanish identically once we restrict to the phase space of the FKT lattice.

References

- [1] O. I. Bogoyavlensky, *Comm. Math. Phys.* **51** (1976), 201.
- [2] Constantinides, K., Generalized full Kostant–Toda lattices, Master thesis, Department of Mathematics and Statistics, University of Cyprus, (2003)
- [3] Damianou P. A. , Paschalis P. , Sophocleous C., A tri-Hamiltonian formulation of the full Kostant-Toda lattice, *Letters in Math. Phys.*, **34**, 17–24 (1995).
- [4] P. A. Damianou, Multiple Hamiltonian structure of Bogoyavlensky–Toda lattices, *Reviews in Math. Phys.*, **16**, 175–241, (2004).
- [5] Deift, P.A., Li, L.C., Nanda, T., Tomei, C., The Toda lattice on a generic orbit is integrable, *Comm. Pure Appl. Math.* **39**, 183-232 (1986).
- [6] Ercolani N.M., Flaschka H., Singer S., The Geometry of the full Kostant-Toda Lattice, Colloque Verdier, Progress in Mathematics Series, Birkhaeuser Verlag, 181-225, (1994).
- [7] Falqui, G., Magri, F. and Pedroni, M.: bihamiltonian geometry, Darboux coverings, and linearization of the KP hierarchy, *Comm. Math. Phys.* **197** 303–324 (1998)
- [8] Faybusovich L. and Gekhtman. M.: Elementary Toda orbits and integrable lattices. *J. Math. Phys.*, **41** 2905-2921, (2000)
- [9] Flaschka, H., On the Toda lattice, *Phys. Rev.* **9**, 1924-1925 (1974).
- [10] Flaschka H.: On the Toda lattice II. Inverse-scattering solution, *Progr. Theor. Phys.* **51**, 703–716 (1974)
- [11] Flaschka, H., Integrable systems and torus actions. Lecture notes. University of Arizona (1992).
- [12] Fokas, A.S. and Fuchssteiner B., The Hierarchy of the Benjamin-Ono equations, *Phys. Lett.* **86A** 341-345 (1981).

- [13] Fuchssteiner, B. Mastersymmetries and higher order time-dependent symmetries and conserved densities of nonlinear evolution equations. *Progr. Theor. Phys.* **70**, 1508-1522 (1983).
- [14] Henon M.: Integrals of the Toda lattice, *Phys. Rev. B* **9**, 1921–1923 (1974)
- [15] Kostant, B., The solution to a generalized Toda lattice and representation theory, *Adv. Math.* **34**, 195-338 (1979)
- [16] B. Kupershmidt, *Asterisque* **123**, (1985)
- [17] Magri, F., A simple model of the integrable Hamiltonian equation. *J. Math. Phys.* **19**, 1156-1162 (1978)
- [18] Manakov S.: Complete integrability and stochastization of discrete dynamical systems, *Zh. Exp. Teor. Fiz.* **67**, 543–555 (1974)
- [19] Meucci A. : Compatible Lie algebroids and the periodic Toda lattice, *J. Geom. and Phys.* **35** 273–287 (2000)
- [20] Meucci A. : Toda Equations, bi-Hamiltonian Systems, and compatible Lie Algebroids, *Math. Phys. Analysis and Geom.*, **4** 131–146 (2001)
- [21] Moser J.: Finitely many mass points on the line under the influence of an exponential potential—an integrable system, *Lect. Notes Phys.* **38**, 97–101 (1976)
- [22] Moser J.: Three integrable Hamiltonian systems connected with isospectral deformations, *Adv. Math.* **16**, 197–220 (1975)
- [23] Singer S., The Geometry of the full Toda Lattice, Ph.D. Dissertation, Courant Institute, (1991).
- [24] Toda M.: One-dimensional dual transformation, *J. Phys. Soc. Japan* **22**, 431–436 (1967)