

# Imbrex Geometries

Jeroen Schillewaert \*

Department of Mathematics  
Imperial College  
London, England  
jschillewaert@gmail.com

Hendrik Van Maldeghem

Department of Mathematics  
Ghent University  
Ghent, Belgium  
hvm@cage.ugent.be

## Abstract

We introduce an axiom on strong parapolar spaces of diameter 2, which arises naturally in the framework of Hjelmslev geometries. This way, we characterize the Hjelmslev-Moufang plane and its relatives (line Grassmannians, certain half-spin geometries and Segre geometries). At the same time we provide a more general framework for a lemma of Cohen, which is widely used to study parapolar spaces. As an application, if the geometries are embedded in a projective space, we provide a common characterization of (projections of) Segre varieties, line Grassmann varieties, half-spin varieties of low rank, and the exceptional variety  $\mathcal{E}_{6,1}$  by means of a local condition on tangent spaces.

## 1 Introduction

Springer and Veldkamp [12] introduced the *Hjelmslev-Moufang planes* as geometries resembling Hjelmslev planes (because lines can meet in more than one point) and Moufang planes (because the coordinatizing structure is also an octonion algebra, though split, whereas the usual Moufang projective planes are defined over non-split ones). In the present paper we push the analogy a little bit further, using the more modern notion of *parapolar spaces*. Indeed, although the Hjelmslev-Moufang planes are not Hjelmslev planes themselves, we show that they satisfy a far more intuitive property of “realistic geometry”, as Hjelmslev [5] himself was aiming at, and that property follows from the definition of Hjelmslev planes, but it does not characterize Hjelmslev planes. The basic observation made by Hjelmslev was that, if one draws lines “close” to each other (meaning that the sharp angle they define is very small), then it is hard to identify the intersection point, and it looks as if the lines have a little segment in common. Dually, if two points are very close to each other, then the joining line is hard to identify. Hjelmslev,

---

\*Research supported by Marie Curie IEF grant GELATI (EC grant nr 328178)

and later Klingenberg [7], included these observations in their geometries by introducing a neighbor relation which precisely indicates when two points and two lines are close to one another. However, in the approach of Hjelmlev and Klingenberg, the neighbor relation is an equivalence relation, and the equivalence classes define an ordinary projective or affine plane (the “underlying” plane). A consequence of this is the following intuitive property:

(Imb) *Let  $x$  be a point not neighboring two neighboring points  $y_1, y_2$ . Then there are unique lines  $L_i$  joining  $x$  and  $y_i$ ,  $i = 1, 2$ , and  $L_1$  is a neighbor of  $L_2$ .*

The proof is very simple: in the underlying plane,  $y_1$  and  $y_2$  define the same point, distinct from the point defined by  $x$ , hence the lines  $L_1$  and  $L_2$  must define the same lines and are consequently contained in the same equivalence class.

We propose to take (Imb) as an axiom and combine this with the theory of parapolar spaces, where we want to view the symplecta as the lines of our geometry. In the approach of Hjelmlev, any two points are joined by at least one line; hence we consider strong parapolar spaces of diameter 2. Next, one must define the notion of “neighboring”. To do this, we refer back to the observation of Hjelmlev: if two points define a unique line, then these points are far enough from one another. Translated to the framework of parapolar spaces, two points at distance 2 define a unique symplecton. Consequently it is natural to define that two points are neighboring if they are collinear in the parapolar space. For the dual, we take into account the residual nature of parapolar spaces associated to building geometries. More precisely, in the situations we are interested in, the residue in a point is again a parapolar space, and with the definition of neighboring points above, the neighbor relation on points is preserved under taking residues (i.e., if  $y_1, y_2$  are two (non-)neighboring points collinear to a point  $x$ , then in the residue at  $x$ , the points defined by  $y_1$  and  $y_2$  are (non-)neighboring). So we also want the neighboring relation between symplecta to be preserved under taking residues. The only sensible way to define two symplecta to be neighbors then is when they intersect in a maximal singular subspace. This also implies that we should consider parapolar spaces of constant symplectic rank.

As a side remark, we mention that, in fact, the situation just described resembles in a certain sense better the reality than was the case with the Hjelmlev and Klingenberg planes. Indeed, in our case the neighbor relation is certainly not transitive, and this is more realistic: if one draws a number of points consecutively close to each other, then the first point may well be far from the last one.

Now we translate the property (Imb) to the framework of parapolar spaces with the neighboring relation as just derived. So let  $x$  be a point of a parapolar space of diameter 2, then  $y_1, y_2$  are two collinear points (collinearity in the parapolar space), both at distance 2 from  $x$ . If there was a point on the line  $y_1y_2$  collinear to  $x$ , then the symplecta through  $x, y_1$  and  $x, y_2$ , which we denote by  $\xi(x, y_1)$  and  $\xi(x, y_2)$ , respectively, would coincide, and hence be neighboring trivially.

So we may assume that every point of the line  $y_1y_2$  is at distance 2 from  $x$ . Then the symplecta  $\xi(x, y_1)$  and  $\xi(x, y_2)$  intersect in a maximal singular subspace of both of them.

In conclusion, in this paper we will study strong parapolar spaces with diameter 2 satisfying the following additional property, which we again refer to as (Imb), as an abbreviation of “Imbrex”.

(Imb) *Let  $x$  be a point not collinear with any point of the line  $L$ . Let  $y_1, y_2$  be distinct points on  $L$ . Then the symplecta  $\xi(x, y_1)$  and  $\xi(x, y_2)$  intersect in a singular subspace, which is maximal for both of them.*

In various places we shall use the fact that all maximal singular subspaces of a polar space have the same dimension. To ensure this we add the axiom that every sequence of nested singular subspaces is finite (it is the close analogue of the axiom for polar spaces ensuring finite rank, and all main examples satisfy it), and we call these parapolar spaces *imbrex geometries*. In fact, if the symplectic rank is at least 3, then one can classify imbrex geometries using a rather powerful theorem of Cohen & Cooperstein [3, 4] as updated by Shult [10]. The merit of Property (Imb), however, lies in the fact that it allows to study the symplectic rank 2 case in a more general and conceptual way than was done before. Moreover, it is ready-made to generalize the characterization of Segre varieties in [15] to the other varieties in the second row of the split version of the extended Freudenthal-Tits Magic Square (FTMS), i.e., line Grassmannians of projective spaces and the variety associated to buildings of type  $E_6$ , i.e., the variety associated to Springer and Veldkamp’ Hjelmslev-Moufang planes (in contrast, the restricted FTMS just contains the Segre variety of two projective planes, and the line Grassmannian of projective 5-space).

The paper is organized as follows. After introducing the preliminaries in Section 2, we investigate in Section 3 imbrex geometries of symplectic rank 2. We show that, if the symplecta are thick generalized quadrangles, then the maximal singular subspaces contain a lot of non-closed O’Nan configurations (the latter is a set of four lines such that each pair intersects in a point, except for exactly one pair). This immediately implies a fundamental lemma of Cohen [3], which was originally proved only for the case of classical generalized quadrangles, and later generalized by Shult and K. Thas [11] for all Moufang quadrangles (in fact, their proof shows that only the “strong transitivity” property of Moufang quadrangles is needed). In our setting, there is no restriction on the generalized quadrangles, and we also provide examples of imbrex geometries of symplectic rank 2 with thick symplecta (the maximal singular subspaces of these imbrex geometries are not isomorphic to projective spaces!). In this way, Cohen’s lemma for thick symplecta is turned into a positive result, rather than merely showing nonexistence. Also, our result reveals the true geometric reason why the thick case in Cohen’s setting cannot exist: it is not because the quadrangle has remarkable transitivity properties or enjoys the structure of pseudo-quadratic forms, but it is because in projective spaces all O’Nan configurations are closed (projective spaces are in fact characterized by that property).

In Section 4, we classify all imbrex geometries of symplectic rank at least 3. This will be an application of impressive work of Cohen and Cooperstein [3, 4].

In Section 5, we apply our results to the theory of *Mazzocca-Melone sets*. Roughly, a Mazzocca-Melone set is a set of points in a projective space satisfying a far reaching generalization of the conditions stated by Mazzocca & Melone [8] that originally characterized finite quadric Veronesean varieties. In [9] we provided a characterization of the Severi varieties over an arbitrary field. These precisely correspond to the varieties of the second row of the split version of the restricted FTMS using Mazzocca-Melone sets. We here propose to alter one of the axioms from a global condition to a local one, inspired by the characterization of all Segre varieties  $\mathcal{S}_{m,n}(\mathbb{K})$  (see [15]). The thus defined *local Mazzocca-Melone sets* will be classified in Section 5, and will entail a characterization of the varieties in the second row of the split version of the extended FTMS.

## 2 Preliminaries

In this section we briefly introduce polar and parapolar spaces thus fixing notation. More information about parapolar spaces can be found in [10].

### 2.1 Polar spaces

Polar spaces have been introduced by Veldkamp [18], later on included in the theory of buildings by Tits [16], and around the same time the axioms have been simplified by Buekenhout & Shult [2]. It is the latter point of view we take here.

Let  $\Gamma = (\mathcal{P}, \mathcal{L}, *)$  be a point-line geometry ( $\mathcal{P}$  is the set of points,  $\mathcal{L}$  the set of lines, and  $*$  a symmetric incidence relation). We will not consider geometries with repeated lines, so from now on we view  $\mathcal{L}$  as a subset of the power set of  $\mathcal{P}$ , and  $*$  is inclusion made symmetric. The *incidence graph* is the bipartite graph on  $\mathcal{P} \cup \mathcal{L}$  with  $*$  as adjacency relation. The dual of  $\Gamma$  is the point-line geometry  $(\mathcal{L}, \mathcal{P}, *)$ . A *subspace* of  $\Gamma$  is a subset  $S$  of the point set such that, if two points  $a, b$  belong to  $S$ , then all lines containing both  $a$  and  $b$  are contained in  $S$ . Points contained in a common line will be called *collinear*, dually, lines sharing at least one point are called *concurrent*. A *singular* subspace is a subspace every two points of which are collinear. Note that the empty set and a single point are legible singular subspaces. Now,  $\Gamma$  is called a *polar space of rank  $r$*  if the following conditions hold.

- (PS1) Every line contains at least 3 points.
- (PS2) No point is collinear with all other points.

- (PS3) Every nested sequence of singular subspaces has at most length  $r + 1$  and there exists such sequence of length  $r + 1$ .
- (PS4) For any point  $x$  and any line  $L$ , either one or all points on  $L$  are collinear with  $x$ .

A *generalized quadrangle* is a polar space of rank 2, the dual of a polar space of rank 2, or the point-line geometry defined by a  $(2 \times N)$ -grid (with  $N$  any cardinal number at least 2), or the dual of the latter. A generalized quadrangle is *thick* if it is both a polar space and a dual polar space, equivalently, if every element is incident with at least 3 other elements. If  $T \subseteq \mathcal{L}$ , then we denote by  $T^\perp$  the set of lines concurrent with every member of  $T$ . A pair of non-concurrent lines  $\{L, M\}$  of a generalized quadrangle is called *regular* if  $(\{L, M\}^\perp)^\perp = \{L', M'\}^\perp$ , for some distinct  $L', M' \in \{L, M\}^\perp$ . If for a line  $L$  every pair  $\{L, M\}$  with  $M$  non concurrent with  $L$  is regular, then the line  $L$  is called *regular*. A *subquadrangle* of the generalized quadrangle  $\Gamma = (\mathcal{P}, \mathcal{L}, *)$  is a subset of  $\mathcal{P}$ , together with a subset of  $\mathcal{L}$  on which  $*$  induces a generalized quadrangle (it is not necessarily a subspace!). A subquadrangle  $\Gamma'$  is *ideal* if for every point  $x$  in  $\Gamma'$ , every line of  $\Gamma$  incident with  $x$  also belongs to  $\Gamma'$ .

We note that polar spaces are partial linear spaces, i.e., two collinear points  $x, y$  determine exactly one line, which we sometimes denote by  $xy$ .

## 2.2 Parapolar spaces

Parapolar spaces were introduced to capture the spherical buildings of exceptional type (*spherical buildings* comprise projective spaces, polar spaces and the ones of exceptional type). Since we only need strong parapolar spaces of diameter 2, we only introduce these.

Let again  $\Gamma = (\mathcal{P}, \mathcal{L}, *)$  be a point-line geometry. A subspace  $S$  is called *convex* if for any pair of points  $\{x, y\} \subseteq S$ , every point contained in or incident with a line of any shortest path between  $x$  and  $y$  (in the incidence graph) is contained in  $S$ . Also,  $\Gamma$  is called *connected* if its incidence graph is connected. Now,  $\Gamma$  is called a *strong parapolar space of diameter 2* if the following two assertions hold:

- (PPS1)  $\mathcal{P}, \mathcal{L}$  is a connected space such that for every point  $x$  and every line  $L$  either 0, or exactly one or all points of  $L$  are collinear with  $x$  and all three possibilities occur.
- (PPS2) For every pair of distinct non-collinear points  $x$  and  $y$  in  $\mathcal{P}$ , the smallest convex subspace containing  $x$  and  $y$  is a polar space of rank at least 2.

The convex subspaces of (PPS2) are called the *symplecta*, or briefly, the *symps*, of  $\Gamma$ . By Axiom (PPS2), any two non-collinear points  $x, y \in \mathcal{P}$  are contained in a unique symp and we denote it by  $\xi(x, y)$ . Also, (PPS1) implies that there are at least two distinct symps.

Usually, the third axiom of a parapolar space reads

(PPS3) Every line is contained in at least one symplecton,

but this axiom is automatically satisfied in our case (we restricted to *strong* and *diameter 2*, see [9]).

For any integer  $k > 1$ , a parapolar space is said to have *symplectic rank  $k$*  (*symplectic rank at least  $k$* ) if and only if every symplecton has polar rank  $k$  (at least  $k$ ).

It follows from Corollary 13.3.3 of [10] that any symplecton is uniquely determined by any pair of noncollinear points contained in it. We will use this without further notice. From (PPS2) we immediately obtain the following two lemmas.

**Lemma 2.1 (The Quadrangle Lemma)** *Let  $L_1, L_2, L_3, L_4$  be four (not necessarily pairwise distinct) singular lines such that  $L_i$  and  $L_{i+1}$  (where  $L_5 = L_1$ ) share a (not necessarily unique) point  $p_i$ ,  $i = 1, 2, 3, 4$ , and suppose that  $p_1$  and  $p_3$  are not collinear. Then  $L_1, L_2, L_3, L_4$  are contained in a unique common symp.*

**Lemma 2.2** *Let  $p \in \mathcal{P}$  and let  $H$  be a symp not containing  $p$ . Then the set of points of  $H$  collinear with  $p$  constitutes a singular subspace of  $H$ .*

### 2.3 Imbrex geometries

We call a strong parapolar space with diameter 2 an *imbrex geometry* if it additionally satisfies the axiom (Imb) stated in the introduction. We now show that all symps have the same rank. This common rank will then be called the *symplectic rank* of the imbrex geometry.

**Lemma 2.3** *In an imbrex geometry, all symps have the same rank.*

**Proof** Let  $H, H'$  be two symps. We first note that (PPS2) implies that the graph of all symps, adjacent when they meet in a nonempty subspace, is connected. Hence we may assume that  $H \cap H'$  contains some point  $x$ . Let  $y \in H$  and  $y' \in H'$  be two points, both not collinear to  $x$ , and hence  $y' \notin H$ ,  $y \notin H'$ . If  $y$  and  $y'$  are collinear, then suppose that the line  $yy'$  contains a point  $z$  collinear to  $x$ . It follows that the symp  $\xi(x, y) = H$  contains  $z$  and hence  $y'$ , a contradiction. Hence by (Imb), the symps  $\xi(x, y) = H$  and  $\xi(x, y') = H'$  have the same rank. So we may assume that  $y$  and  $y'$  are not collinear. The symp  $\xi(y, y')$  shares a singular subspace  $S$  with  $H$  and a singular subspace  $S'$  with  $H'$ . We distinguish two possibilities.

1. Suppose the rank of  $\xi(y, y')$  is at least 3, or  $\xi(y, y')$  is a thick generalized quadrangle. Then it is easy to see that we can find a point  $z$  not in  $S \cup S'$ , and so not in  $H \cup H'$ , but collinear to both  $y, y'$ . Since  $x$  is not collinear to  $z$  by Lemma 2.1, as before, the lines  $yz$  and  $y'z$  do not contain any point collinear to  $x$ . Hence the symps  $\xi(x, y) = H$  and  $\xi(x, z)$  have the same rank, as well as  $\xi(x, y') = H'$  and  $\xi(x, z)$ .

2. Suppose  $\xi(y, y')$  is a non-thick generalized quadrangle, i.e., a grid. If we cannot find a point  $z$  as in the previous case, then the only possibility is that  $S$  and  $S'$  are disjoint lines of  $\xi(y, y')$ . Since both  $S$  and  $S'$  contain at least three points, collinearity defines a bijection between  $S$  and  $S'$ , and  $x$  is collinear to exactly one point of  $S$  and one point of  $S'$ , we can find collinear points  $u \in S$  and  $u' \in S'$  such that  $x$  is not collinear to either  $u$  or  $u'$ . Note that  $u = y$  and  $u' = y'$  are allowed. By (Imb), the symps  $\xi(x, u) = H$  and  $\xi(x, u') = H'$  intersect in a maximal singular subspace of both, implying  $H$  and  $H'$  have the same rank.  $\square$

### 3 Imbrex geometries of symplectic rank 2

Throughout this section, we let  $\Gamma = (\mathcal{P}, \mathcal{L}, *)$  be an imbrex geometry of symplectic rank 2. Our main aim is to show that, if at least one symp is thick, then every maximal singular subspace of  $\Gamma$  contains at least one *non-closed O'Nan configuration*, i.e., four distinct lines pairwise intersecting in a point, except for one pair, which is disjoint. But we will also determine the structure of  $\Gamma$  in case all symps are non-thick. Hence for the time being, we do not assume that at least one symp is thick.

**Lemma 3.1** *Let  $x, q_1, q_2 \in \mathcal{P}$  with  $q_1$  collinear to  $q_2$ . Suppose no point of the line  $q_1q_2$  is collinear to  $x$ . Then some point of the line  $L := \xi(x, q_1) \cap \xi(x, q_2)$  is collinear with all points on  $q_1q_2$ .*

**Proof** Note that  $L$  is indeed a line by (Imb). Now, in the generalized quadrangle  $\xi(x, q_i)$  there is a unique point  $z_i$  on  $L$  collinear to  $q_i$ ,  $i = 1, 2$ . If  $z_1 \neq z_2$ , then Lemma 2.1 yields  $\xi(x, q_1) = \xi(x, q_2)$ , implying that  $x$  is collinear with a point of  $L$ , a contradiction. Hence  $z_1 = z_2 \in L$  is collinear to both  $q_1$  and  $q_2$ . Axiom (PPS1) completes the proof.  $\square$

**Lemma 3.2** *Let  $L \in \mathcal{L}$  be arbitrary. Then the set  $B$  of points collinear with at least two distinct points of  $L$  is a maximal singular subspace of  $\Gamma$ . Conversely, every maximal singular subspace arises in this way.*

**Proof** First note that by (PPS1) a point of  $B$  is collinear to all points of  $L$ . Now let  $y_1, y_2 \in B$  be arbitrary. If  $y_1$  and  $y_2$  are collinear, then (PPS1) again implies that every point of the joining line belongs to  $B$ . Hence  $B$  is a subspace. There remains to show that  $y_1$  and  $y_2$  are collinear. Suppose not, then Lemma 2.1 yields a symp containing  $y_1, y_2$  and  $L$ , contradicting the fact that the symplectic rank is 2. Hence  $y_1$  is collinear with  $y_2$ , and  $B$  is singular. Clearly,  $B$  is maximal. Also, the converse is easy noting that every maximal singular subspace must contain a line since every point is contained in a line.  $\square$

**Lemma 3.3** *For each point  $r$ , there exists a line  $L$  no point of which is collinear to  $r$ .*

**Proof** Let  $r \in \mathcal{P}$  be arbitrary. Consider a symp  $H$  through  $r$  and a point  $y \notin H$  not collinear to  $r$  (this is possible since there are at least two symps, and clearly not all points of a second symp outside  $H$  can be collinear to  $r$ ). Also, by Lemma 2.2, not all points of  $H$  not collinear to  $r$  are collinear to  $y$ ; hence we find a point  $z \in H$  such that  $r, z$  and  $y$  are pairwise non-collinear. Consider two different lines  $L_1, L_2$  in  $\xi(y, z)$  through  $y$ . If  $r$  is collinear with points  $p_1, p_2$  on  $L_1, L_2$ , respectively, then  $r \in \xi(p_1, p_2) = \xi(y, z)$ , implying  $H = \xi(r, z) = \xi(y, z)$  contains  $y$ , a contradiction. Hence at least one of  $L_1$  or  $L_2$  contains no point collinear to  $r$ .  $\square$

We define a new point-line geometry  $\Delta = (\mathcal{P}, \mathcal{B}, \star)$ , where we call the elements of  $\mathcal{B}$  *blocks* to avoid confusion with the lines of  $\Gamma$ , where  $\mathcal{B}$  is the family of maximal singular subspaces of  $\Gamma$ , and where  $\star$  is containment made symmetric. We will need the following auxiliary results.

**Lemma 3.4** (i) *Two blocks intersect at most in one point.*

(ii) *A point  $p$  not contained in a block  $B$  is collinear with at least one point  $r \in B$ .*

**Proof** Clearly, (i) follows from Lemma 3.2.

For (ii) consider a line  $L \subseteq B$ . If  $p$  is collinear with a point of  $L$  we are done. Otherwise, by Lemma 3.1, there is a block  $B'$  containing  $L$  such that  $p$  is collinear with a point of  $B'$ . But by (i),  $B = B'$ , establishing the proof.  $\square$

**Lemma 3.5** (i) *The point-line geometry  $\Delta$  is a generalized quadrangle where each line contains at least three points.*

(ii) *Every symp is an ideal subquadrangle of  $\Delta$ . In particular, the symplecta are either all thick or all non-thick.*

**Proof** We show that  $\Delta$  is a polar space of rank 2. By Lemma 3.2 every line is contained in a block and (PS1), (PS2) and (PS3) follow easily. As for (PS4), consider a point  $x$  and a block  $B$  with  $x \notin B$ . If  $x$  were collinear to at least two points of  $B$ , then we would find a maximal subspace intersecting  $B$  in at least two points, a contradiction, hence by (ii) of Lemma 3.4  $x$  is collinear with exactly one point of  $B$ .

For (ii), consider a point  $x$  and a symp  $H$  through  $x$ . We need to show that each block  $B$  containing  $x$  intersects  $H$  in a line. Consider  $y \in H$  not collinear with  $x$ . By (ii) of Lemma 3.4,  $y$  is collinear with a point  $z \in B$  and hence it follows that  $z \in \xi(x, y) = H$ , establishing the proof (noting that every line, and hence also every block, contains at least three points, hence if a symp contains at least one point contained in three lines, then all points are contained in at least three lines, and the same holds for  $\Delta$ ).  $\square$

As a corollary, one can classify all imbrex geometries with symplectic rank 2 for which at least one symp is non-thick.



**Proposition 3.6** *If at least one symp is non-thick, then  $\Gamma$  is the product geometry of two linear spaces. Conversely, the product geometry of two arbitrary linear spaces is an imbrex geometry with symplectic rank 2, where the symps are the grids obtained by restriction to the product of two lines.*

**Proof** If some symp is non-thick, then they are all non-thick by the previous lemma. So  $\Delta$  is also a non-thick generalized quadrangle, hence a grid. Since the blocks endowed with the lines contained in it are linear spaces, the first assertion follows. The second assertion is easy.  $\square$

A natural question is whether a block of  $\Delta$  coincides with a line of  $\Gamma$ . Lemma 3.1 implies that, whenever for some line  $L$  of  $\Gamma$ , there is a point of  $\Gamma$  not collinear with any point on  $L$ , then  $L$  is properly contained in a block of  $\Delta$ . Hence, if for every line  $L$  of  $\Gamma$ , there is a point of  $\Gamma$  not collinear with any point of  $L$ , then every line is properly contained in a block. Now, the converse is also true, and we have the following result.

**Proposition 3.7** *The following are equivalent.*

- (1) *Some line of  $\Gamma$  coincides with some block of  $\Delta$ .*
- (2) *For some line  $L$  of  $\Gamma$  every point of  $\Delta$  is collinear to some point of  $L$ .*
- (3) *The quadrangle  $\Delta$  is isomorphic to the product geometry of a line  $L$  of  $\Gamma$  with some block of  $\Delta$  properly containing lines of  $\Gamma$ .*

**Proof** By the above considerations we only need to show that (1) and (3) follow from (2). So let  $L$  be a line of  $\Gamma$  such that every point of  $\Gamma$  is collinear with a point of  $L$ . We first claim that every symp  $H$  containing any point  $x$  of  $L$  contains  $L$ . Indeed, if  $y \in H$  is a point not collinear with  $x$ , then by assumption  $\xi(x, y)$  must contain a second point of  $L$  (namely, the one collinear with  $y$ ).

Now let  $H$  be a symp with  $L \subseteq H$ . Since  $\mathcal{P} \neq H$ , there is some point  $z$  outside  $H$ , and hence some symp  $H'$  containing  $z$  and a point of  $L$ . By the previous claim,  $H'$  contains  $L$ . Let  $x, y \in L$ ,  $x \neq y$ . We show that  $H'$  is not thick. Suppose for a contradiction that  $H'$  is thick. Select a point  $p \in H' \setminus H$  collinear to  $x$  but not collinear to  $y$ . Let  $U$  be the set of points of  $H'$  collinear to  $p$ . By Lemma 2.2,  $U$  is a subspace containing  $x$ , but not  $y$ . Since  $H'$  is not a grid, there is a point  $p' \in H' \setminus H$  collinear to  $x$  and not lying in  $U$ . Hence  $p'$  and  $p$  are not collinear and, by convexity,  $H'' := \xi(p, p')$  contains  $x$ . But it does not contain  $y$  because the intersection  $H \cap H''$  would otherwise not be a singular subspace, as it would contain the non-collinear points  $p$  and  $y$ . This contradicts our first claim. Hence Lemma 3.5(ii) yields that all symps are grids.

Proposition 3.6 now implies that  $\Delta$  is isomorphic to the product of two blocks, say  $B_1$  and  $B_2$ , and we can assume that  $L \subseteq B_1$ . If  $L$  is properly contained in  $B_1$ , say  $x \in B_1 \setminus L$ , then every point  $z$  not in  $B_1$  but collinear with  $x$  has the property that it is collinear with no point of  $L$ , a contradiction. This shows (1) and (3).  $\square$

In the rest of this section, we may assume that the symps are thick.

Let  $B$  be a block of  $\Delta$  and let  $H$  be a symp of  $\Gamma$  with  $B \cap H$  not a line of  $H$ . Since  $H$  is ideal,  $B$  does not contain any point of  $H$ . Let  $u \in H$  be arbitrary. Then there is a unique block  $B_u$  through  $u$  intersecting  $B$  in a point. The block  $B_u$  intersects  $H$  in a line  $L_u$  and we have a mapping  $u \mapsto L_u$  from the point set of  $H$  to the line set of  $H$  (but this mapping depends on  $B$ ). For  $u, v \in H$ , we easily have  $L_u \cap L_v = \emptyset$  or  $L_u = L_v$ . Hence we obtain a partition of  $H$  into lines, which is usually called a *spread* of  $H$ . Here, the spread is said to be *induced (by  $B$ )*. These induced spreads have an interesting property, which is trivial for non-thick symps, hence we may state it nevertheless in general.

**Lemma 3.8** (i) *Every pair of non-concurrent lines of any symp of  $\Gamma$  is regular.*

(ii) *If  $L, M$  are two distinct lines of some induced spread  $\mathcal{S}$  in some symp  $H$ , then every member of  $(\{L, M\}^\perp)^\perp$  belongs to the spread. Moreover, if the spread is induced by the block  $B$ , then the point-line geometry  $\beta$  induced on  $B$  by the lines of  $\Gamma$  contained in  $B$  contains a subspace which is isomorphic to the point-line geometry  $\sigma$  induced on  $\mathcal{S}$  by the “double perps”, i.e., the point-line geometry with point set  $\mathcal{S}$  and lines the sets  $(\{L, M\}^\perp)^\perp$ , with  $L, M \in \mathcal{S}$ .*

**Proof** Consider a symp  $H$  and let  $L_1, L_2$  be two non-concurrent lines of  $H$ . Let  $B_1, B_2$  be the blocks containing  $L_1, L_2$ , respectively. Let  $x_1 \in B_1 \setminus L_1$ . Then there is a unique point  $x_2 \in B_2$  collinear with  $x_1$ . Since  $H$  is a subquadrangle of  $\Delta$ , the point  $x_2$  does not belong to  $L_2$ . We note that the block  $B$  containing  $x_1$  and  $x_2$  does not contain a point of  $H$  because if  $u \in B \cap H$ , then the unique point on  $B_1$  collinear with  $u$  would be contained in  $L_1$  since  $H$  is a subquadrangle. But that point is  $x_1$ , a contradiction.

Let  $M_1, M_2 \in \{L_1, L_2\}^\perp$  be distinct. Put  $y_{i,j} = L_i \cap M_j$ . The symp  $\xi(x_1, y_{2,j})$ ,  $j = 1, 2$ , contains the lines  $x_1x_2$  and  $M_j$ . Hence, for any point  $x \in x_1x_2$ , there is a unique point  $y_j \in M_j$  collinear to  $x$ . If  $y_1$  were not collinear to  $y_2$ , then  $H = \xi(y_1, y_2)$  would contain  $x$ , a contradiction. Hence  $y_1$  and  $y_2$  are collinear. Varying  $M_2$  over  $\{L_1, L_2\}^\perp \setminus \{M_1\}$ , we see that the line  $y_1y_2$  belongs to  $(\{L_1, L_2\}^\perp)^\perp$ . This shows  $(\{L_1, L_2\}^\perp)^\perp = \{M_1, M_2\}^\perp$ , and hence the pair  $\{L_1, L_2\}$  is regular.

But, as is obvious from the previous paragraph, all elements of  $\{M_1, M_2\}^\perp$  belong to the spread induced by  $B$ , and the corresponding blocks intersect  $B$  precisely in (all) points of the line  $x_1x_2$ . Hence the mapping  $y_1y_2 \mapsto x$  defines an injective morphism from  $\sigma$  to  $\beta$ . This completes the proof of the lemma.  $\square$

We now arrive at the crux of this section.

**Theorem 3.9** *If  $\Gamma$  has thick symps, then every maximal singular subspace of  $\Gamma$  contains a non-closed O’Nan configuration. In particular, no maximal singular subspace is a projective space.*




Figure 1: Proof of Lemma 3.9

**Proof** Let  $H$  be any symp and let  $\mathcal{S}$  be a spread induced by some block of  $\Delta$  outside  $H$ . By the previous lemma, it suffices to show that the geometry  $\sigma$  with point set  $\mathcal{S}$  and lines the double perps contains non-closed O’Nan configurations.

Consider lines  $X, L_1, L_2, M_1 \in \mathcal{S}$  such that  $X \in (\{L_1, L_2\}^\perp)^\perp =: R$ ,  $M_1 \notin R$ , see Figure 1. The fact that we can choose  $M_1 \notin R$  follows from the fact that  $H$  is thick and hence there is at least one point  $x_1$  of  $H$  not on any line of  $R$ ; the spread line through  $x_1$  is a good choice for  $M_1$ . Note that  $M_1$  does not contain any point of any line of  $R$  since all members of  $R$  belong to  $\mathcal{S}$ .

Consider a line  $N \in R^\perp$  (and note  $N \cap M_1 = \emptyset$  since all points of  $N$  belong to members of  $R$ ), and the points  $p_1 = N \cap L_1$  and  $p_2 = N \cap L_2$ . Let  $q_1$  be the unique point of  $M_1$  collinear to  $p_1$ . Consider the unique line  $K$  incident with  $q_1$  and concurrent with  $X$ . Let  $q_2$  be the unique point of  $K$  collinear to  $p_2$ . Let  $M_2 \in (\{X, M_1\}^\perp)^\perp$  be such that  $q_2 \in M_2$ . Suppose first there exists  $Y \in (\{L_1, M_1\}^\perp)^\perp \cap (\{L_2, M_2\}^\perp)^\perp \cap \mathcal{L}$ . Then  $Y \in \mathcal{S}$  by Lemma 3.8(ii).

As  $p_i q_i \in \{L_i, M_i\}^\perp$ ,  $i = 1, 2$ , the line  $Y$  meets both  $p_1 q_1$  and  $p_2 q_2$ . Hence  $Y, N, K \in \{p_1 q_1, p_2 q_2\}^\perp$ . Since  $X$  intersects both  $N$  and  $K$ , we have  $X \in (\{p_1 q_1, p_2 q_2\}^\perp)^\perp$ . By regularity  $Y$  is concurrent with  $X$ , a contradiction to  $X, Y \in \mathcal{S}$ . Hence  $(\{L_1, M_1\}^\perp)^\perp \cap (\{L_2, M_2\}^\perp)^\perp = \emptyset$ , and the four double perps  $(\{L_1, L_2\}^\perp)^\perp$ ,  $(\{L_1, M_1\}^\perp)^\perp$ ,  $(\{M_1, M_2\}^\perp)^\perp$  and  $(\{L_2, M_2\}^\perp)^\perp$  form a non-closed O’Nan configuration.  $\square$

Theorem 3.9 has been shown by Cohen in [3], Proposition 4.2, in case the symplecta are *classical* generalized quadrangles, i.e., for generalized quadrangles occurring as point-residues in polar spaces of rank 3, with an extensive and explicit calculation. In fact, Proposition 4.2 of [3] does not refer to the imbrex geometry and treats the following more general Problem (\*): *does a spread  $\mathcal{S}$  in thick generalized quadrangle  $\Gamma$  exist with the property that every pair of lines in  $\mathcal{S}$  is regular, and the geometry defined by  $\mathcal{S}$  and the double perps  $(\{L, M\}^\perp)^\perp$ ,  $L, M \in \mathcal{S}$ , is a projective space.* Then Cohen proves that, if  $\Gamma$  is a classical generalized quadrangle, the answer to (\*) is negative. More recently, Shult & K. Thas [11] have found a more elegant proof only using

the *Moufang property* of such quadrangles (and in fact, they only use the *strong transitivity*, i.e., the transitivity of the automorphism group on the set of pairs of incident point-line pairs  $(p, L), (q, M)$  such that  $p, q$  are not collinear and  $L, M$  not concurrent). Hence for strongly transitive quadrangles, the answer to (\*) is also negative. Finally, our proof above only uses the fact that for some line  $L$  of the spread, there exists at least one line  $M$  not in the spread such that  $\{L, M\}$  is a regular pair (in our proof,  $L$  is the line  $X$  and  $M$  is the line  $p_2q_2$ ).

Note that in the finite case the answer to (\*) is always negative our assumptions coincide with those of Cohen and Shult & K. Thas in the above references.

**Examples.** We now present a class of examples of imbrex geometries of symplectic rank 2 with thick symps. In this case, the maximal singular subspaces can be seen as higher dimensional unitals.

Let  $\Delta$  be any generalized quadrangle whose dual  $\Omega$  is thick and embedded in some (possibly infinite-dimensional) projective space  $\mathbb{P}(\mathbb{L})$  over the skew field  $\mathbb{L}$ , i.e., the blocks of  $\Delta$  are points of  $\mathbb{P}(\mathbb{L})$  and the points of  $\Delta$  are lines of  $\mathbb{P}(\mathbb{L})$  (with natural incidence). All such  $\Omega$  are classified, see [13, 14] (every classical generalized quadrangle in the above sense qualifies). Now we require that every plane of  $\mathbb{P}(\mathbb{L})$  containing at least two lines of  $\Omega$  contains at least three lines of  $\Omega$ . Let  $\Gamma = (\mathcal{P}, \mathcal{L}, *)$  be the point-line geometry with same point set as  $\Delta$ , and where the lines correspond to planes of  $\mathbb{P}(\mathbb{L})$  containing at least two lines of  $\Omega$ , with natural incidence. Since such planes contain at least three lines of  $\Omega$ , the geometry  $\Gamma$  has thick lines (i.e., each line has at least three points). Consider Axiom (PPS2) and let  $L, M$  be the lines of  $\mathbb{P}(\mathbb{L})$  corresponding to two non-collinear points of  $\Gamma$ . Then  $\langle L, M \rangle$  is 3-dimensional. Now  $\langle L, M \rangle$  induces a subquadrangle  $H$  of  $\Omega$ , and  $H$  is generated by  $L, M$ . Hence (PPS2) holds with the symplecta being the subquadrangles of  $\Delta$  corresponding to 3-spaces of  $\mathbb{P}(\mathbb{L})$  containing two non-intersecting lines of  $\Omega$  (and these quadrangles are automatically thick). Now (PS1) is easy to check if we assume  $n \geq 4$  (otherwise there is only one symp).

We now check the Axiom (Imb). Translated to  $\Omega$ , we have to show that for given lines  $L, M_1, M_2$  of  $\Omega$  such that  $M_1$  and  $M_2$  intersect and  $L$  does not intersect the plane  $\langle M_1, M_2 \rangle$ , the 3-spaces  $\langle L, M_1 \rangle$  and  $\langle L, M_2 \rangle$  intersect in a plane  $\pi$  containing at least two lines of  $\Omega$ . Clearly  $\pi = \langle L, M_1 \cap M_2 \rangle$  and this contains, besides  $L$ , also the line through  $M_1 \cap M_2$  intersecting  $L$ .

So, in conclusion, every embedded generalized quadrangle spanning a projective space of dimension at least 4 and such that the subquadrangles induced by 3-spaces are thick, gives rise to an example of an imbrex geometry of symplectic rank 2.

In the finite case, there is just one class of such examples, namely where  $\Omega$  is the Hermitian quadrangle denoted by  $Q(4, q^2)$ . The maximal singular subspaces of  $\Gamma$  are in this case classical unitals. In the infinite case, all quadrangles arising from a  $\sigma$ -quadratic form, with  $\sigma$  nontrivial and with nontrivial isotropic kernel, qualify.

Similarly, other classes of examples are the exceptional Moufang quadrangles of types  $F_4, E_6, E_7$  and  $E_8$ . Using the terminology of Appendix C of [17], if we let points correspond to the isotropic orbit in the Tits diagram which corresponds to the multiple root in the root system of type  $BC_2$ , then the symplecta are the ideal subquadrangles belonging to a root system of type  $C_2$ . Again, all axioms of an imbrex geometry hold.

## 4 Imbrex geometries of symplectic rank at least 3

In this section,  $\Gamma = (\mathcal{P}, \mathcal{L}, *)$  is an imbrex geometry of symplectic rank  $r$  at least 3. Note that by (PPS3),  $r$  is finite.

**Lemma 4.1** *Let  $(x, H)$  be a non-incident point-symplecton pair in  $\Gamma$ . If  $x$  is collinear with all points of a line  $L \subseteq H$ , then  $x$  is collinear with all points of a maximal singular subspace of  $H$ .*

**Proof** Let  $p \in H$  be a point which is not collinear to all points of  $L$ , and let  $H' = \xi(p, x)$ . Then by convexity  $H'$  contains the unique point  $p'$  on  $L$  collinear to  $p$  in  $H$ . Select  $x' \in L$  different from  $p'$ . Then we claim that no point on the line  $xx'$  is collinear to  $p$ . Indeed, suppose some point  $y \in xx'$  is collinear to  $p$ , then  $y \in H'$ , and hence also  $x' \in H'$ , contradicting  $H = \xi(p, x')$ . Our claim is proved.

Now Axiom (Imb) implies that  $H$  and  $H'$  intersect in a maximal singular subspace  $U$ . In  $H'$ , the point  $x$  is collinear with an  $(r-1)$ -subspace  $U'$  of  $U$ , which does not contain  $x'$  as  $x' \notin H'$ . Hence  $x$  is collinear with all points of  $U'$  and with  $x'$ , and these must then generate a maximal singular subspace since  $r$  is finite.  $\square$

Now we can apply Theorem 15.4.5 of [10]. The latter theorem is an updated summary of the work done by Cohen [3] and Cohen & Cooperstein [4]. Since we assume diameter 2, our conclusion will only contain a restricted list of geometries. Without going into too much detail, we define the *Lie incidence geometry*  $X_{n,i}(\mathbb{K})$ , where  $X \in \{A, D, E\}$ , as the  $i$ -Grassmannian point-line geometry related to the building of type  $X_n$  over the skew field  $\mathbb{K}$  (which is automatically a field for the cases D and E), and we use Bourbaki labeling [1] for the nodes of the corresponding diagram, and hence for the subscripts. For instance, the Lie incidence geometry  $A_{n,i}$  is the ordinary Grassmannian geometry of all  $i$ -spaces of the  $(n+1)$ -dimensional vector space over  $\mathbb{K}$ .

**Corollary 4.2** *An imbrex geometry  $\Gamma$  of symplectic rank  $r \geq 3$  is exactly one of the following.*

- ( $r = 3$ ) *The Lie incidence geometry  $A_{n,2}(\mathbb{L})$ , for  $\mathbb{L}$  any skew field,  $n \geq 4$ .*
- ( $r = 4$ ) *The Lie incidence geometry  $D_{5,5}(\mathbb{K})$ , for any field  $\mathbb{K}$ .*
- ( $r = 5$ ) *The Lie incidence geometry  $E_{6,1}(\mathbb{K})$ , for any field  $\mathbb{K}$ .*

**Proof** If  $\Gamma$  is an imbrex geometry of symplectic rank at least 3, then the assertion follows from Theorem 15.4.5 of [10]. Conversely, if  $\Gamma$  is one of the mentioned geometries, then we will prove in the next section that their universal embedding is a local Mazzocca-Melone set (for definitions, see the next section), and this will immediately imply that they satisfy (Imb). Hence we postpone the details to the next section.  $\square$

## 5 Local Mazzocca-Melone sets

We now apply the classification of the previous section to the theory of Mazzocca-Melone sets, initiated in [9]. We introduce pre-Mazzocca-Melone sets and local Mazzocca-Melone sets in the next subsection.

### 5.1 Definition of local Mazzocca-Melone sets

Let  $N, d, r$  be natural numbers. Let  $X$  be a spanning point set of  $\mathbb{P}^n(\mathbb{K})$ , with  $\mathbb{K}$  any skew field, and let  $\Xi$  be a collection of  $(d+1)$ -spaces of  $\mathbb{P}^n(\mathbb{K})$ ,  $d \geq 2$ , such that, for any  $\xi \in \Xi$ , the intersection  $\xi \cap X =: X(\xi)$  is a polar space of rank  $r$ ,  $r \geq 2$ , in  $\xi$  (and then, for  $x \in X(\xi)$ , we denote the tangent space at  $x$  to  $X(\xi)$  by  $T_x(X(\xi))$  or sometimes simply by  $T_x(\xi)$ ). Also, we say that two points of  $X$  are *X-collinear* if all points of the line they span in  $\mathbb{P}^n(\mathbb{K})$  are contained in  $X$ . We call  $(X, \Xi)$  a *pre-Mazzocca-Melone set (of type  $(d, r)$ )* if (MM1) and (MM2) are satisfied, and a *local Mazzocca-Melone set* if on top (LMM3) holds.

- (MM1) Any pair of points  $x$  and  $y$  of  $X$  which are not  $X$ -collinear lies in at least one element of  $\Xi$ .
- (MM2) If  $\xi_1, \xi_2 \in \Xi$ , with  $\xi_1 \neq \xi_2$ , then  $\xi_1 \cap \xi_2 \subset X$ .

It follows from (MM1) and (MM2) that any pair of points  $x$  and  $y$  of  $X$  which are not  $X$ -collinear lies in exactly one member of  $\Xi$ , and we denote that member by  $[x, y]$ . The pre-Mazzocca-Melone set  $(X, \Xi)$  is called *proper* if  $|\Xi| \geq 2$ . It follows from these axioms that proper pre-Mazzocca-Melone sets define parapolar spaces where the symps are the polar spaces  $X(\xi)$ ,  $\xi \in \Xi$ , see [9].

- (LMM3) If  $x \in X$  and  $L \subseteq X$  is a line of  $\mathbb{P}^n(\mathbb{K})$  such that no point of  $L$  is  $X$ -collinear with  $x$ , then all  $d$ -spaces  $T_x([x, y])$ ,  $y \in L$ , generate a subspace  $T_{x,L}$  of  $\mathbb{P}^n(\mathbb{K})$  of dimension at most  $2d - r + 1$ .

A local Mazzocca-Melone set is *proper* if it is a proper pre-Mazzocca-Melone set. Non-proper pre-Mazzocca-Melone sets are just embedded polar spaces. In this section we shall classify all proper local Mazzocca-Melone sets of type  $(d, r)$ , for all  $d, r \geq 2$ .

## 5.2 Examples of proper local Mazzocca-Melone sets

In [15], it is proved that the *Segre variety*  $\mathcal{S}_{p,q}(\mathbb{K})$ ,  $p \geq 1$  and  $q \geq 1$ , satisfies the axioms of a local Mazzocca-Melone set (although it was not called as such in [15]).

Now, the Lie incidence geometries  $A_{n,2}(\mathbb{K})$ , for any field  $\mathbb{K}$ ,  $n \geq 4$ ,  $D_{5,5}(\mathbb{K})$ , for any field  $\mathbb{K}$ , and  $E_{6,1}(\mathbb{K})$ , for any field  $\mathbb{K}$ , all admit the so-called *universal embedding*, which is a pre-Mazzocca-Melone set, as proved in [9]. We denote these pre-Mazzocca-Melone sets by  $\mathcal{A}_{p,2}(\mathbb{L})$ ,  $p \geq 4$ ,  $\mathcal{D}_{5,5}(\mathbb{K})$  and  $\mathcal{E}_{6,1}(\mathbb{K})$ , respectively satisfy the above axioms. We use slightly different notation here since we view them as varieties.

We now show that these are local Mazzocca-Melone sets. By inclusion of the appropriate sets, it suffices to show (LMM3) for the minimal cases, namely,  $\mathcal{A}_{4,2}(\mathbb{K})$ ,  $\mathcal{D}_{5,5}(\mathbb{K})$  and  $\mathcal{E}_{6,1}(\mathbb{K})$ . Note that in this list, a preceding one is the *residue* of the next one, i.e., if  $x$  is a point in one of these sets  $X$ , then the lines of  $X$  through  $x$  form, together with the tangent spaces to the symps at  $x$ , a pre-Mazzocca-Melone set in a subspace of the quotient projective space with respect to  $x$  isomorphic to the previous Mazzocca-Melone set in the list. Hence the assertion will be proved if we show that the validity of (LMM3) for a given set follows from the validity of that axiom in any residue, adding  $\mathcal{S}_{1,2}(\mathbb{K})$  in front of that list. The latter indeed satisfies (LMM3) by [15], see the first sentence of this section.

So suppose (LMM3) is valid in the residue  $(X_p, \Xi_p)$  of  $(X, \Xi)$  at  $p \in X$ , for all  $p \in X$ , with  $(X, \Xi)$  one of the pre-Mazzocca-Melone sets  $\mathcal{A}_{4,2}(\mathbb{L})$ ,  $\mathcal{D}_{5,5}(\mathbb{K})$  or  $\mathcal{E}_{6,1}(\mathbb{K})$ . Let  $r - 1$  be the dimension of the maximal singular subspaces of the symps (having rank  $r$ ) of  $(X, \Xi)$  (which are then hyperbolic quadrics in  $(2r - 1)$ -dimensional spaces, and  $d = 2r - 2$ ). Let  $x \in X$  and let  $L$  be a line contained in  $X$  no point of which is  $X$ -collinear with  $x$ . Let  $H$  and  $H'$  be two distinct symps containing  $x$  and a (different) point of  $L$ , say  $y, y'$ , respectively. We first claim that  $H \cap H'$  has dimension  $r - 1$  (hence that (Imb) holds). Since  $y'$  is collinear with the point  $y$  of  $H$ , it is collinear with an  $(r - 1)$ -space  $U$  of  $H$  (this can be checked in all instances directly from the definitions of the corresponding Lie incidence geometries). There is a unique  $(r - 2)$ -space  $W \subseteq U$  all of whose points are collinear with  $x$ . By the Quadrangle Lemma, the singular  $(r - 1)$ -space  $\langle x, W \rangle$  is contained in  $H'$ , showing our claim.

Moreover, since both  $y$  and  $y'$  are collinear with all points of  $W$ , all points of  $L$  are collinear with all points of  $W$  and  $\langle L, W \rangle$  is a singular  $r$ -space. The Quadrangle Lemma implies that any symp  $H^*$  through  $x$  and a point of  $L$  contains  $W$ . Since  $r \geq 3$ , we can select two distinct points  $q, q'$  of  $W$ . Clearly,  $\langle H^* \rangle$  is generated by  $T_q(H^*)$  and  $T_{q'}(H^*)$ . Hence  $\langle T_{x,L}, L \rangle$  is generated by all  $T_q(H^*)$  and  $T_{q'}(H^*)$ , for  $H^*$  running through all symps through  $x$  and a point of  $L$ . By induction, and since in the case  $d = 2r - 2$  equality is reached in (LMM3), the dimension of the span of all  $T_q(H^*)$  for  $H^*$  as above, is equal to  $2(2r - 4) - (r - 1) + 1 + (2 + 1) = 3r - 3$  (in the residue at  $q$ , the point  $\langle q, x \rangle$  is not  $X_q$ -collinear to any point of  $\langle q, L \rangle$ ). Similarly for the span of all  $T_{q'}(H^*)$ . The intersection of those two spaces is, by (double) induction  $2(2r - 6) - (r - 2) + 1 + (2 + 2) = 3r - 5$

(look in the residue of the line  $\langle q, q' \rangle$ ; for  $r = 3$ , there is no “double” induction, but then this can be seen directly in  $\mathcal{S}_{1,2}(\mathbb{K})$ ). Hence the dimension of  $\langle T_{x,L}, L \rangle$  is equal to  $3r - 1$ . Hence, since  $L$  does not meet  $T_x(H)$ , we see that  $\dim(T_{x,L}) = 3r - 3 = 2(2r - 2) - r + 1$ , exactly what we had to prove.

### 5.3 Classification of Local Mazzocca-Melone sets

Using the results on imbrex geometries we now classify all local Mazzocca-Melone sets, up to projection from a suitable subspace (i.e., a subspace containing no point in the span of two symplecta).

**Theorem 5.1** *A proper local Mazzocca-Melone set of type  $(d, r)$ ,  $d, r \geq 2$  is projectively equivalent to a projection from a suitable subspace (in the above sense) of one of the following pre-Mazzocca-Melone sets.*

$d = 2$ : the Segre variety  $\mathcal{S}_{m,n}(\mathbb{K})$ ,  $m \geq 1$  and  $n \geq 1$ , but  $nm > 1$ ,

$d = 4$ :  $\mathcal{A}_{n,2}(\mathbb{K})$ ,  $n \geq 4$ ,

$d = 6$ :  $\mathcal{D}_{5,5}(\mathbb{K})$ ,

$d = 8$ :  $\mathcal{E}_{6,1}(\mathbb{K})$ .

**Proof** Clearly, a local Mazzocca-Melone set is an imbrex geometry. So we can apply the results of Section 3. Suppose first that  $r = 2$ . Then the symplectic rank as a parapolar space is 2, and as the geometry is embedded, all singular subspaces are projective spaces and hence do not contain non-closed O’Nan configurations. By Lemma 3.5(ii), Theorem 3.9 and [15], we are done. If  $r \geq 3$ , we use Corollary 4.2, and the fact that all embeddings arise from the absolute universal embedding by suitable projection, see [6].  $\square$

### 5.4 Weakening the (Imb) axiom

If we would not care about the residual property of (Imb), then an interesting option would be to weaken it to the following condition.

(Imb\*) *Let  $x$  be a point not collinear with any point of the line  $L$ . Let  $y_1, y_2$  be distinct points on  $L$ . Then the symplecta  $\xi(x, y_1)$  and  $\xi(x, y_2)$  intersect in a singular subspace of dimension at least 1.*



This does not guarantee constant symplectic rank from the beginning, and we were not able to classify strong parapolar spaces of diameter 2 and symplectic rank at least 3 under Condition (Imb<sup>\*</sup>). However, it would be interesting to do so, since, from the point of view of Hjelmslev, this is the weakest condition one can ask. We pose it here as an open problem.

**Acknowledgement** The authors would like to thank the referees for their very detailed reports which improved this paper, and the Mathematisches Forschungsinstitut Oberwolfach for providing them with the best possible research environment imaginable during a stay as Oberwolfach Leibniz Fellow (JS) and Visiting Senior Researcher (HVM) in June 2013.

## References

- [1] N. Bourbaki, *Groupes et Algèbres de Lie*, Chapters 4,5 and 6. *Actu. Sci. Ind.* **1337**, Hermann, Paris, 1968.
- [2] F. Buekenhout & E.E. Shult, On the foundations of polar geometry. *Geom. Dedicata* **3** (1974), 155–170.
- [3] A. Cohen, On a theorem of Cooperstein. *European J. Combin.* **4** (1983), 107–126.
- [4] A. Cohen & B. Cooperstein, A characterization of some geometries of Lie type. *Geom. Dedicata* **15** (1983), 73–105.
- [5] J. Hjelmslev, Die natürliche Geometrie. *Abh. Mathem. Sem. Univ. Hamburg (2)* (1923), 1–36.
- [6] A. Kasikova & E.E. Shult, Absolute embeddings of point-line geometries. *J. Algebra*, **238** (2001), 265–291.
- [7] W. Klingenberg, Projektive und affine Ebenen mit Nachbarelementen. *Math. Z.* **60** (1954), 384–406.
- [8] F. Mazzocca & N. Melone, Caps and Veronese varieties in projective Galois spaces. *Discrete Math.* **48** (1984), 243–252.
- [9] J. Schillewaert & H. Van Maldeghem, Severi varieties over arbitrary fields. *Submitted*.
- [10] E. E. Shult, *Points and Lines, Characterizing the Classical Geometries*. Universitext, Springer-Verlag, Berlin Heidelberg, 2011.
- [11] E. E. Shult & K. Thas, A theorem of Cohen on parapolar spaces. *Combinatorica* **30** (2010), 435–444.
- [12] T. A. Springer & F. D. Veldkamp, On Hjelmslev-Moufang planes. *Math. Z.* **107** (1968), 249–263.
- [13] A. Steinbach & H. Van Maldeghem, Generalized quadrangles weakly embedded of degree  $> 2$  in projective space. *Forum Math.* **11** (1999), 139–176.

- [14] A. Steinbach & H. Van Maldeghem, Generalized quadrangles weakly embedded of degree 2 in projective space. *Pacific J. Math.* **193** (2000), 227–248.
- [15] J. A. Thas & H. Van Maldeghem, Characterizations of Segre varieties, *J. Combin. Theory Ser. (A)* **120** (2013), 795–802.
- [16] J. Tits, *Buildings of Spherical Type and Finite BN-Pairs*. Lecture Notes in Mathematics **386**. Springer-Verlag, Berlin-New York, 1974.
- [17] H. Van Maldeghem, *Generalized Polygons*, Monographs in Mathematics **93**, Birkhäuser Verlag, Basel, 1998.
- [18] F.D. Veldkamp, Polar geometry. I, II, III, IV, V. *Nederl. Akad. Wetensch. Proc. Ser. A* **62**; **63**, *Indag. Math.* **21** (1959), 512–551 **22** (1959) 207–212.