

The smallest sets of points meeting all generators of $H(2n, q^2)$, $n \geq 2$

Jan De Beule

Ghent University, Department of Pure Mathematics and Computer Algebra,
Krijgslaan 281, 9000 Gent, Belgium.

Joint work with: Klaus Metsch

Universität Gießen, Mathematisches Institut, Arndtsstraße 2, D-35392 Gießen,
Germany

It is known that the Hermitian varieties $H(2n, q^2)$, $n \geq 2$, have no ovoids. The question arises how the smallest sets of points meeting every generator look like.

We start with the case $H(4, q^2)$. Supposing that \mathcal{K} is a minimal blocking set of size $q^5 + \delta$, $\delta \leq q^2$, we obtain a contradiction in several steps if $\delta < q^2$. The main step is to consider intersections with hyperplanes. The considered $H(3, q^2)$ also contain lines as generators and hence the subsets of \mathcal{K} in the corresponding hyperplanes must block these generators and contain a minimal number of points. This leads to $\delta = q^2$ and hence $|\mathcal{K}| \geq q^5 + q^2$. If we have equality, a short extra argument shows that \mathcal{K} is the set of points of a cone with base $H(2, q^2)$ and vertex a point of $H(4, q^2)$, minus the point itself. This is quite classical and analogue results are known for other polar spaces.

Having the result for $H(4, q^2)$, it is now possible to characterise the smallest minimal blocking sets of $H(2n, q^2)$, $n > 2$. Looking in quotient geometries and using the result for $H(4, q^2)$, together with standard combinatorial arguments, proves that \mathcal{K} is a *truncated cone* with base a Hermitian curve $H(2, q^2)$ and vertex an $(n - 2)$ -dimensional subspace contained in $H(2n, q^2)$

References

- [1] J. De Beule and L. Storme. The smallest minimal blocking sets of $Q(6, q)$, q even. *J. Combin. Des.*, 11(4):290–303, 2003.
- [2] J. De Beule and L. Storme. The smallest minimal blocking sets of $Q(2n, q)$, for small odd q . *Discrete Math.*, submitted.
- [3] K. Metsch. Small point sets that meet all generators of $W(2n + 1, q)$. *Des., Codes Cryptogr.*, to appear
- [4] K. Metsch. The sets closest to ovoids in $Q^-(2n+1, q)$. *Bull. Belg. Math. Soc. Simon Stevin*, 5(2-3):389–392, 1998. Finite geometry and combinatorics (Deinze, 1997).
- [5] K. Metsch. On blocking sets of quadrics. *J. Geom.*, 67(1-2):188–207, 2000. Second Pythagorean Conference (Pythagoreion, 1999).