

On Cameron-Liebler line classes with large parameter

J. De Beule

(joint work with Jeroen Demeyer, Klaus Metsch and Morgan Rodgers)

Department of Mathematics
Ghent University

February 13, 2012
Giornate di Geometria 2012, Vicenza

definitions

Definition

A set $\mathcal{L}$ of lines of $\text{PG}(3, q)$ is a *Cameron-Liebler* line class with parameter x if and only if $|\mathcal{L} \cap \mathcal{S}| = x$ for every spread $\mathcal{S}$ of $\text{PG}(3, q)$.

- Introduced in an attempt to classify collineation groups of $\text{PG}(3, q)$ that have equally many point orbits and line orbits.
- different equivalent definitions.

Definition

A set $\mathcal{L}$ of lines of $\text{PG}(3, q)$ is a *Cameron-Liebler* line class with parameter x if and only if for every line l

$$|\{m \in \mathcal{L} \setminus \{l\} \mid m \cap l \neq \emptyset\}| = (q+1)x + (q^2-1)\chi_{\mathcal{L}}(l)$$

definitions

Definition

A set $\mathcal{L}$ of lines of $\text{PG}(3, q)$ is a *Cameron-Liebler* line class with parameter x if and only if $|\mathcal{L} \cap \mathcal{S}| = x$ for every spread $\mathcal{S}$ of $\text{PG}(3, q)$.

- Introduced in an attempt to classify collineation groups of $\text{PG}(3, q)$ that have equally many point orbits and line orbits.
- different equivalent definitions.

Definition

A set $\mathcal{L}$ of lines of $\text{PG}(3, q)$ is a *Cameron-Liebler* line class with parameter x if and only if for every line l

$$|\{m \in \mathcal{L} \setminus \{l\} \mid m \cap l \neq \emptyset\}| = (q+1)x + (q^2-1)\chi_{\mathcal{L}}(l)$$

definitions

- $\theta_r(q) := \frac{q^{r+1}-1}{q-1}$

Definition

An x -tight set $\mathcal{L}$ of a finite classical polar space $\mathcal{P}$ of rank $r \geq 2$, is a set of $x\theta_{r-1}(q)$ points, such that

$$|P^\perp \cap \mathcal{L}| = \begin{cases} x\theta_{r-2}(q) + q^{r-1} & \text{if } P \in \mathcal{L} \\ x\theta_{r-2}(q) & \text{if } P \notin \mathcal{L}. \end{cases}$$

A Cameron-Liebler line class of $\text{PG}(3, q)$ with parameter x , is equivalent to an x -tight set of $\text{Q}^+(5, q)$.

definitions

- $\theta_r(q) := \frac{q^{r+1}-1}{q-1}$

Definition

An x -tight set $\mathcal{L}$ of a finite classical polar space $\mathcal{P}$ of rank $r \geq 2$, is a set of $x\theta_{r-1}(q)$ points, such that

$$|P^\perp \cap \mathcal{L}| = \begin{cases} x\theta_{r-2}(q) + q^{r-1} & \text{if } P \in \mathcal{L} \\ x\theta_{r-2}(q) & \text{if } P \notin \mathcal{L}. \end{cases}$$

A Cameron-Liebler line class of $\text{PG}(3, q)$ with parameter x , is equivalent to an x -tight set of $\text{Q}^+(5, q)$.

non-existence

Theorem (K. Metsch)

A Cameron-Liebler line class in $\text{PG}(3, q)$ with parameter x does not exist for $2 < x \leq q$.

Constructions

Constructions of Cameron-Liebler line classes:

- Bruen, Drudge: q odd, $x = \frac{q^2+1}{2}$
- Govaerts, Penttila: $q = 4$, $x \in \{4, 5\}$.

The quest for new examples

- $q = 2 \pmod 3, x = \frac{(q+1)^2}{3}$
- $q = 3^h, x = \frac{(q^2-1)}{2}$
- For all examples, the group $C_3 : C_{q^2+q+1}$ is a subgroup of the automorphism group

Recent examples for $q \not\equiv 1 \pmod 3$

Morgan Rodgers found examples for $q \leq 200$:

- $q \equiv 1 \pmod 4$: $x = \frac{q^2-1}{2}$
- $q \equiv 2 \pmod 4$: $x = \frac{(q+1)^2}{3}$

Algebraic combinatorics

Theorem (Bamberg, Kelly, Law, Penttila)

Let A be the collinearity matrix of $Q^+(5, q)$, and let $\mathcal{L}$ be an x -tight set with characteristic vector χ . Then

$$\chi - \frac{x}{q^2 - 1} \mathbf{j}$$

is an eigenvector of A with eigenvalue $q^2 - 1$, $\mathbf{j}$ the all one vector.

Using the group

- $G = C_{q^2+q+1}$
- orbits on points of $\text{PG}(3, q)$: $\pi_\infty, \{(1, 0, 0, 0)\}$, $q - 1$ orbits of length $q^2 + q + 1$
- orbits on lines of $\text{PG}(3, q)$: lines through $(1, 0, 0, 0)$, lines in π_∞ , $q^2 - 1$ orbits of length $q^2 + q + 1$.
- reconstruct the example, and investigate the intersection properties of the line class and the point orbits.

Some observations

- The $q - 1$ point orbits are third degree surfaces in $\text{PG}(3, q)$.
- The C_3 is generated by the Frobenius automorphism from $\mathbb{F}_{q^3} \rightarrow \mathbb{F}_q$.
- Some examples seems to have a larger automorphism group.

Bruen-Drudge construction

- Choose an elliptic quadric $Q^-(3, q)$ in $PG(3, q)$.
- There are $\frac{(q^2+1)q^2}{2}$ secant lines
- There are $q + 1$ tangent lines through each point of $Q^-(3, q)$, choose half of them for each point
- the secant lines together with the chosen tangent lines is a Cameron-Liebler line class with parameter $x = \frac{q^2+1}{2}$.

Algebraic description

- We use $\mathbb{F}_{q^3}$ to represent $\text{AG}(3, q)$.
- The non-trivial point orbits are now

$$\{\beta u^i \mid \beta \in \mathbb{F}_q \setminus \{0\}, i = 0 \dots q^2 + q\},$$

where u is an element of order $q^2 + q + 1$ in $\mathbb{F}_{q^3}$.

- Notice: the Frobenius automorphism from $\mathbb{F}_{q^3} \rightarrow \mathbb{F}_q$ stabilizes the point orbits.

Combinatorics of the third degree surface

Suppose $q \neq 3^h$

- lines through 0: $q^2 + q + 1$
- lines at ∞ : $q^2 + q + 1$
- lines meeting in 0 points: $\frac{q^2 - q - 2}{3}(q^2 + q + 1)$
- lines meeting in 1 point: $\frac{q^2 - q - 2}{2}(q^2 + q + 1)$
- lines meeting in 2 points: $(q + 1)(q^2 + q + 1)$
- lines meeting in 3 points: $\frac{q^2 - q - 2}{6}(q^2 + q + 1)$

Combinatorics of the third degree surface

Suppose $q \neq 3^h$

- lines through 0: $q^2 + q + 1$
- lines at ∞ : $q^2 + q + 1$
- lines meeting in 0 points: $\frac{q^2 - q - 2}{3}(q^2 + q + 1)$
- lines meeting in 1 point: $\frac{q^2 - q - 2}{2}(q^2 + q + 1)$
- lines meeting in 2 points: $(q + 1)(q^2 + q + 1)$
- lines meeting in 3 points: $\frac{q^2 - q - 2}{6}(q^2 + q + 1)$

Combinatorics of the third degree surface

Suppose $q = 3^h$

- lines through 0: $q^2 + q + 1$: meet in a point of multiplicity 3.
- lines at ∞ : $q^2 + q + 1$
- lines meeting in 0 points: $\frac{q^2 - q - 2}{3}(q^2 + q + 1)$
- lines meeting in 1 point: $\frac{q^2 - q}{2}(q^2 + q + 1)$ (never multiplicity 3)
- lines meeting in 2 points: $q(q^2 + q + 1)$
- lines meeting in 3 points: $\frac{q^2 - q}{6}(q^2 + q + 1)$

Combinatorics of the third degree surface

Suppose $q = 3^h$

- lines through 0: $q^2 + q + 1$: meet in a point of multiplicity 3.
- lines at ∞ : $q^2 + q + 1$
- lines meeting in 0 points: $\frac{q^2 - q - 2}{3}(q^2 + q + 1)$
- lines meeting in 1 point: $\frac{q^2 - q}{2}(q^2 + q + 1)$ (never multiplicity 3)
- lines meeting in 2 points: $q(q^2 + q + 1)$
- lines meeting in 3 points: $\frac{q^2 - q}{6}(q^2 + q + 1)$

Combinatorics of the third degree surface

Suppose $q \neq 3^h$

- lines through 0: $q^2 + q + 1$
- lines at ∞ : $q^2 + q + 1$
- lines meeting in 0 points: $\frac{q^2 - q - 2}{3}(q^2 + q + 1)$
- lines meeting in 1 point: $\frac{q^2 - q - 2}{2}(q^2 + q + 1)$
- lines meeting in 2 points: $(q + 1)(q^2 + q + 1)$
- lines meeting in 3 points: $\frac{q^2 - q - 2}{6}(q^2 + q + 1)$

Combinatorics of the third degree surface

Suppose $q \neq 3^h$

- lines through 0: $q^2 + q + 1$
- lines at ∞ : $q^2 + q + 1$
- lines meeting in 0 points: $\frac{q^2 - q - 2}{3}(q^2 + q + 1)$
- lines meeting in 1 point: $\frac{q^2 - q - 2}{2}(q^2 + q + 1)$
- lines meeting in 2 points: $(q + 1)(q^2 + q + 1)$
- lines meeting in 3 points: $\frac{q^2 - q - 2}{6}(q^2 + q + 1)$

Combinatorics of the third degree surface

Suppose $q \neq 3^h$: all lines behave the same:

- 0 points of $\frac{q-2}{3}$ surfaces
- 1 point of $\frac{q-2}{2}$ surfaces
- 2 points of 1 surface
- 3 points of $\frac{q-2}{6}$ surfaces

Combinatorics of the third degree surface

Suppose $q = 3^h$: two line types:

Type (I): q^2 lines:

- 0 points of $\frac{q-3}{3}$ surfaces
- 1 point of $\frac{q-1}{2}$ surfaces
- 2 points of 1 surface
- 3 points of $\frac{q-3}{6}$ surfaces

Type (II): q lines:

- 0 points of $\frac{2q-3}{3}$ surfaces
- 3 points of $\frac{q}{3}$ surfaces

Final objectives

- describe infinite families of Cameron-Liebler line classes for $q = 2^h$, $x = \frac{(q+1)^2}{3}$
- describe infinite families of Cameron-Liebler line classes for $q = 3^h$, $x = \frac{q^2-1}{2}$
- investigate new examples: $q = 27$, $x = \frac{(q+1)^2}{2}$, this is probably also a member of an infinite family.
- describe more infinite families for $q = p^h$, $p \notin \{2, 3\}$.