

Pseudocomplementation in (normal) subgroup lattices

Tom De Medts Marius Tărnăuceanu

February 2, 2009

Abstract

The goal of this paper is to study finite groups admitting a pseudocomplemented subgroup lattice (PK-groups) or a pseudocomplemented normal subgroup lattice (PKN-groups). In particular, we obtain a complete classification of finite PK-groups and of finite nilpotent PKN-groups. We also study groups with a Stone normal subgroup lattice, and we classify finite groups for which every subgroup has a Stone normal subgroup lattice. Finally, we obtain a complete classification of finite groups for which every subgroup is monolithic.

MSC2000: primary 06D15, 20D30; secondary 20D15, 20D20

Key words: pseudocomplement, subgroup lattice, normal subgroup lattice, monolithic group, Stone lattice

1 Introduction

It is an interesting question in group theory in how far the structure of the subgroup lattice of a group determines the structure of the group itself. This question in its pure form is quite old [11, 2], and M. Suzuki spent his early research years on this problem [14, 15]. Since then, many characterizations and classifications have been obtained for groups for which the subgroup lattice or normal subgroup lattice has certain lattice-theoretic properties. The possibly most famous result in this direction is Ore's result that a group

is locally cyclic if and only if its lattice of subgroups is distributive [9]. We refer to Suzuki's book [17], Schmidt's book [12] or the more recent book [23] by the second author for more information about this theory.

In the current paper, we investigate (finite) groups which have a subgroup lattice or normal subgroup lattice which is pseudocomplemented; these are called *PK-groups* and *PKN-groups*, respectively. We obtain a complete classification of finite PK-groups and of finite nilpotent PKN-groups, thereby vastly improving earlier results [5, 19]. We will also classify groups with the property that each subgroup is itself a PKN-group, and we will give some results about groups for which the normal subgroup lattice is a so-called Stone lattice. In particular, we will classify finite groups for which the normal subgroup lattice of every subgroup is a Stone lattice, and as a corollary, we obtain a complete classification of finite groups for which every subgroup is monolithic.

Let (L, \wedge, \vee) be a bounded lattice with top 1 and bottom 0, and let $a \in L$. An element $b \in L$ is called a *complement* of a if $a \vee b = 1$ and $a \wedge b = 0$, and the lattice L is called *complemented* if every element of L has a complement. An element $a^* \in L$ is called a *pseudocomplement* of a if the following two conditions are satisfied:

- (i) $a \wedge a^* = 0$;
- (ii) $a \wedge x = 0$ ($x \in L$) implies $x \leq a^*$.

Any element of L can have at most one pseudocomplement. We say that L is a *pseudocomplemented lattice* if every element of L has a pseudocomplement. Note that the terminology is slightly misleading, since a complement is not necessarily a pseudocomplement; in fact, a complement of an element need not be unique.

In a pseudocomplemented lattice L the set $S(L) = \{x^* \mid x \in L\}$ forms a lattice (called the *skeleton* of L), which is a \wedge -subsemilattice of L and in which the join is defined by $x \sqcup y = (x \vee y)^{**} = (x^* \wedge y^*)^*$. The lattice $(S(L), \wedge, \sqcup)$ is in fact a boolean lattice [7, Theorem I.6.4]; this result was first proved for complete distributive lattices by V. Glivenko [6].

A lattice L is called *distributive* if the identity

$$a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$$

holds for all $a, b, c \in L$. A pseudocomplemented distributive lattice L for which the skeleton $S(L)$ is a sublattice of L is called a *Stone lattice*.

Let (G, \cdot, e) be a group (where e denotes the identity of G). Then the set $L(G)$ consisting of all subgroups of G is a complete bounded lattice with top G and bottom $\{e\}$, called the *subgroup lattice* of G . The *normal subgroup lattice* $N(G)$ of G is the modular sublattice of $L(G)$ constituted by all normal subgroups of G . Many classes of groups determined by different properties of their subgroup lattices or normal subgroup lattices (such as modularity, distributivity, complementation, . . . and so on) have been identified. In the present paper our aim is to investigate the groups G for which the lattice $L(G)$ or $N(G)$ is pseudocomplemented.

The paper is organized as follows. In Section 2 we study the class of finite PK-groups, i.e. the groups with pseudocomplemented subgroup lattice. In Section 3 we look at finite PKN-groups, i.e. the groups with pseudocomplemented normal subgroup lattice. Section 4 deals with groups for which all subgroups have pseudocomplemented lattices of normal subgroups. In the final section 5, finite groups whose normal subgroup lattices are Stone lattices will be investigated.

Most of our notation is standard and will not be repeated here. Basic definitions and results on lattices and groups can be found in [3, 7] and [1, 18], respectively. For subgroup lattice concepts we refer the reader to [12, 17, 23].

We only mention that by a *generalized quaternion group*, we mean a group of order 2^t for some natural number t , defined by the presentation

$$Q_{2^t} = \langle a, b \mid a^{2^{t-2}} = b^2, a^{2^{t-1}} = 1, b^{-1}ab = a^{-1} \rangle$$

and *not* the more general notion of groups of order $4n$ of the form $Q_{4n} = \langle a, b \mid a^n = b^2, a^{2n} = 1, b^{-1}ab = a^{-1} \rangle$ which is used by some authors.

2 PK-groups

Definition 2.1. A group G is called a *PK-group* if its lattice of subgroups $L(G)$ is pseudocomplemented, and it is called a *PKN-group* if its lattice of normal subgroups $N(G)$ is pseudocomplemented.

PK-groups were introduced and studied in [19] and [20]. Our goal in this section is to give a complete classification of finite PK-groups. For the sake

of readability, we will include a complete proof, even though some parts have already appeared (in a different form) in [19] or [20].

Theorem 2.2. *Let G be a finite group of order $2^t m$ with m odd. Then G is a PK-group if and only if it is cyclic or it is isomorphic to the direct product of a cyclic group of order m and a generalized quaternion group of order 2^t .*

We start with the following easy but important lemma.

Lemma 2.3. (i) *Every subgroup of a PK-group is again a PK-group;*
(ii) *every cyclic group is a PK-group;*
(iii) *a group of order p^2 , p prime, is a PK-group if and only if it is cyclic;*
(iv) *every generalized quaternion group is a PK-group.*

Proof. (i) Let G be PK-group and let $H \leq G$. Let A be an arbitrary subgroup of H and denote its pseudocomplement in G by A^* . Then $A^* \cap H$ is a pseudocomplement of A in H .

(ii) Let G be a finite cyclic group, and let A be an arbitrary subgroup of G . Let s be the largest divisor of $|G|$ coprime to $|A|$, and let A^* be the unique subgroup of G of order s . Then A^* is the pseudocomplement of A in G .

Now let G be the infinite cyclic group $(\mathbb{Z}, +)$; then any two non-trivial subgroups of G intersect non-trivially. So let A be an arbitrary subgroup of G , then $A^* = \{0\}$ is a pseudocomplement of A in G .

(iii) Let G be a PK-group of order p^2 , and assume that G is not cyclic. Then $G \cong C_p \times C_p$, hence G has $p + 1$ subgroups of order p . Take any one of them, say A , and let A^* be its pseudocomplement. Now take two distinct subgroups $B_1, B_2 \leq G$ of order p different from A ; then $A \cap B_i = 1$, hence by the definition of A^* we have $B_i \leq A^*$ implying $G = \langle B_1, B_2 \rangle \leq A^*$ and hence $A \cap A^* = A \neq 1$, a contradiction.

(iv) Let G be a generalized quaternion group. It is well known that such a group contains a unique involution z ; see, for example, [1, Chapter 8, Exercise 3(7)]. But then every non-trivial subgroup A of G contains z , and hence $A^* = 1$ is a pseudocomplement for A in G . \square

Corollary 2.4. *Let G be a finite PK-group. Then every Sylow subgroup of G is either cyclic or generalized quaternion.*

Proof. Let p be any prime divisor of $|G|$, and let S be a Sylow p -subgroup of G ; by Lemma 2.3(i and iii), every subgroup of S of order p^2 is cyclic. Equivalently, S has a unique subgroup of order p , since the center $Z(S)$ of S always contains a subgroup of order p . If p is odd, this implies that S is cyclic, and if $p = 2$, this implies that S is either cyclic or generalized quaternion; see, for example, [1, Chapter 8, Exercise 4]. \square

Before we proceed, we show that the PK and PKN properties behave well with respect to direct products of groups of coprime order.

Proposition 2.5. *Let $G \cong G_1 \times \cdots \times G_n$ with $\gcd(|G_i|, |G_j|) = 1$ for all $i, j \in \{1, \dots, n\}$ with $i \neq j$.*

- (i) *G is a PK-group if and only if each G_i is a PK-group;*
- (ii) *G is a PKN-group if and only if each G_i is a PKN-group.*

Proof. We only prove (ii); the proof of (i) is similar. So assume first that G is a PKN-group, let $i \in \{1, 2, \dots, n\}$ and let $N \trianglelefteq G_i$. Then $N \trianglelefteq G$, hence N has a pseudocomplement N^* in G ; it is clear that $N^* \cap G_i$ is a pseudocomplement of N in G_i .

Conversely, assume that each G_i is a PKN-group, and let $N \trianglelefteq G$; since $\gcd(|G_i|, |G_j|) = 1$ for all $i, j \in \{1, \dots, n\}$ with $i \neq j$, we can write $N = N_1 \times \cdots \times N_n$, where $N_i \trianglelefteq G_i$, for all $i \in \{1, \dots, n\}$. For each i , let N_i^* be a pseudocomplement of N_i in G_i . Then $N^* = N_1^* \times \cdots \times N_n^*$ is a pseudocomplement of N in G . \square

Remark 2.6. It is clear from Lemma 2.3(iii) that the coprimeness condition cannot be omitted.

We are now ready to prove Theorem 2.2. We are grateful to the anonymous referee of an earlier version of this article for the short argument; our original proof was much more involved.

Proof of Theorem 2.2. Assume first that G is either cyclic or isomorphic to the direct product of a cyclic group of order m and a generalized quaternion group of order 2^t . Then it is clear from Lemma 2.3(ii and iv) together with Proposition 2.5(i) that G is a PK-group.

Assume now that G is a PK-group. We will show that G is nilpotent; the result will then follow immediately from Corollary 2.4 since a finite nilpotent group is the direct product of its Sylow subgroups.

So let P be an arbitrary Sylow p -subgroup of G for some prime divisor p of $|G|$, and let N be a pseudocomplement for P in G . Moreover, let R be the subgroup of G generated by all p' -elements of G ; then R is characteristic in G . Since N contains every p' -element of G , we have $R \leq N$. Because $N \cap P = 1$, however, we have $R \cap P = 1$ as well, and hence R intersects every Sylow p -subgroup trivially. We conclude that R is a Hall p' -subgroup of G , and since it is normal, it is, in fact, a normal p -complement.

This holds for every prime divisor p of $|G|$, and hence G has a normal p -complement for each of its prime divisors. It follows that every Sylow subgroup of G is normal (as it is the intersection of normal Hall subgroups), and thus G is nilpotent. \square

3 PKN-groups

In contrast to the situation of PK-groups, a (normal) subgroup of a PKN-group is not necessarily a PKN-group. For example, the dihedral group D_8 of order 8 is a PKN-group, but has a normal subgroup $C_2 \times C_2$, which is not a PKN-group. This simple fact makes the study of PKN-groups considerably harder.

Some of the results in this section have already appeared in [19, 20, 21, 22], but we believe it is interesting to include them nevertheless, and the proofs are somewhat more polished than in *loc. cit.*

We start with an easy but useful observation.

Proposition 3.1. *Let G be a finite PKN-group. Then the center $Z(G)$ of G is cyclic.*

Proof. Let $H \leq Z(G)$ be arbitrary; then $H \trianglelefteq G$, so it has a pseudocomplement H^* in $N(G)$. It is clear that $Z(G) \cap H^*$ is a pseudocomplement of H in $L(Z(G))$. Hence $Z(G)$ is a PK-group; the result now follows from Theorem 2.2 since $Z(G)$ is abelian. \square

Proposition 2.5(ii) shows that the PKN property behaves well with respect to direct products of coprime order. For arbitrary direct products, we still have the following facts.

Lemma 3.2. *Let $G = G_1 \times \cdots \times G_n$ be a PKN-group. Then*

- (i) *each G_i is a PKN-group;*
- (ii) *if $N \trianglelefteq G$ is such that $N \cap G_i = 1$ for all i , then $N = 1$.*

Proof. (i) Let $i \in \{1, 2, \dots, n\}$ and let $N \trianglelefteq G_i$. Then $N \trianglelefteq G$, hence N has a pseudocomplement N^* in G ; it is clear that $N^* \cap G_i$ is a pseudocomplement of N in G_i .

- (ii) Let N^* be the pseudocomplement of N in G . Since $N \cap G_i = 1$, we have $G_i \leq N^*$ for all i , hence $N^* = G$ and since $N \cap N^* = 1$ this implies $N = 1$. \square

Remark 3.3. By a well-known result of J. Wiegold [26], a group has a complemented lattice of normal subgroups (such a group is called an nD -group – see [24]) if and only if it is a direct product of simple groups. So an nD -group is not always a PKN-group. A sufficient condition for a direct product G of simple groups to become a PKN-group is that no two its minimal abelian factors are isomorphic. In this case $N(G)$ is a boolean lattice (see [25, Lemma 1]) and hence a pseudocomplemented lattice.

In any normal subgroup lattice $N(G)$ we have $1^* = G$ and $G^* = 1$. A special type of PKN-groups G is obtained when the skeleton $S(N(G))$ only contains the subgroups 1 and G .

Definition 3.4. We say that a PKN-group G is *elementary* if $S(N(G)) = \{1, G\}$.

A characterization of finite elementary PKN-groups is given by the following result.

Proposition 3.5. *A finite group G is an elementary PKN-group if and only if it is a monolithic group, i.e. a group with a unique minimal normal subgroup.*

Proof. Suppose that G is an elementary PKN-group and assume that it has at least two minimal normal subgroups M_1 and M_2 . Because $M_1 \cap M_2 = 1$, it follows that $M_2 \leq M_1^*$ and therefore $M_1^* \neq 1$, a contradiction. Conversely, suppose that G is a monolithic group and let M be its unique minimal normal subgroup. Then for every $H \trianglelefteq G$ with $H \neq 1$, we have $M \leq H$. This shows that $H^* = 1$. Hence G is an elementary PKN-group. \square

Remark 3.6. This result is no longer true for infinite groups, since an infinite group may have no minimal normal subgroups at all. For example, the infinite cyclic group $(\mathbb{Z}, +)$ is elementary PKN, but has no minimal normal subgroups.

Remark 3.7. Using Proposition 3.5, many classes of elementary PKN-groups can be obtained. For example, any group whose normal subgroup lattice is a chain is an elementary PKN-group. In particular, simple groups, symmetric groups, cyclic p -groups or finite groups of order $p^n q^m$ (p, q distinct primes) with cyclic Sylow subgroups and trivial center [12, Exercise 3, p. 497] are all elementary PKN-groups; see also Theorem 5.9 and Theorem 5.12 below. We also mention that any proper semidirect product of type $G = \langle a \rangle M$, where M is a maximal normal subgroup of G having a fully ordered lattice of normal subgroups, is an elementary PKN-group.

Obviously, elementary PKN-groups G have the property that the lattice $S(N(G))$ is a sublattice of $N(G)$. Our next result gives a characterization of finite PKN-groups for which this property holds.

Proposition 3.8. *For a finite PKN-group G , the following two properties are equivalent:*

- (a) $S(N(G))$ is a sublattice of $N(G)$;
- (b) G is a direct product of elementary PKN-groups.

Proof. For each normal subgroup $N \trianglelefteq G$, we will denote its pseudocomplement in G by N^* . We claim that (a) is equivalent with the condition

$$(A \cap B)^* = A^* B^* \quad \text{for all } A, B \in S(N(G)). \quad (3.1)$$

Indeed, observe that $S(N(G)) = \{A \trianglelefteq G \mid A^{**} = A\}$. If $S(N(G))$ is a sublattice of $N(G)$, then for all $A, B \in S(N(G))$, we have $AB = A \sqcup B =$

$(A^* \cap B^*)^*$, so by replacing A with A^* and B with B^* we obtain (3.1). Conversely, if (3.1) holds, then for all $A, B \in S(N(G))$, we have $AB = A^{**}B^{**} = (A^* \cap B^*)^* \in S(N(G))$, so $S(N(G))$ is a sublattice of $N(G)$.

(a) \Rightarrow (b). To prove (b), we will use induction on $|G|$. Let M_1, M_2, \dots, M_s be the minimal normal subgroups of G . If $s = 1$, then G itself is an elementary PKN-group, so we may assume $s \geq 2$. Then $M_1^* \cap M_1^{**} = 1$, and by equation (3.1),

$$G = 1^* = (M_1^* \cap M_1^{**})^* = M_1^{**}M_1^{***} = M_1^{**}M_1^*,$$

hence G is the direct product of M_1^{**} and M_1^* . In particular, every minimal normal subgroup of M_1^{**} is also a minimal normal subgroup of G .

Now $M_i \leq M_1^*$, for all $i \in \{2, \dots, s\}$, hence $M_i \cap M_1^{**} = 1$. Therefore M_1^{**} has a unique minimal normal subgroup, namely M_1 , and hence M_1^{**} is an elementary PKN-group.

By Lemma 3.2(i), M_1^* is again a PKN-group. It remains to show that $S(N(M_1^*))$ is a sublattice of $N(M_1^*)$, since the result will then follow by the induction hypothesis. We have

$$S(N(M_1^*)) = \{K^* \cap M_1^* \mid K \in N(G)\} = \{A \cap M_1^* \mid A \in S(N(G))\}.$$

For any two elements $A \cap M_1^*$ and $B \cap M_1^*$ of $S(N(M_1^*))$, we have

$$\begin{aligned} (A \cap M_1^*) \cap (B \cap M_1^*) &= (A \cap B) \cap M_1^* \in S(N(M_1^*)), \\ (A \cap M_1^*)(B \cap M_1^*) &= (AB) \cap M_1^* \in S(N(M_1^*)), \end{aligned}$$

where we have used the fact that $S(N(G))$ is a distributive lattice. This shows that $S(N(M_1^*))$ is a sublattice of $N(M_1^*)$, as claimed.

(b) \Rightarrow (a). Let $G = G_1 \times \dots \times G_n$, where each G_i is an elementary PKN-group. Let $K \in S(N(G))$ be arbitrary, and let $i \in \{1, \dots, n\}$. Since $K \cap G_i \in S(N(G_i)) = \{1, G_i\}$, we have either $K \cap G_i = 1$ (and then $G_i \leq K^*$) or $G_i \leq K$ (and then $K^* \cap G_i = 1$). Let

$$\begin{aligned} X &= \{i \in \{1, \dots, n\} \mid K \cap G_i = 1\} = \{i \in \{1, \dots, n\} \mid G_i \leq K^*\}, \\ Y &= \{i \in \{1, \dots, n\} \mid K^* \cap G_i = 1\} = \{i \in \{1, \dots, n\} \mid G_i \leq K\}, \end{aligned}$$

and write $A = \prod_{i \in X} G_i$ and $B = \prod_{i \in Y} G_i$. Then $G = A \times B$; by Lemma 3.2(i), both A and B are PKN-groups. Observe that $B \leq K$. By the modular law,

$$K = K \cap G = K \cap (AB) = B(K \cap A).$$

However, by Lemma 3.2(ii) applied on the PKN-group A , we have $K \cap A = 1$. Hence $K = B = \prod_{i \in Y} G_i$.

Now take $K_1, K_2 \in S(N(G))$; then there are subsets Y and Z of $\{1, \dots, n\}$ such that $K_1 = \prod_{i \in Y} G_i$ and $K_2 = \prod_{i \in Z} G_i$. Since $S(N(G))$ is always a \wedge -subsemilattice of $N(G)$, we have $K_1 \cap K_2 \in S(N(G))$ (in fact $K_1 \cap K_2 = \prod_{i \in Y \cap Z} G_i$). On the other hand, it is obvious that $K_1 K_2 = \prod_{i \in Y \cup Z} G_i \in S(N(G))$. Hence $S(N(G))$ is a sublattice of $N(G)$. \square

Remark 3.9. If the equivalent properties of Proposition 3.8 are satisfied, then $S(N(G))$ is in fact a direct product of chains of length 1; this follows from the proof of (b) \Rightarrow (a).

A complete classification of finite PKN-groups seems to be out of reach at this point, but we are able to classify all finite *nilpotent* PKN-groups.

Theorem 3.10. *Let G be a finite nilpotent group. Then G is a PKN-group if and only if its center $Z(G)$ is cyclic.*

Proof. If G is a PKN-group, then $Z(G)$ is cyclic by Proposition 3.1. So assume that G is a finite nilpotent group with cyclic center. Then G is isomorphic to the direct product of its Sylow subgroups, $G \cong \prod_{i=1}^k G_i$, where each G_i is a p_i -group with cyclic center. By Proposition 2.5(ii), G is a PKN-group if and only if each G_i is a PKN-group.

So we may assume without loss of generality that G is a p -group with cyclic center $Z(G)$. But then the unique subgroup A of $Z(G)$ of order p is in fact the unique minimal normal subgroup of G . By Proposition 3.5, G is a PKN-group (in fact an elementary PKN-group). \square

Corollary 3.11. *Let G be a finite nilpotent PKN-group. Then*

- (i) G is the direct product of elementary PKN-groups;
- (ii) $S(N(G))$ is a sublattice of $N(G)$.

Proof. Part (i) follows from the fact that a PKN- p -group is elementary PKN; part (ii) then follows from Proposition 3.8. \square

4 PKN*-groups

As we have observed, the class of PKN-groups is not closed under (normal) subgroups. It therefore makes sense to introduce the following class of groups.

Definition 4.1. A group G is called a *PKN*-group* if all of its subgroups are PKN-groups.

Our goal in this section is to classify finite PKN*-groups.

Theorem 4.2. *Let G be a finite group. Then the following are equivalent.*

- (a) G is a PKN*-group;
- (b) for each prime divisor p of $|G|$, every Sylow p -subgroup of G has a unique subgroup of order p ;
- (c) all Sylow subgroups of G of odd order are cyclic and all Sylow 2-subgroups of G are cyclic or generalized quaternion.

Proof. The fact that (b) and (c) are equivalent is well known; see, for example, [1, Chapter 8, Exercise 4]. So it suffices to show that (a) and (b) are equivalent.

Assume first that G is a PKN*-group; then every subgroup of order p^2 (with p prime) is itself a PKN-group, and hence such a subgroup is cyclic by Lemma 2.3(iii). As in the proof of Corollary 2.4, it follows that every Sylow p -subgroup has a unique subgroup of order p . This shows that (a) implies (b).

Conversely, suppose that G is a finite group such that every Sylow p -subgroup of G has a unique subgroup of order p . We claim that for every two normal subgroups $U, V \trianglelefteq G$, the equivalence

$$U \cap V = 1 \iff \gcd(|U|, |V|) = 1 \tag{4.1}$$

holds. Indeed, if $\gcd(|U|, |V|) = 1$, then of course $U \cap V = 1$. So assume $\gcd(|U|, |V|) \neq 1$, then there is a prime p dividing both $|U|$ and $|V|$. Let P

be a Sylow p -subgroup of G , and let A and B be Sylow p -subgroups of U and V , respectively. Then there exist $x, y \in G$ such that $1 \neq A^x \leq P$ and $1 \neq B^y \leq P$. Since P has exactly one subgroup of order p , this implies that $A^x \cap B^y \neq 1$ and so $U \cap V = U^x \cap V^y \neq 1$. This proves the equivalence (4.1).

Now, let N be an arbitrary normal subgroup of G , and let U be the largest normal subgroup of G with order coprime to $|N|$. Then of course $N \cap U = 1$, and by (4.1), every subgroup of G which intersects N trivially, has order coprime to $|N|$ and is therefore contained in U . Hence U is a pseudocomplement of N in G , and we conclude that G is a PKN-group. Since the condition (b) is inherited to subgroups, it follows that every subgroup of G is a PKN-group as well, and hence G is a PKN*-group. This shows that (b) implies (a). \square

Remark 4.3. The finite groups with all Sylow subgroups cyclic, are always solvable, and these groups have, in fact, a very precise structure. They are known as ZM-groups; see also Definition 5.5 and Theorem 5.8 below. An important (lattice theoretical) property of the ZM-groups is that these groups are exactly the finite groups whose poset of conjugacy classes of subgroups forms a distributive lattice [4, Theorem A].

On the other hand, the groups satisfying condition (c) of Theorem 4.2 are not always solvable, but the non-solvable ones are also well understood, by the following deep result by Suzuki.

Theorem 4.4 (Suzuki [16]). *Let G be a non-solvable finite group such that every Sylow subgroup of G is either cyclic or generalized quaternion. Then G contains a normal subgroup G_1 such that $[G : G_1] \leq 2$ and $G_1 \cong Z \times L$, where Z is a solvable group whose Sylow subgroups are all cyclic, and $L \cong \mathrm{SL}(2, p)$ for some odd prime p .*

In particular, for each prime p , the group $\mathrm{SL}(2, p)$ is a PKN*-group.

5 Groups whose normal subgroup lattices are Stone lattices

In this final section we present some results concerning finite groups G for which the lattice $N(G)$ is a Stone lattice, i.e. a distributive pseudocomplemented lattice such that $S(N(G))$ is a sublattice of $N(G)$. As a corollary,

we will obtain a complete classification of the finite groups for which every subgroup is monolithic, i.e. has a unique minimal normal subgroup.

We mention that the structure of groups with distributive lattices of normal subgroups, the so-called *DLN-groups*, is not known, but there are some characterizations of these groups (see [10] or [12, §9.1]). We only mention the following theorem, which is a generalization of [10, Theorem 4.1].

Theorem 5.1. *Let $G = G_1 \times \cdots \times G_k$ be a finite group. Then G is a DLN-group if and only if each direct factor G_i is a DLN-group and for all $i \neq j$, there are no central chief factors in G_i and in G_j the orders of which coincide.*

We are now able to establish the following characterization of finite groups whose normal subgroup lattice is a Stone lattice.

Theorem 5.2. *Let G be a finite group. Then $N(G)$ is a Stone lattice if and only if G is a direct product $G = G_1 \times \cdots \times G_k$ of monolithic DLN-groups G_i , and for all $i \neq j$, there are no central chief factors in G_i and in G_j the orders of which coincide.*

Proof. Assume first that $N(G)$ is a Stone lattice. Then G is a PKN-group and $S(N(G))$ is a sublattice of $N(G)$, so by Proposition 3.8, G is a direct product of monolithic groups: $G = G_1 \times \cdots \times G_k$. Since G is a DLN-group, the conclusion now follows from Theorem 5.1.

Conversely, assume that $G = G_1 \times \cdots \times G_k$, where each G_i is a monolithic DLN-group, and for all $i \neq j$, there are no central chief factors in G_i and in G_j the orders of which coincide. Then by Theorem 5.1, $N(G)$ is a distributive lattice. We now claim that G is a PKN-group. Indeed, let $N \trianglelefteq G$. By distributivity of $N(G)$, we have $N = \prod_{i=1}^k (N \cap G_i)$. Without loss of generality, we may assume that $N \cap G_i \neq 1$ for $i \in \{1, \dots, s\}$ and $N \cap G_i = 1$ for $i \in \{s+1, \dots, k\}$; in particular, $N \leq \prod_{i=1}^s G_i$. Let $N^* = \prod_{i=s+1}^k G_i$; we claim that N^* is a pseudocomplement of N in $N(G)$. Indeed, we clearly have $N \cap N^* = 1$. Assume that $A \trianglelefteq G$ is such that $N \cap A = 1$. For each $i \in \{1, \dots, s\}$, the group G_i is monolithic, so since $N \cap G_i \neq 1$ we must have $A \cap G_i = 1$. But then by distributivity, $A = \prod_{i=1}^k (A \cap G_i) \leq N^*$, which proves that N^* is a pseudocomplement of N as claimed. It now follows from Proposition 3.8 that $S(N(G))$ is a sublattice of $N(G)$, and hence $N(G)$ is a Stone lattice. \square

Remark 5.3. The structure of finite monolithic DLN-groups is not known either. This class of groups obviously includes the cyclic p -groups, but there exist a lot of non-cyclic groups which are monolithic DLN-groups (for example, the non-cyclic elementary PKN-groups presented in Remark 3.7). Nevertheless, it is easy to see that the cyclic p -groups are the only finite p -groups that are monolithic DLN-groups. Moreover, because a direct product of k cyclic p -groups is a PKN-group if and only if $k = 1$, the cyclic p -groups are also the only finite p -groups G for which $N(G)$ is a Stone lattice.

The previous remark allows us to classify finite nilpotent groups whose normal subgroup lattice is a Stone lattice.

Corollary 5.4. *Let G be a finite nilpotent group. Then $N(G)$ is a Stone lattice if and only if G is cyclic.*

Proof. Since G is nilpotent, it can be written as a direct product of its Sylow subgroups: $G = G_1 \times \cdots \times G_k$. If G is cyclic, then each of the groups G_i is a cyclic p_i -group, and hence a monolithic DLN-group; it follows from Theorem 5.2 that $N(G)$ is a Stone lattice.

Conversely, assume that $N(G)$ is a Stone lattice. Since the orders of the Sylow subgroups are coprime, Theorem 5.2 implies that each of the lattices $N(G_i)$ is a Stone lattice; by Remark 5.3, each G_i is cyclic. Hence G is also cyclic. \square

We will now investigate finite groups for which the normal subgroup lattice of every subgroup is a Stone lattice. We start with a reduction; let us first recall the following definition which we have encountered in Remark 4.3.

Definition 5.5. A finite group G is called a *Zassenhaus metacyclic group*, or *ZM-group* for short, if all Sylow subgroups of G are cyclic.

Proposition 5.6. *Let G be a finite group. Then the following two properties are equivalent:*

- (a) $N(H)$ is a Stone lattice for all subgroups H of G ;
- (b) $G \cong G_1 \times \cdots \times G_k$, where the G_i are monolithic ZM-groups of coprime orders, with the property that $N(H_i)$ is a Stone lattice for all subgroups H_i of G_i .

Proof. (a) \Rightarrow (b). Let H be a Sylow subgroup of G . Then $N(H)$ is a Stone lattice and therefore, by Corollary 5.4, H is cyclic. Thus G is a ZM-group. Since $N(G)$ is a Stone lattice, Theorem 5.2 implies that G can be written as a direct product of monolithic DLN-groups G_i , $i \in \{1, \dots, k\}$. Since every subgroup of a ZM-group is itself a ZM-group, the G_i are in fact ZM-groups. Suppose that there are $i \neq j$ such that $|G_i|$ and $|G_j|$ have a common prime divisor p . Then there are a subgroup M_i of order p in G_i , and a subgroup M_j of order p in G_j . By our hypothesis, the normal subgroup lattice of $H = M_i \times M_j \cong C_p \times C_p$ is a Stone lattice, a contradiction.

(b) \Rightarrow (a). Let H be an arbitrary subgroup of G . Since the direct factors G_i have coprime orders, $H \cong H_1 \times \dots \times H_k$ for certain subgroups $H_i \leq G_i$. Clearly H is a ZM-group, so by Remark 4.3 it is a DLN-group. Since $N(H_i)$ is a Stone lattice for each i , Proposition 3.8 implies that each H_i is a direct product of elementary PKN-groups, and hence the same is true for H . Moreover, each H_i is a PKN-group, and since the H_i have coprime orders, Proposition 2.5(ii) implies that H is a PKN-group. We conclude that $N(H)$ is a Stone lattice. \square

In view of Proposition 5.6(b), we will now digress on monolithic ZM-groups. The structure of ZM-groups has been completely determined by Zassenhaus.

Definition 5.7. A triple (m, n, r) satisfying the conditions

$$\gcd(m, n) = \gcd(m, r - 1) = 1 \quad \text{and} \quad r^n \equiv 1 \pmod{m}$$

will be called a *ZM-triple*, and the corresponding group

$$\langle a, b \mid a^m = b^n = 1, b^{-1}ab = a^r \rangle$$

will be denoted by $\mathbf{ZM}(m, n, r)$.

Theorem 5.8 (Zassenhaus). *Let G be a ZM-group. Then there exists a ZM-triple (m, n, r) such that $G \cong \mathbf{ZM}(m, n, r)$. We have $|G| = mn$ and $G' = \langle a \rangle$ (so $|G'| = m$), and G/G' is cyclic of order n .*

Conversely, every group isomorphic to $\mathbf{ZM}(m, n, r)$ is a ZM-group.

Proof. See, for example, [8, IV, Satz 2.11]. \square

We can now determine the structure of monolithic ZM-groups.

Theorem 5.9. *Let $G \cong \text{ZM}(m, n, r)$ be a ZM-group. Then the following properties are equivalent:*

- (a) G is monolithic;
- (b) either $m = 1$ and n is a prime power, or m is a prime power and $r^d \not\equiv 1 \pmod{m}$ for all $1 \leq d < n$;
- (c) either $|G|$ is a prime power, or $|G'|$ is a prime power and $Z(G) = 1$.

Proof. (a) \Rightarrow (b). Assume that $G \cong \text{ZM}(m, n, r)$ is monolithic. If $m = 1$, then G is cyclic of order n ; for each prime divisor p of n , G has a minimal normal subgroup of order p ; hence n is a prime power.

So assume that $m \neq 1$, and let p be a prime divisor of m . By the defining relations of G in Definition 5.7, we see that every subgroup of $\langle a \rangle$ is normalized by b and thus is a normal subgroup of G . In particular, G has a minimal normal subgroup of order p .

Hence m can have only one prime divisor, i.e. m is a prime power; let N be the minimal normal subgroup contained in $\langle a \rangle$. Suppose now that there is some $d < n$ such that $r^d \equiv 1 \pmod{m}$. Since $r^n \equiv 1 \pmod{m}$ as well, we may in fact assume that $d \mid n$; assume furthermore that n/d is prime. Let $H = \langle b^d \rangle$; this is a subgroup of G of prime order. By the defining relation $ab = ba^r$, we get

$$ab^d = b^d a^{r^d} = b^d a$$

since $r^d \equiv 1 \pmod{m}$; this shows that $b^d \in Z(G)$, so in particular H is a minimal normal subgroup of G , different from N . This contradicts the fact that G is monolithic.

- (b) \Rightarrow (c). If $m = 1$ and n is a prime power, then $|G| = mn$ is a prime power. So assume that $m = |G'|$ is a prime power and $r^d \not\equiv 1 \pmod{m}$ for all $1 \leq d < n$. Let $g = b^s a^t$ with $0 \leq s \leq n - 1$ and $0 \leq t \leq m - 1$ be an arbitrary element of G ; we compute

$$[g, a] = a^{1-r^s}, \quad [g, b] = a^{t(r-1)}.$$

Assume now that $g \in Z(G)$; then we must have $r^s \equiv 1 \pmod{m}$ and $m \mid t(r-1)$. Our assumption $r^d \not\equiv 1 \pmod{m}$ for all $1 \leq d < n$ implies

$s = 0$; the fact that $\gcd(m, r - 1)$ and $t < m$ implies $t = 0$. Hence $Z(G) = 1$.

(c) \Rightarrow (a). If $|G|$ is a prime power, then G is a cyclic group of prime power order and hence it is monolithic. So assume that $|G'|$ is a prime power and $Z(G) = 1$. Then G' has a unique subgroup P of prime order, which is therefore a minimal normal subgroup of G . Let $N \trianglelefteq G$ be an arbitrary non-trivial normal subgroup of G ; it is sufficient to show that $N \cap G' \neq 1$, since this will imply that $P \leq N$. So suppose that $N \cap G' = 1$. Then $[N, G] = 1$, and hence $N \leq Z(G)$. But now $Z(G) = 1$ implies $N = 1$, a contradiction. \square

We now claim that every monolithic ZM-group has the property that the normal subgroup lattice of any subgroup is a Stone lattice. This statement is obvious when the group is cyclic of prime power order, so we only consider non-abelian ZM-groups.

Theorem 5.10. *Let $G \cong \text{ZM}(m, n, r)$ be a monolithic non-abelian ZM-group with $m = p^k$ for some prime p and some $k \geq 1$. Then*

- (i) $n \mid p - 1$;
- (ii) *the order of each element of G is a divisor of m or a divisor of n ;*
- (iii) *$N(H)$ is a Stone lattice for each subgroup H of G . More precisely, every subgroup of G is either cyclic or monolithic.*

Proof. Let $G \cong \text{ZM}(m, n, r)$ be a monolithic ZM-group with generators a and b as in Definition 5.7. By Theorem 5.9, $m = p^k$ for some prime p and some number $k \geq 1$ since we assume G to be non-abelian.

- (i) By Theorem 5.9(b), the order of r modulo p^k is precisely n . In particular, $n \mid \phi(p^k) = p^{k-1}(p - 1)$, where ϕ is the Euler totient function; since $\gcd(n, p) = 1$, we have $n \mid p - 1$.
- (ii) By the defining relations in Definition 5.7, it is not very hard to compute, using induction on d , that

$$(b^s a^t)^d = b^{sd} a^{t(1+r^s+r^{2s}+\dots+r^{(d-1)s})}, \quad (5.1)$$

for all natural numbers s, t, d . Suppose that G contains an element $g = b^s a^t$ of order pq , where q is a prime dividing n . Then b^s must have

order q , hence we have $s = c \cdot n/q$ for some $c \in \{1, \dots, q-1\}$. Let $x = r^s$; then the order of x modulo p^k is q . We have

$$g^q = a^{t(1+x+x^2+\dots+x^{(q-1)})}, \quad g^{pq} = a^{t(1+x+x^2+\dots+x^{(pq-1)})}.$$

Since g has order pq , this implies

$$t(1+x+x^2+\dots+x^{(q-1)}) \not\equiv 0 \pmod{p^k}, \quad (5.2)$$

$$t(1+x+x^2+\dots+x^{(pq-1)}) \equiv 0 \pmod{p^k}. \quad (5.3)$$

Also observe that by (i), we have $q \mid p-1$, so in particular $x^p \equiv x \pmod{p^k}$. Let $y = 1+x+\dots+x^{p-1}$; then $y(x-1) = x^p - 1 \equiv x - 1 \pmod{p^k}$, hence $p^k \mid (y-1)(x-1)$. However, $x \not\equiv 1 \pmod{p^k}$, and therefore $p \mid y-1$. In particular, $\gcd(y, p) = 1$.

Since $x^p \equiv x \pmod{p^k}$, we have

$$\begin{aligned} 1+x+x^2+\dots+x^{(pq-1)} \\ &= (1+x+x^2+\dots+x^{(p-1)})(1+x^p+x^{2p}+\dots+x^{(q-1)p}) \\ &\equiv y(1+x+x^2+\dots+x^{(q-1)}) \pmod{p^k}; \end{aligned}$$

since $\gcd(y, p) = 1$, this implies the equivalence

$$\begin{aligned} t(1+x+x^2+\dots+x^{(pq-1)}) &\equiv 0 \pmod{p^k} \\ \iff t(1+x+x^2+\dots+x^{(q-1)}) &\equiv 0 \pmod{p^k}. \end{aligned}$$

This contradicts equations (5.2) and (5.3), and (ii) follows.

- (iii) Let H be an arbitrary subgroup of G . If H is abelian, then it is cyclic, so Corollary 5.4 implies that $N(H)$ is a Stone lattice. So we may assume that H is non-abelian; in particular, the derived subgroup H' is non-trivial, and since $H' \leq G' = \langle a \rangle$, this implies that $g = a^{p^{k-1}} \in H$, and therefore $p \mid |H|$. If p were the only prime dividing $|H|$, then H would be cyclic, contradicting the fact that H is non-abelian; hence there is some prime $q \mid n$ with $q \mid |H|$. Let $h \in H$ be an element of order q .

Suppose that $Z(H) \neq 1$. If $Z(H)$ contains an element x of order p , then xh has order pq , contradicting (ii). If $Z(H)$ does not contain elements of order p , then it must contain an element y of order q' for some prime $q' \mid n$, but then yg has order pq' , again contradicting (ii). We conclude that $Z(H) = 1$, and hence by Theorem 5.9, H is a monolithic ZM-group; in particular, $N(H)$ is a Stone lattice. \square

We now obtain the complete classification of all groups with the property that each subgroup has a Stone normal subgroup lattice as an easy corollary.

Theorem 5.11. *Let G be a finite group. Then the following two properties are equivalent:*

- (a) $N(H)$ is a Stone lattice for all subgroups H of G ;
- (b) $G \cong G_1 \times \cdots \times G_k$, where the G_i are monolithic ZM-groups of coprime orders.

Proof. This follows immediately from Proposition 5.6 and Theorem 5.10. \square

We end this section by classifying all finite groups such that every subgroup is monolithic.

Theorem 5.12. *Let G be a finite group; then the following two properties are equivalent:*

- (a) every subgroup of G is monolithic;
- (b) either G is a cyclic group of order p^k , or G is a ZM-group of order $p^k q^\ell$ with $Z(G) = 1$, where p, q are distinct primes.

Proof. (a) \Rightarrow (b). Assume that every subgroup of G is monolithic. Then every Sylow subgroup of G is cyclic, hence G is a monolithic ZM-group, say $G \cong \text{ZM}(m, n, r)$. Assume that $|G|$ is not a prime power; then by Theorem 5.9, $Z(G) = 1$ and m is a prime power p^k . On the other hand, the group $\langle b \rangle$ is a cyclic group of order n ; such a group can only be monolithic if n is a prime power q^ℓ .

(b) \Rightarrow (a). If G is a cyclic group of prime power order, then each of its subgroups is monolithic; so assume that G is a ZM-group of order $p^k q^\ell$ with $Z(G) = 1$. By Theorem 5.9, G is monolithic. Now let H be an arbitrary subgroup of G . Then by Theorem 5.10(iii), H is cyclic or monolithic. If it is monolithic, we are done, so assume H is cyclic. By Theorem 5.10(ii), the order of each element is either a power of p or a power of q , so it follows that H is a cyclic p -group or a cyclic q -group. In both cases, H is monolithic. \square

References

- [1] M. Aschbacher, “Finite group theory”, Second edition, Cambridge Studies in Advanced Mathematics **10**, Cambridge University Press, Cambridge, 2000.
- [2] R. Baer, The Significance of the System of Subgroups for the Structure of the Group, *Amer. J. Math.* **61** (1939), no. 1, 1–44.
- [3] G. Birkhoff, “Lattice theory”, Amer. Math. Soc., Providence, R.I., 1967.
- [4] R. Brandl, G. Cutolo and S. Rinauro, Posets of subgroups of groups and distributivity, *Boll. U.M.I. (7)* **9-A** (1995), 217–223.
- [5] Gr. G. Călugăreanu, Abelian groups with pseudocomplemented lattice of subgroups, *Studia Univ. Babeş-Bolyai Cluj-Napoca (3)* **31** (1986), 39–41.
- [6] V. Glivenko, Sur quelques points de la logique de M. Brouwer, *Bull. Acad. Sc. Belgique* **15** (1929), 183–188.
- [7] G. Grätzer, “General lattice theory”, Academic Press, New York, 1978.
- [8] B. Huppert, “Endliche Gruppen I”, Die Grundlehren der Mathematischen Wissenschaften, Band **134**, Springer-Verlag, Berlin, New York, 1967.
- [9] O. Ore, Structures and group theory II, *Duke Math. J.* **4** (1938), 247–269.
- [10] G. Pazderski, On groups for which the lattice of normal subgroups is distributive, *Beiträge zur Algebra und Geometrie* **24** (1987), 185–200.
- [11] A. Rottlaender, Nachweis der Existenz nicht-isomorpher Gruppen von gleicher Situation der Untergruppen, *Math. Z.* **28** (1928), no. 1, 641–653.
- [12] R. Schmidt, “Subgroup lattices of groups”, de Gruyter Expositions in Mathematics **14**, de Gruyter, Berlin, 1994.
- [13] R. Schmidt, New results and methods in the theory of subgroup lattices of groups, *Quaderni di Mat.* **8** (Topics in Infinite Groups), Eds. M. Curzio, F. De Giovanni, 2000, 239–275.

- [14] M. Suzuki, On the lattice of subgroups of finite groups, *Trans. Amer. Math. Soc.* **70** (1951), 345–371.
- [15] M. Suzuki, On the L -homomorphisms of finite groups, *Trans. Amer. Math. Soc.* **70** (1951), 372–386.
- [16] M. Suzuki, On finite groups with cyclic Sylow subgroups for all odd primes, *Amer. J. Math.* **77** (1955), 657–691.
- [17] M. Suzuki, “Structure of a group and the structure of its lattice of subgroups”, *Ergebnisse der Mathematik und ihrer Grenzgebiete, Neue Folge, Heft 10*, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1956.
- [18] M. Suzuki, “Group theory”, I, II, Springer-Verlag, Berlin, 1982, 1986.
- [19] M. Tărnăuceanu, On groups whose lattices of subgroups are pseudo-complemented, *Fuzzy Systems & Artificial Intelligence (2)* **10** (2004), 45–49.
- [20] M. Tărnăuceanu, Pseudocomplemented groups, *An. Univ. “Al.I. Cuza”, Iași, Math., (1)* **51** (2005), 201–205.
- [21] M. Tărnăuceanu, Complementation in subgroup lattices, *Sci. Ann. Univ. Agric. Sci. Vet. Med.* **49** (2006), 285–303.
- [22] M. Tărnăuceanu, Complementation in normal subgroup lattices, *Sci. Ann. Univ. Agric. Sci. Vet. Med.* **49** (2006), 305–322.
- [23] M. Tărnăuceanu, “Groups determined by posets of subgroups”, Ed. Matrix Rom, București, 2006.
- [24] K. H. Toh, Groups with normal subgroups possessing subnormal complements, *Proc. Amer. Math. Soc. (2)* **41** (1973), 378–380.
- [25] G. Whitson, Finite groups whose subgroup, composition subgroup, or normal subgroup lattice is an ortholattice, *Algebra Universalis* **8** (1978), 123–127.
- [26] J. Wiegold, On direct factors in groups, *J. London Math. Soc.* **35** (1960), 310–320.

Tom De Medts
Department of Pure Mathematics
and Computer Algebra
Ghent University
Galglaan 2, B-9000 Gent, Belgium
e-mail: tdemedts@cage.UGent.be

Marius Tărnăuceanu
Faculty of Mathematics
“Al.I. Cuza” University
Iași, Romania
e-mail: tarnauc@uaic.ro