

James Parkinson and Hendrik Van Maldeghem*

Automorphisms and opposition in spherical buildings of classical type

DOI 10.1515/advgeom-2024-0012. Received 12 March, 2023

Abstract: An automorphism of a spherical building is called domestic if it maps no chamber to an opposite chamber. In this paper we classify domestic automorphisms of spherical buildings of classical type.

Keywords: Classical spherical building, polar space, opposition diagram, domestic automorphism.

2010 Mathematics Subject Classification: 20E42, 51E24, 51A50, 20E45



Communicated by: R. Weiss

Introduction

The study of the geometry of fixed elements of automorphisms of spherical buildings is a well-established and beautiful topic; see [9]. Recently a complementary theory concerning the “opposite geometry” $\text{Opp}(\theta)$, consisting of those elements mapped to opposite elements by an automorphism, has been developed; see [11; 12; 13; 14]. The generic situation is that this opposite geometry is rather large, and typically contains many chambers of the building. The more delicate and interesting situation is when the opposite geometry $\text{Opp}(\theta)$ contains no chamber, in which case the automorphism θ is called *domestic*. This paper is part of series of papers directed towards classifying, as precisely as possible, the class of domestic automorphisms of spherical buildings, with the focus of this paper being on the buildings of classical types.

Central to the study of the domesticity and the opposite geometry is the notion of the *opposition diagram* $\text{Diag}(\theta)$ of an automorphism θ , introduced in [11; 12]. This diagram encodes the types of the simplices of the building that are mapped onto opposite simplices by θ , and one surprising aspect of the theory is that there are relatively few permitted opposition diagrams for a given spherical building. This opens up the possibility of classifying the class of domestic automorphisms with each diagram. In [13] and [14] we carried out this program for split spherical buildings of exceptional type (with the exception of the three opposition diagrams $E_{7,3}$, $E_{7,4}$ and $E_{8,4}$ that will be dealt with in future work), and for all Moufang hexagons. In the present paper we carry out the program for spherical buildings of classical type (both split and non-split) of rank at least 3.

Before stating our main results, we briefly expand on some of the above concepts (see Section 1 for remaining concepts). Let Δ be a thick spherical building of irreducible type (W, S) with Coxeter diagram Γ , and let θ be an automorphism of Δ . Let $\pi \in \text{Aut}(\Gamma)$ denote the automorphism of Γ induced by θ (thus $\pi = \text{id}$ if and only if θ is type preserving). Let $\pi_0 \in \text{Aut}(\Gamma)$ be the automorphism of Γ induced by the longest element w_0 . The *distinguished orbits* are the nontrivial subsets of S that are minimal subject to being preserved by $\pi_0 \circ \pi$. In particular, the distinguished orbits are all singletons if and only if $\pi_0 \circ \pi = \text{id}$. The *opposition diagram* $\text{Diag}(\theta)$ of θ is drawn by encircling those distinguished orbits $J \subseteq S$ of Γ such that there exists a type J simplex mapped onto an opposite simplex. The opposition diagram $\text{Diag}(\theta)$ is drawn “straight” if $\pi_0 \circ \pi = \text{id}$, and “bent” in the usual way otherwise.

Thus, for example, the two opposition diagrams (a)  (b)  represent (a) a non-type preserving automorphism of a D_7 building mapping type 2 and type 4 vertices to opposite vertices, and (b) a type preserving automorphism of a D_7 building mapping type 2 and type 4 vertices to opposite ones,

James Parkinson, University of Sydney, Sydney, Australia, email: jamesp@maths.usyd.edu.au

*Corresponding author: Hendrik Van Maldeghem, Ghent University, Ghent, Belgium, email: Hendrik.VanMaldeghem@UGent.be

and type $\{6, 7\}$ simplices onto opposite simplices (recall that the opposition relation π_0 on D_n is type preserving if n is even, and interchanges types $n - 1$ and n if n is odd).

In [11; 12] we showed that if θ is an automorphism of a thick irreducible spherical building then $\text{Diag}(\theta)$ belongs to a very restricted list of possible diagrams. These diagrams are called the *admissible diagrams*, and are defined by simple combinatorial rules. We will not require the formal definition of admissible diagrams, instead the explicit list of admissible diagrams of classical type given in Tables 1 and 2 is sufficient for the purposes of this paper (see [11] for the formal details).

Symbol	Diagram	Conditions
${}^2A_{n,i}^1$		$0 \leq i \leq n/2$
${}^1A_{n,(n-1)/2}^2$		n odd
${}^1A_{n,n}^1$		
${}^1B_{n,i}^1$ or ${}^1C_{n,i}^1$		$0 \leq i \leq n$
${}^1B_{n,i}^2$ or ${}^1C_{n,i}^2$		$0 \leq i \leq n/2$
${}^1D_{n,i}^1$		$0 \leq i < n$ and $i = n \pmod 2$
${}^1D_{n,n}^1$		
${}^2D_{n,i}^1$		$0 \leq i < n - 1$ and $i = n + 1 \pmod 2$
${}^2D_{n,n-1}^1$		
${}^1D_{n,i}^2$		n even and $0 \leq i < n/2$
${}^1D_{n,n/2}^2$		n even
${}^2D_{n,i}^2$		n odd and $0 \leq i < (n - 1)/2$
${}^2D_{n,(n-1)/2}^2$		n odd

Table 1: Admissible diagrams of classical type, general rank

As in the tables, each admissible diagram of classical type is denoted by a symbol ${}^tX_{n,i}^j$, where

- (1) $X \in \{A, B, C, D\}$ is the Coxeter type of Γ , and n is the rank;
- (2) $t \in \{1, 2, 3\}$ is the order of the automorphism $\pi_0 \circ \pi$ of Γ (the “twisting”);
- (3) i is the number of distinguished orbits encircled in the diagram; and
- (4) $j \in \{1, 2\}$ is the (graph) distance between successive encircled distinguished orbits in J .

Symbol	Diagram	Symbol	Diagram	Symbol	Diagram
${}^2B_{2,1}^1$ or ${}^2C_{2,1}^1$		${}^3D_{4,1}^2$		${}^3D_{4,2}^1$	

Table 2: Admissible diagrams of classical type, special rank

Domestic automorphisms of generalised polygons (rank 2 buildings) are studied in [19; 10], and domestic trialities of buildings of type D_4 are studied in [22]; the diagrams in Table 2 are not discussed further in this paper. For the diagrams ${}^tX_{n,i}^j$ in Table 1 it turns out that the twisting index $t \in \{1, 2\}$ is determined by the indices n, i, j (and X). This is clear if $X \in \{A, B, C\}$, and for type D note that $t \in \{1, 2\}$ is determined by the condition $t \equiv n + ij + 1 \pmod{2}$. With this in mind, it is usually convenient to omit the twisting index t from the notation ${}^tX_{n,i}^j$, without risk of confusion.

An automorphism θ is called *capped* if whenever there exist both type J_1 and J_2 simplices in $\text{Opp}(\theta)$, then there exists a type $J_1 \cup J_2$ simplex in $\text{Opp}(\theta)$. In [11, Theorem 1] we proved that every automorphism of a large spherical building is capped (a thick spherical building of rank at least 3 is called *large* if it contains no Fano plane residues, and *small* otherwise). Since automorphisms of small buildings are studied in [12] we shall, when required, restrict attention to large spherical buildings in this paper, and by doing so we may thus assume that all automorphisms are capped. In this case an automorphism is domestic if and only if its opposition diagram has at least one distinguished orbit not encircled.

Domestic automorphisms of large buildings of type A_n have been completely classified in previous work (see Theorem 1.5 of this paper for a summary). Thus the main focus of this paper is on buildings of types B_n, C_n and D_n , where the situation is considerably more complicated. Such buildings may naturally be regarded in a standard and uniform way as *polar spaces* $\Pi = (\mathcal{P}, \Omega)$ of rank n , where \mathcal{P} is the set of points of Π (the type 1 vertices of the building in standard Bourbaki labelling), and Ω is the set of singular subspaces of Π (see Section 1.6 for more details). The singular subspaces of (projective) dimension $n - 1$ are called the *maximal singular subspaces*.

Since we consider thick buildings, the associated polar spaces have thick lines (meaning that each line contains at least 3 points). We adopt the convention that buildings of type C_n are those corresponding to symplectic polar spaces (that is, split buildings of type C_n), buildings of type D_n are those corresponding to the non-thick polar spaces (meaning that each singular $(n - 2)$ -space is contained in precisely 2 maximal singular subspaces), and buildings associated to all other polar spaces are considered to be of type B_n (however note that this convention is somewhat non-standard as buildings of absolute type 2A_n are often considered as “type C_n ” elsewhere). Thus for buildings of type B_n and C_n the type i vertices ($1 \leq i \leq n$) correspond to the singular subspaces of projective dimension $i - 1$. This is also true for buildings of type D_n for $1 \leq i \leq n - 2$, while the type $n - 1$ and n vertices of the building correspond to the two types of maximal singular subspaces. Points x, y of a polar space are opposite one another if and only if they are distinct and not collinear. Singular subspaces $X, Y \in \Omega$ of a polar space are opposite one another if and only if they have the same dimension and for each point $x \in X$ there is a point $y \in Y$ with x and y opposite.

The automorphisms of spherical buildings with opposition diagrams in Table 1 are all collineations of the associated polar space Π (however, note that in the D_n case, the automorphism θ may interchange the type $n - 1$ and n vertices of the building). A collineation θ of a polar space is called *point-domestic* if it maps no point (equivalently, no type 1 vertex of the building) to an opposite. More generally, θ is called *i -domestic* ($0 \leq i \leq n - 1$) if it maps no singular subspace of projective dimension i to an opposite (hence point-domestic is equivalent to 0-domestic). Domestic automorphisms of rank 3 polar spaces are classified in [18, Theorem 7.2], and so, when required, we may restrict our attention to rank at least 4, and moreover, from the comments above, we may restrict to large polar spaces.

It is convenient to divide the class of domestic collineations of a polar space into three mutually disjoint classes, as follows.

Definition. A domestic collineation θ of a polar space is said to be

- (1) of class I if it is not point-domestic.
- (2) of class II if it is point-domestic, and has a fixed point.
- (3) of class III if it is point-domestic, and has no fixed points.

Note that if θ is of class I, then by the classification of admissible diagrams, we have $\text{Diag}(\theta) = X_{n,i}^1$, where $1 \leq i < n$ (with $i < n - 1$ in the $X = D$ case). Similarly, if θ is of class II or class III then $\text{Diag}(\theta) = X_{n,i}^2$ with $0 \leq i \leq n/2$ (where $X \in \{B, C, D\}$).

The main results of this paper are summarised as follows (for undefined notions, see Section 1). To begin with, consider class I collineations of a polar space $\Pi = (\mathcal{P}, \Omega)$. We first show that such automorphisms are characterised by big fixed point sets (see Definition 1.8).

Theorem 1. *If θ is a class I automorphism of a large polar space Π with diagram $\text{Diag}(\theta) = X_{n,i}^1$ (with $X \in \{B, C, D\}$) then θ fixes pointwise a subspace of corank i . In particular, the fixed point set of θ is a possibly degenerate and possibly non-thick polar space (possibly with lines of size 2). Conversely, if θ is a collineation of Π mapping at least one point to an opposite and fixing pointwise a subspace of corank i and fixing no subspace of corank $j < i$, then $\text{Diag}(\theta) = X_{n,i}^1$ (with $X \in \{B, C, D\}$).*

On an intuitive level, Theorem 1 says that domestic automorphisms that are not point-domestic are i -domestic for sufficiently large i for the following reason: by fixing a subspace of suitably high dimension it is forced that each singular subspace of suitably high dimension i contains a fixed point, and thus is not mapped onto an opposite singular subspace.

In certain cases we are able to be more precise. For example, we prove the following (see Definition 1.8 for the definition of elations).

Theorem 2. *Let $\Pi = C_{n,1}(\mathbb{K})$ be the (large) symplectic polar space over a field \mathbb{K} with $n \geq 3$ and let θ be a collineation of Π . Then:*

- (1) $\text{Diag}(\theta) = C_{n,1}^1$ if and only if θ is a central elation (that is, a long root elation).
- (2) $\text{Diag}(\theta) = C_{n,2}^1$ if and only if θ is either
 - (a) a product of two nontrivial perpendicular long root elations, or
 - (b) a nontrivial member of the group generated by two opposite long root groups and not conjugate to a long root elation.

In a similar way we also classify those collineations of thick non-embeddable polar spaces with diagram $C_{3,1}^1$ (see Theorem 2.4) and those of split polar spaces with diagram $B_{n,1}^1$ and $D_{n,1}^1$ (see Proposition 2.5, Theorem 2.6 and Remark 2.7 for the statements).

The situation for class II and class III automorphisms of polar spaces (that is, point-domestic collineations) is more complicated, but richer. Recall that the diagrams of such (nontrivial) automorphisms are of the form $X_{n,i}^2$ with $X \in \{B, C, D\}$ and $1 \leq i \leq \lfloor n/2 \rfloor$. The automorphisms whose diagrams have the fewest encircled nodes (that is $i = 1$) admit a complete classification.

Theorem 3. *Let $\Pi = (\mathcal{P}, \Omega)$ be a polar space of rank $n \geq 3$ and let θ be a collineation of Π .*

- (1) *If Π is of symplectic type then θ has opposition diagram $C_{n,1}^2$ if and only if θ is an axial collineation (in the case $\text{char}(\mathbb{K}) = 2$) or a specific involutive homology (if $\text{char}(\mathbb{K}) \neq 2$).*
- (2) *If Π is not of symplectic type then θ has diagram $X_{n,1}^2$, $X \in \{B, D\}$, if and only if θ is an axial elation (and so Π is an orthogonal polar space), or the rank is 3 and θ is an ideal Baer collineation.*

In the split case (that is, for those arising from Chevalley groups over commutative fields), it is possible to completely classify the point-domestic automorphisms with diagram not equal to one of the “extreme diagrams” $B_{n,n/2}^2$, $C_{n,n/2}^2$, $D_{n,n/2}^2$ (with n even) or $D_{n,(n-1)/2}^2$ (with n odd).

Theorem 4. *Let θ be a point-domestic collineation of a parabolic, symplectic (with characteristic not 2), or hyperbolic polar space Δ with opposition diagram $B_{n,i}^2$, $1 \leq i \leq (n-1)/2$, $C_{n,i}^2$, $1 \leq i \leq (n-1)/2$, or $D_{n,i}^2$, $1 \leq i \leq (n-2)/2$, respectively. Then θ is the product of i pairwise orthogonal long root elations, an $(I_{2n-2i}, -I_{2i})$ -homology, or the product of i pairwise orthogonal long root elations, respectively. The conclusion also holds for $C_{n,n/2}^2$ if it is assumed that θ fixes at least one point.*

Remarkably, when the polar space of rank $n \geq 3$ is Hermitian, there is only one type of automorphisms of class II, and they have opposition diagram $B_{n,m}^2$ with $2m \in \{n, n+1\}$ (for the precise statement, we refer to Proposition 3.18).

The most beautiful situation is a domestic automorphism that fixes no points (class III). These are very restricted, as illustrated by the following theorem. In particular, note that every class III automorphism must have an “extreme” opposition diagram $X_{n,n/2}^2$ with n even and $X \in \{B, C, D\}$.

Theorem 5. *Let Δ be a spherical building of type B_n, C_n or D_n with $n \geq 4$ and let $\Pi = (\mathcal{P}, \Omega)$ be the associated polar space defined over the field \mathbb{K} . Suppose that there exists a point-domestic collineation θ of Π with no fixed points. Let \mathcal{Q} denote the set of fixed lines of θ , and let Ω' denote the set of fixed singular subspaces. Then*

- (1) n is even,
- (2) $\text{Diag}(\theta) = X_{n,n/2}^2$ with $X \in \{B, C, D\}$, and
- (3) the pair $\Pi' = (\mathcal{Q}, \Omega')$ is a polar space defined over either a quadratic extension of \mathbb{K} , or over a quaternion division algebra \mathbb{H} which is 2-dimensional over \mathbb{K} (and \mathbb{K} is 2-dimensional over the centre of \mathbb{H}).

In the split case we can be even more explicit.

Theorem 6. *Let Δ be a split polar space of rank $n \geq 4$ over a field \mathbb{K} . Let θ be a point-domestic collineation of Δ with no fixed points. Then n is even, and Δ is either a symplectic polar space (type C_n), or a hyperbolic polar space (type D_n), and θ has opposition diagram $C_{n,n/2}^2$ or $D_{n,n/2}^2$, respectively. Moreover, the fixed element structure of θ is either*

- (1) a symplectic polar space over a quadratic extension of \mathbb{K} , or
- (2) a minimal Hermitian polar space, or a mixed polar space,

respectively. In each case, the points of the fixed element structure are the lines of Δ that are fixed by θ .

We can be more precise in each case, as follows.

Theorem 7. *Let Δ be a symplectic polar space of rank $n \geq 4$ defined over a field \mathbb{K} . Let θ be a point-domestic collineation of Δ with no fixed points. Let \mathbb{L} be the quadratic extension of \mathbb{K} over which the fixed element structure of θ is defined (according to Theorem 6). Then θ is an involution, and if $\text{char}(\mathbb{K}) = 2$ then \mathbb{L} is an inseparable extension of \mathbb{K} . Conversely, if \mathbb{K} admits a quadratic extension \mathbb{L}/\mathbb{K} , inseparable in the case $\text{char}(\mathbb{K}) = 2$, then Δ admits a point-domestic collineation θ without fixed points as above.*

Theorem 8. *Let Δ be a hyperbolic polar space of rank $n \geq 4$ defined over a field \mathbb{K} . Let θ be a point-domestic collineation of Δ with no fixed points. Let $\text{PG}(2n - 1, \mathbb{K})$ be the ambient projective space. Then θ naturally extends to $\text{PG}(2n - 1, \mathbb{K})$ pointwise fixing a spread \mathcal{S} which defines a projective space $\text{PG}(n - 1, \mathbb{L})$ over a quadratic extension \mathbb{L} of \mathbb{K} , and every collineation of $\text{PG}(2n - 1, \mathbb{K})$ which pointwise fixes \mathcal{S} is a point-domestic collineation of Δ without fixed points. Moreover:*

- (1) If \mathbb{L}/\mathbb{K} is separable then the fixed element polar space (cf. Theorem 6) is minimal Hermitian.
- (2) If \mathbb{L}/\mathbb{K} is inseparable then the fixed element polar space is the subspace of the symplectic polar space over \mathbb{L} obtained by restricting the long root relations to \mathbb{K} .

Conversely, if \mathbb{K} admits a quadratic extension \mathbb{L}/\mathbb{K} then there exists a point-domestic collineation of Δ without fixed points as above.

For non-split polar spaces the situation depends highly on the underlying field and the form defining the polar space. The neatest case is as follows.

Theorem 9. *Let Δ be a minimal Hermitian polar space of rank $n \geq 4$ defined over a field \mathbb{K} , with corresponding field involution σ and field extension \mathbb{K}/\mathbb{F} . Let θ be a point-domestic collineation of Δ with no fixed points. Let $\text{PG}(2n - 1, \mathbb{K})$ be the ambient projective space. Then θ is an involution and naturally extends to $\text{PG}(2n - 1, \mathbb{K})$ pointwise fixing a spread \mathcal{S} which defines a projective space $\text{PG}(n - 1, \mathbb{H})$ over a quaternion division algebra \mathbb{H} over its centre \mathbb{F} , with \mathbb{K} a 2-dimensional subalgebra of \mathbb{H} . The only nontrivial collineation of Δ fixing every member of \mathcal{S} in Δ is θ . The fixed element polar space is a minimal quaternion polar space related to a non-standard involution of \mathbb{H} pointwise fixing a 3-dimensional subalgebra over \mathbb{F} .*

Conversely, if \mathbb{K} admits a quaternion extension \mathbb{H} such that the centre \mathbb{F} of \mathbb{H} is a subfield of \mathbb{K} and \mathbb{K}/\mathbb{F} is a separable quadratic field extension, then there exists a point-domestic collineation of Δ without fixed points as above.

The remaining cases are orthogonal and Hermitian polar spaces whose standard form has a nontrivial anisotropic part. Since these contain the hyperbolic and minimal Hermitian polar spaces fixed under the given point-domestic collineation, the previous two theorems give necessary conditions for the existence of collineations of class III. We provide examples, and some instances where such collineations do not exist (see Section 3.3).

In this paper we shall also classify those admissible diagrams that can be obtained as the opposition diagram of a unipotent element of a split spherical building (that is, an element conjugate to an element of U^+), extending the corresponding result [13, Theorem 5] for buildings of exceptional type. In [13] we introduced a combinatorial notion of an admissible diagram being *polar closed*. See Section 1.4 for the definition, however for now we note that in the classical case all admissible diagrams other than $A_{n;(n-1)/2}^2$ (n odd), $A_{n;n}^1$, $B_{n;i}^1$ or $D_{n;i}^1$ (with i odd and $1 \leq i < n$), $C_{n;i}^2$ (with $1 \leq i \leq n/2$) and those in Table 2 turn out to be polar closed. The classification of admissible diagrams arising as the opposition diagram of a unipotent element is then as follows.

Theorem 10. *Let Δ be a split irreducible spherical building with Dynkin diagram Γ . The admissible Dynkin diagrams of type Γ that can be obtained as opposition diagrams of a unipotent element are precisely the polar closed diagrams.*

In particular, Theorem 10 shows that all polar closed admissible diagrams (of classical type) occur as the opposition diagram of some automorphism of the respective split building (over any field). We will prove the following theorem, completing the proof of [13, Corollary 10], showing that the list of admissible diagrams contains no redundancies.

Theorem 11. *Let Δ be a split spherical building of irreducible type X_n (with the underlying field assumed to be perfect in the case of B_n , C_n or F_4 in characteristic 2 and G_2 in characteristic 3). Every admissible diagram with underlying Dynkin diagram X_n arises as the opposition diagram of some automorphism of Δ .*

We conclude this introduction with a summary of the structure of this paper. Section 1 provides background on spherical buildings, domestic automorphisms, and polar spaces. The main theorems are proved in the Sections 2–4. More precisely, Theorem 1 is proved at the beginning of Section 2, Theorem 2 follows from Theorems 2.2 and 2.3, and Theorem 3 follows from Theorem 3.7 and Proposition 3.11. Theorem 4 follows from Theorems 3.13 and 3.14, Theorem 5 follows from Corollaries 3.4, 3.6, and 3.21. Theorems 6, 7, 8 and 9 are proved in Theorem 3.22, Proposition 3.23, Proposition 3.24, and Proposition 3.25, respectively. Theorems 10 and 11 are proved in Section 4.

1 Background and definitions

This section contains background on Coxeter groups, spherical buildings, domestic automorphisms, opposition diagrams, Chevalley groups, projective spaces and polar spaces.

1.1 Coxeter groups and spherical buildings

Let (W, S) be a spherical Coxeter system with Coxeter diagram $\Gamma = \Gamma(W, S)$. For $J \subseteq S$ let W_J be the parabolic subgroup generated by J , and let $\Gamma_J = \Gamma(W_J, J)$ be the associated subgraph of Γ . For each $J \subseteq S$ let w_J be the longest element of W_J , and write $w_0 = w_S$. For each $J \subseteq S$ the element w_J induces an automorphism π_J of Γ_J by $\pi_J(s) = w_J s w_J^{-1}$ for $s \in J$. Write $\pi_0 = \pi_S$.

Let Δ be a thick spherical building of type (W, S) , regarded as a simplicial complex, with chamber set $\mathcal{C} = \mathcal{C}(\Delta)$ and W -distance function $\delta : \mathcal{C} \times \mathcal{C} \rightarrow W$. Let $\tau : \Delta \rightarrow 2^S$ be a fixed type map on the simplicial complex Δ

(we adopt Bourbaki [3] conventions for the indexing of the generators of spherical Coxeter systems). Our main references for the theory of buildings are [1; 20], and we assume that the reader is already acquainted with the theory.

Chambers $A, B \in \mathcal{C}$ are *opposite* if they are at maximum distance in the chamber graph, or equivalently if $\delta(A, B) = w_0$. Simplices α, β of Δ are *opposite* if $\tau(\beta) = \pi_0(\tau(\alpha))$ and there exists a chamber A containing α and a chamber B containing β such that A and B are opposite.

The *residue* of a simplex α of Δ is the set of all simplices of Δ which contain α , together with the order relation induced by that on Δ . Then $\text{Res}(\alpha)$ is a building whose Coxeter diagram is obtained from the Coxeter diagram of Δ by removing all nodes which belong to $\tau(\alpha)$.

Let α be a simplex of Δ . The *projection onto α* is the map $\text{proj}_\alpha : \Delta \rightarrow \text{Res}(\alpha)$ defined as follows; see [20, Section 3]. For a simplex β of Δ , we set $\text{proj}_\alpha(\beta)$ to be the unique simplex γ of $\text{Res}(\alpha)$ which is maximal subject to the property that every minimal length gallery from a chamber of $\text{Res}(\beta)$ to $\text{Res}(\alpha)$ ends in a chamber containing γ .

We call the thick irreducible spherical buildings of rank at least 3 with no Fano plane residues *large buildings*, and those containing at least one Fano plane residue are called *small buildings*.

1.2 Automorphisms and opposition diagrams

An *automorphism* of Δ is a simplicial complex automorphism $\theta : \Delta \rightarrow \Delta$. Note that θ does not necessarily preserve types. Indeed, each automorphism $\theta : \Delta \rightarrow \Delta$ induces an automorphism π_θ of Γ by $\delta(A, B) = s$ if and only if $\delta(A^\theta, B^\theta) = \pi_\theta(s)$.

Let θ be an automorphism of Δ . The *opposite geometry* of θ is

$$\text{Opp}(\theta) = \{\alpha \in \Delta \mid \alpha \text{ is opposite } \alpha^\theta\},$$

and the *type* $\text{Typ}(\theta)$ of θ is the union of all subsets $J \subseteq S$ such that there is a type J simplex in $\text{Opp}(\theta)$.

The following fundamental theorem, due to Leeb [8] and Abramenko and Brown [2], shows that $\text{Opp}(\theta)$ is empty if and only if θ is the identity.

Theorem 1.1 ([8; 2]). *Every nontrivial automorphism of a thick spherical buildings maps some simplex onto an opposite simplex.*

The *opposition diagram* $\text{Diag}(\theta)$ of θ is the triple $(\Gamma, \text{Typ}(\theta), \pi_\theta)$, where Γ is the Coxeter diagram of Δ . Less formally, the opposition diagram of θ is depicted by drawing Γ and encircling the nodes of $\text{Typ}(\theta)$, where we encircle nodes in minimal subsets invariant under $\pi_0 \circ \pi_\theta$. We draw the diagram “bent” (in the standard way) if $\pi_0 \circ \pi_\theta \neq 1$ (see the examples in the introduction). An opposition diagram is *empty* if no nodes are encircled (that is, $\text{Typ}(\theta) = \emptyset$), and *full* if all nodes are encircled (that is, $\text{Typ}(\theta) = S$). Note that $\text{Diag}(\theta)$ is empty if and only if θ is the identity (by Theorem 1.1).

Definition 1.2. Let Δ be a spherical building of type (W, S) . Let θ be a nontrivial automorphism of Δ , and let $J \subseteq S$ be stable under the diagram automorphism $\pi_0 \circ \pi_\theta$. Then θ is called

- (1) *capped* if there exists a type $\text{Typ}(\theta)$ simplex in $\text{Opp}(\theta)$, and *uncapped* otherwise;
- (2) *domestic* if $\text{Opp}(\theta)$ contains no chamber;
- (3) *J-domestic* if $\text{Opp}(\theta)$ contains no type J -simplex.

A simplex $\alpha \in \Delta$ is said to be *domestic for θ* if it is not mapped onto an opposite simplex by θ , and is *non-domestic for θ* otherwise. If the automorphism θ is clear from context, we will often simply say α is domestic (or non-domestic).

The requirement that J be stable under $\pi_0 \circ \pi_\theta$ in the above definition is to avoid trivialities. For if J is not stable under $\pi_0 \circ \pi_\theta$ then necessarily there is no type J -simplex mapped to an opposite by θ ; see [11, Lemma 1.3].

The main result of [11] is the following useful fact.

Theorem 1.3 ([11, Theorem 1]). *Every automorphism of a large spherical building is capped.*

Let θ be an automorphism of Δ , and suppose that $\alpha \in \text{Opp}(\theta)$. It follows from [20, Theorem 3.28] that the map $\text{proj}_\alpha : \text{Res}(\alpha^\theta) \rightarrow \text{Res}(\alpha)$ is an isomorphism, and hence we have an isomorphism

$$\theta_\alpha : \text{Res}(\alpha) \xrightarrow{\sim} \text{Res}(\alpha) \quad \text{given by} \quad \theta_\alpha = \text{proj}_\alpha \circ \theta.$$

If π_θ is the automorphism of Γ induced by θ , and if $J = \tau(\alpha)$, then the automorphism of $\Gamma_{S \setminus J}$ induced by θ_α is $\pi_{\theta_\alpha} = \pi_{S \setminus J} \circ \pi_\theta$; see [11, Proposition 1.11]. There is a very useful relationship between domesticity of θ and domesticity of θ_α , as follows (in particular, this relationship facilitates inductive arguments).

Lemma 1.4 ([11, Proposition 1.13]). *Let θ be an automorphism of a spherical building Δ and let $\alpha \in \text{Opp}(\theta)$. If $\beta \in \text{Res}(\alpha)$ then β is opposite β^θ in the building Δ if and only if β is opposite β^{θ_α} in the building $\text{Res}(\alpha)$.*

An admissible diagram $X = (\Gamma, J, \pi)$ is called *type preserving* if $\pi = \text{id}$ (these are the opposition diagrams of type preserving automorphisms). For example, the type preserving diagrams of type D are precisely the diagrams $D_{n,i}^1$ with n and i both even, $D_{n,i}^1$ with n odd and i even, and all diagrams $D_{n,i}^2$.

1.3 Root systems and Chevalley groups

The *split* irreducible spherical buildings are those arising from Chevalley groups over a commutative field \mathbb{K} . In this case there is an associated irreducible crystallographic root system Φ . Let $\alpha_1, \dots, \alpha_n$ be a fixed system of simple roots, and let Φ^+ denote the associated positive roots. Adopting standard notation (from [4; 16]) we write $x_\alpha(a)$ ($\alpha \in \Phi$ and $a \in \mathbb{K}$) for the unipotent generators of the Chevalley group $G = G_\Phi(\mathbb{K})$. Write $U_\alpha = \langle x_\alpha(a) \mid a \in \mathbb{K} \rangle$ for the root group of $\alpha \in \Phi$, and

$$U = \langle U_\alpha \mid \alpha \in \Phi^+ \rangle.$$

By a *unipotent element* we shall mean an element of G conjugate to a member of U .

Let $\Gamma = \Gamma(\Phi)$ be the Dynkin diagram (with the convention that the arrow points towards the short root in the case of double or triple bonds). The *height* of a root $\alpha = k_1\alpha_1 + \dots + k_n\alpha_n$ is $\text{ht}(\alpha) = k_1 + \dots + k_n$, and there is a unique root φ of maximal height (the *highest root of Φ*). The *polar type* of Φ is the subset $\wp \subseteq \{1, 2, \dots, n\}$ given by

$$\wp = \{1 \leq i \leq n \mid \langle \alpha_i, \varphi \rangle \neq 0\}.$$

Thus, in standard Bourbaki labelling, the polar types of A_n, B_n, C_n, D_n are $\{1, n\}, \{2\}, \{1\}, \{2\}$, respectively (a simple way to remember these is to note that the polar type is the set of nodes of Γ to which the additional generator is joined when constructing the associated affine diagram). In the case that $\wp = \{p\}$ is a singleton set, we refer to p as the *polar node*.

The *dual polar type* of a Dynkin diagram is the subset \wp' corresponding to the polar node of the dual diagram. Thus $\wp' = \wp$ in the A_n and D_n cases, while $\wp' = \{1\}$ for B_n , and $\wp' = \{2\}$ for C_n . If $\wp' = \{p'\}$ we refer to p' as the dual polar node.

1.4 Polar closed admissible diagrams

The definition of admissible diagrams naturally extends to admissible Dynkin diagrams (with the additional condition that π preserves arrows on the Dynkin diagram). Let X be an admissible Dynkin diagram, and let $X = X_0, X_1, \dots$ be subdiagrams such that, for $j \geq 1$, the diagram X_j is obtained from X_{j-1} by removing an encircled polar type from one of the connected components of X_{j-1} . This process terminates at some step $j = k$ (that is, X_k has no polar types encircled). We say that X is *polar closed* if X_k is an empty diagram (that is, has no nodes encircled).

For example, the following sequence X_0, X_1, X_2, X_3, X_4 shows that $B_{5,4}^1$ is polar closed



while $B_{5,3}^1$ is not polar closed because

$$X_0 = \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \circ \text{---} \rightarrow X_1 = \text{---} \circ \times \text{---} \circ \text{---} \circ \text{---} \rightarrow X_2 = \text{---} \circ \text{---} \circ \text{---} \circ \text{---}$$

where X_2 is a non-empty diagram with no polar node encircled. In fact, by inspection of the list of diagrams, the admissible Dynkin diagrams of classical type that are not polar closed are precisely the diagrams listed in the introduction.

Suppose that X is polar closed, and let $X = X_0, X_1, \dots, X_k$ be a sequence of subdiagrams obtained by successively removing encircled polar nodes with X_k empty (as in the definition of polar closed). Let $\varphi_1, \dots, \varphi_k \in \Phi^+$ be the highest roots associated to the polar nodes that are removed at each step. For example, the sequence of highest roots corresponding to the above sequence for $B_{5,4}^1$ is (in the basis of simple roots)

$$\varphi_{B_5} = (12222) \mapsto \varphi_{A_1} = (10000) \mapsto \varphi_{B_3} = (00122) \mapsto \varphi_{A_1} = (00100).$$

Note that the sequence X_0, \dots, X_k is not unique (for example, in the $B_{5,4}^1$ sequence given above we may remove the polar node of the B_3 diagram at the second step instead of the A_1 polar node), however it is clear that the set $\{\varphi_1, \dots, \varphi_k\}$ of highest roots is independent of the choices made. Moreover, note that the roots $\varphi_1, \dots, \varphi_k$ are mutually perpendicular (by the definition of the polar type), and hence the group

$$U(X) = \langle U_{\varphi_1}, \dots, U_{\varphi_k} \rangle$$

generated by the associated positive root groups is abelian. We call an element $g \in U(X)$ *generic* if $g = x_{\varphi_1}(a_1) \cdots x_{\varphi_k}(a_k)$ with $a_1, \dots, a_k \neq 0$.

1.5 Projective spaces

Buildings of type A_n are equivalent to projective spaces in a well known way, with type 1 vertices of the building corresponding to points of the projective space, type 2 vertices corresponding to lines, and so on, with type n vertices corresponding to hyperplanes. An automorphism of a projective space is a *collineation* (respectively, *duality*) if the corresponding automorphism of the building of type A_n is type preserving (respectively, interchanges types i and $n - i + 1$ for $i = 1, \dots, n$).

We denote by $PG(n, \mathbb{K})$ the projective space over \mathbb{K} of dimension n over a (possibly noncommutative) field \mathbb{K} . The points, lines, etc, are the 1-spaces, 2-spaces, etc of the vector space \mathbb{K}^{n+1} .

If \mathbb{F} is a subfield of \mathbb{K} , with \mathbb{K} 2-dimensional over \mathbb{F} , then the natural inclusion of $PG(n, \mathbb{F})$ in $PG(n, \mathbb{K})$ is called a *Baer subspace* (however note that Baer subspaces are not “subspaces” in the technical sense of point-line geometry). A *Baer collineation* of $PG(n, \mathbb{K})$ is a collineation that pointwise fixes a Baer subspace.

A *symplectic polarity* of $PG(n, \mathbb{K})$ is a duality θ of the form $U^\theta = \{v \in \mathbb{K}^{n+1} \mid (u, v) = 0 \text{ for all } u \in U\}$, where (\cdot, \cdot) is a non-degenerate symplectic form on \mathbb{K}^{n+1} (necessarily n is odd and \mathbb{K} is commutative; and here U is a subspace).

An $(I_{n-i}, -I_i)$ -homology of $PG(n - 1, \mathbb{K})$, \mathbb{K} a commutative field, is a collineation arising from a linear map in the underlying vector space with diagonal matrix containing $n - i$ times 1 and i times -1 .

Domesticity in buildings of type A_n is well understood (see Theorem 1.5 below).

Theorem 1.5 ([17, Theorem 3.1, 4.3] and [11, Theorem 3.5]). *Let Δ be the building $A_n(\mathbb{K})$, regarded as the projective space $PG(n, \mathbb{K})$, with \mathbb{K} a possibly non-commutative field and $n \geq 2$.*

- (1) *A collineation θ of Δ has opposition diagram $\text{Diag}(\theta) = A_{n,i}^1$ (with $0 \leq i \leq n/2$) if and only if θ pointwise fixes a unique subspace of projective dimension $n - i$ but does not fix any subspace of larger projective dimension pointwise.*
- (2) *If $|\mathbb{K}| > 2$ then Δ admits domestic dualities if and only if n is odd and \mathbb{K} is commutative. In this case, every domestic duality is a symplectic polarity (that is, θ is induced by a non-degenerate symplectic form on \mathbb{K}^{n+1}), with opposition diagram $\text{Diag}(\theta) = A_{n,(n-1)/2}^2$.*

Thus the focus of this paper is on buildings of types B_n, C_n , and D_n .

1.6 Polar spaces

Building of type B_n , C_n , and D_n may all be considered as polar spaces Π , where the points of Π are the type 1 vertices of the building. We recall the basic definitions below.

Definition 1.6 (See [6, Chapter 7]). Let $n > 0$. A *polar space* of rank n is a pair $\Pi = (\mathcal{P}, \mathcal{Q})$ consisting of a set \mathcal{P} (the *points*) and a set \mathcal{Q} of subsets of \mathcal{P} (the *singular subspaces*) satisfying the following axioms:

- (1) A singular subspace $X \in \mathcal{Q}$ together with the singular subspaces contained in it define a (possibly reducible) d -dimensional projective space for some $d \in \{-1, 0, \dots, n-1\}$ (then $d = \dim(X)$ is the *dimension* of X).
- (2) The intersection of any two singular subspaces is again a singular subspace.
- (3) If X is an $(n-1)$ -dimensional singular subspace and if $p \in \mathcal{P} \setminus X$ then there exists a unique $(n-1)$ -dimensional singular subspace Y containing p such that $\dim(X \cap Y) = n-2$. The singular subspace $X \cap Y$ consists of those points q of X for which $\{p, q\}$ is contained in some singular subspace of dimension 1.
- (4) There exist two disjoint singular subspaces of dimension $n-1$.

Let $\Pi = (\mathcal{P}, \mathcal{Q})$ be a polar space of rank n . The singular subspaces of dimension 1 (respectively 2) are called *lines* (respectively *planes*). The singular subspaces of dimension $n-1$ (respectively $n-2$) are called *maximal* (respectively *submaximal*) singular subspaces. A singular subspace of dimension i is called a *singular i -space*. Points $p, q \in \mathcal{P}$ are *collinear* if they are contained in a line; note that if two distinct points are contained in a singular subspace then they are collinear by the first axiom. The (unique) line determined by two distinct points x, y is denoted $\langle x, y \rangle$. The symbol $\langle \cdot \rangle$ will be used throughout and will have the obvious meaning of generation.

Let \mathcal{L} denote the set of lines of $\Pi = (\mathcal{P}, \mathcal{Q})$. Taking incidence to be containment, one may regard Π as a point line geometry $\Pi' = (\mathcal{P}, \mathcal{L})$. Indeed, there is an equivalent set of axioms, due to Buekenhout and Shult, that characterise polar spaces in terms of this incidence structure; see [6, § 7.2]. In particular, Π' satisfies the following condition (the Buekenhout–Shult one-or-all axiom): *If $p \in \mathcal{P}$ and $L \in \mathcal{L}$ are not incident, then either one or all points of L are collinear with p .* The set \mathcal{Q} can be recovered from the point line geometry $\Pi' = (\mathcal{P}, \mathcal{L})$ as the set of all $X \subseteq \mathcal{P}$ that are *subspaces* of Π' (that is, every line that has at least two distinct points in X has all of its points in X) and *singular* (that is, X consists of mutually collinear points). Thus we will identify Π and Π' .

We write $p \perp q$ to indicate that points p and q are collinear and we write p^\perp for the set of all points collinear to p (including p itself). If $X \subseteq \mathcal{P}$ then we write $X^\perp = \bigcap_{p \in X} p^\perp$.

Points $x, y \in \mathcal{P}$ are *opposite* (in the building theoretic sense) one another if and only if they are not collinear. More generally, opposition of singular subspaces is given by the following obvious lemma.

Lemma 1.7. *Singular spaces $X, Y \in \mathcal{Q}$ are opposite if and only if they have the same dimension, are disjoint, and for each point $x \in X$ there is a point $y \in Y$ such that x and y are opposite. In particular, maximal singular subspaces X, Y are opposite if and only if they are disjoint.*

We recall the following definitions.

Definition 1.8. Let Π be a polar space of rank n .

- (1) A *subspace of corank t* , with $0 \leq t \leq n-1$, of a polar space of rank n is a subspace having non-empty intersection with every singular subspace of dimension t , and such that it is disjoint from at least one singular subspace of dimension $t-1$.
- (2) A *geometric hyperplane* is a subspace of corank 1, that is, a subspace with the property that every line contains at least one point of the subspace, and at least one line contains exactly one point of it. It is called *singular* if it coincides with p^\perp for some point p ; otherwise it is called *non-degenerate*, and then the point line geometry induced in it is a polar space.
- (3) A subspace of corank 2 will also be called a *geometric subhyperplane*.
- (4) A *collineation* of Π is a permutation of the point set \mathcal{P} that preserves the line set \mathcal{L} .
- (5) A *central collineation* (with *centre* $p \in \mathcal{P}$) is a collineation that fixes every point collinear to p .

- (6) An *axial collineation* (with axis $L \in \mathcal{L}$) is a collineation θ that stabilises each line that intersects L . Necessarily θ fixes each point collinear with all points of L .
- (7) Central collineations θ_i ($i = 1, 2$) with centres p_i ($i = 1, 2$) are *perpendicular* if p_1 and p_2 are collinear.
- (8) Axial collineations θ_i ($i = 1, 2$) with centres L_i ($i = 1, 2$) are *perpendicular* if either L_1 and L_2 intersect in a point and are not coplanar (we call these of *the first kind*), or L_1 and L_2 are contained in the same 3-space and do not intersect (we call these of *the second kind*).
- (9) Two central collineations, or two axial collineations, are *opposite* if their centres are opposite.
- (10) An *ideal Baer collineation* of polar space of rank 3 is a collineation that has at least one fixed line, induces a Baer collineation in every fixed plane, and fixes every plane through each fixed line.

We will always assume, unless explicitly stated otherwise, that polar spaces have thick lines. That is, every line contains at least 3 points. We call a polar space of rank n *non-thick* if every $(n - 2)$ -dimensional singular subspace is contained in exactly two maximal singular subspaces (of dimension $n - 1$), and thick otherwise. Thick polar spaces correspond to buildings of type B_n and C_n (the chambers of the building are maximal flags of singular subspaces in the polar space, and in this way the type $i + 1$ vertices of the building correspond to singular subspaces of dimension i for $0 \leq i \leq n - 1$). In the non-thick case every $(n - 2)$ -dimensional singular subspace is contained in exactly two maximal singular subspaces, and there are two types of maximal singular subspaces (members of the same type intersect each other in subspaces of even codimension). The so called “oriflamme complex” of such a polar space is then a thick building of type D_n .

We may regard an automorphism θ of a building Δ of type B_n, C_n or D_n as an automorphism, also denoted θ , of the associated polar space $\Pi = (\mathcal{P}, \mathcal{Q})$. Since neither dualities of B_2/C_2 buildings nor trialities of D_4 buildings play a role in this work, all such automorphisms will be collineations of the polar space Π (however, note that in the D_n case, the automorphism θ may interchange the type $n - 1$ and n vertices of the building).

All thick polar spaces, with the exception of one class in rank 3, are embeddable in projective space. More precisely, they arise as the isotropic geometry of non-degenerate quadratic forms, alternating forms, and Hermitian forms, or are a subspace thereof (see [20]); in the latter case the underlying field is non-commutative and has characteristic 2 and they are described by a pseudo-quadratic form, but we will not need this in this work, since, for example, we will prove results that exclude non-commutative fields, such as Proposition 3.15. The polar spaces related to quadratic forms are called *orthogonal polar spaces* and arise as non-degenerate quadrics in projective space.

The split spherical buildings of type D_n correspond to *hyperbolic quadrics* (with the standard equation $X_{-2n}X_{2n} + X_{-2n+1}X_{2n-1} + \dots + X_{-1}X_1 = 0$), and those of type B_n correspond to the *parabolic quadrics* (with standard equation $X_{-2n}X_{2n} + X_{-2n+1}X_{2n-1} + \dots + X_{-1}X_1 = X_0^2$). The split buildings of type C_n are related to a non-degenerate alternating form in standard form $X_{-2n}Y_{2n} - X_{2n}Y_{-2n} + X_{-2n+1}Y_{2n-1} - X_{2n-1}Y_{-2n+1} + \dots + X_{-1}Y_1 - X_1Y_{-1}$. Note that these arise as the fixed point geometry of symplectic polarities in projective space. It is well-known that, if the field \mathbb{K} is perfect of characteristic 2, then the polar spaces of type B_n and C_n are isomorphic; if the field is not perfect (but still has characteristic 2), then the polar space of type B_n is a polar subspace of the one of type C_n . For ease of formulation, we will call a symplectic polar space in characteristic distinct from 2 a *proper symplectic polar space*.

Polar spaces associated to a non-degenerate σ -Hermitian form, with σ an involution of the underlying field \mathbb{K} , are called Hermitian polar spaces. They are called *minimal Hermitian* if the dimension of the projective space is $2n - 1$, where n is the rank of the polar space. The standard equation of a minimal Hermitian polar space is $X_{-2n}^\sigma X_{2n} + X_{-2n}^\sigma X_{-2n} + X_{-2n+1}^\sigma X_{2n-1} + X_{2n-1}^\sigma X_{-2n+1} + \dots + X_{-1}^\sigma X_1 + X_1^\sigma X_{-1} = 0$.

In the language of polar spaces it is convenient to use “(projective) dimension as a singular subspace” rather than “type of a vertex of the building”, and therefore there is a shift of indexing. If θ is an automorphism of Π we say that θ is *i-domestic* if θ maps no singular subspace of dimension i to an opposite. Thus *i-domesticity* means domestic on type $i + 1$ vertices of the building. There is an exception in non-thick polar spaces, where there are two types of maximal singular subspaces. These both have projective dimension $n - 1$, and we will call them $(n - 1)'$ -spaces and $(n - 1)$ -spaces, and these correspond to the vertices of types $n - 1$ and n in the D_n building. The expressions *point-domestic* and *line-domestic*, etc, have the obvious meanings.

2 Class I collineations

Recall that the class I collineations of a polar space are those domestic collineations that map at least one point to an opposite point. Thus the opposition diagram of a class I collineation is $X_{n,i}^1$ for some $0 \leq i < n$ (where $X \in \{B, C, D\}$). Theorem 1 (proved below) shows that these collineations are characterised by large structured fixed point sets.

Lemma 2.1. *Let Π be a polar space of rank n and let $0 \leq i \leq n - 1$. Suppose that θ is i -domestic. Moreover,*

- (1) *if $i < n - 1$ suppose that θ is $(i + 1)$ -domestic, and*
- (2) *if $i > 0$ suppose that θ is not $(i - 1)$ -domestic.*

Then θ fixes pointwise a geometric subspace of corank i .

Proof. The statement in the case $i < n - 1$ is precisely [18, Theorem 6.1]. If $i = n - 1$ then we follow the ideas of the proof of [18, Theorem 6.1]. Only the first step of the proof in [18, Theorem 6.1] uses the fact that there exist singular subspaces of dimension $i + 1$. Since here $i = n - 1$ (and so no such singular subspaces exist), we provide a different argument for this step of the proof for this case.

Let U be a non-domestic submaximal singular subspace and let M be any maximal singular subspace containing U . By $(n - 1)$ -domesticity, M^θ intersects M in a point p . We now claim that $p^\theta = p$. Suppose for a contradiction that $p^\theta \neq p$ and choose a submaximal singular subspace U_0 contained in M , with $p \in U_0$ and $p^{\theta^{-1}} \notin U_0$. Then U_0^θ contains p^θ but not p . Let M_0 be any maximal singular subspace containing U_0 , but distinct from M . Then M_0 and M_0^θ have a point p_0 in common (by $(n - 1)$ -domesticity). Clearly, $p_0 \notin M^\theta$. But p_0 is collinear with all points of U_0^θ (since it belongs to M_0^θ) and also to p (as it belongs to M_0), which implies that p_0 is collinear to all points of M^θ , a contradiction. This proves the claim.

It follows that $U^\perp \cap (U^\theta)^\perp$ is fixed pointwise, and the remainder of the proof of [18, Theorem 6.1] now applies verbatim. □

We can now prove Theorem 1.

Proof of Theorem 1. Let θ be a nontrivial domestic collineation of a polar space Π with opposition diagram $X_{n,i}^1$ with $X \in \{B, C, D\}$, $n \geq 2$, and $1 \leq i \leq n$. Since we assume that Π is large the collineation θ is capped, and so since θ is domestic we have $i < n$. Thus θ is domestic on type $i + 1$ vertices of the associated building, that is, θ is i -domestic (recall the index shift for projective dimension). Then the hypothesis of Lemma 2.1 holds, and hence θ fixes pointwise a geometric subspace of corank i .

For the converse, we assume that θ maps at least one point to an opposite point, fixes a geometric subspace of corank i , and fixes no geometric subspace of any smaller corank. If θ maps no singular $(i - 1)$ -space to an opposite then θ has diagram $X_{n,j}^1$ for some $j < i$ (by the classification of admissible diagrams, and the fact that a point is mapped to opposite). Thus by the previous paragraph θ fixes a geometric subspace of corank $j < i$, which is a contradiction. It remains to show that θ does not map any singular i -space to an opposite. This is clear from the definition of subspaces of corank i : the fixed structure intersects every singular i -space, hence no such singular subspace is mapped to an opposite. □

The simplest situation occurs with opposition diagram $X_{n,1}^1$, since this is equivalent to a collineation pointwise fixing exactly a geometric hyperplane. It then depends on the structure of the possible geometric hyperplanes whether or not some general statements can be made about such collineations. In particular, if the polar space is split or non-embeddable, we can nail them all down. If the polar space is orthogonal (arises from a quadric) in characteristic distinct from 2, then the geometric hyperplane must be non-degenerate. In the next paragraphs, we give more precise details for these cases. We start with symplectic polar spaces, where we can completely classify the collineations with diagrams $C_{n,1}^1$ and $C_{n,2}^1$. Consider first the case $C_{n,1}^1$.

Theorem 2.2. *Every collineation θ of the symplectic polar space $\Delta = C_{n,1}(\mathbb{K})$ with polar opposition diagram (that is, $\text{Diag}(\theta) = C_{n,1}^1$) is a central elation.*

Proof. By [18, Theorem 5.1] the collineation θ fixes pointwise a geometric hyperplane H . If H spans the whole ambient $(2n - 1)$ -dimensional projective space, then θ is the identity (since every collineation of $C_{n,1}(\mathbb{K})$ is inherited from a collineation of the ambient projective space). Hence H is contained in a projective hyperplane of the ambient projective space, and hence coincides with the perp of some point p . Since θ fixes p^\perp pointwise, dually, θ fixes p subspace-wise, that is, θ fixes all subspaces through p . Hence θ is the central collineation with centre p (and axis p^\perp in the ambient projective space). \square

The opposition diagram $C_{n,2}^1$ is the *polar-copolar* diagram of type C_n (extending the terminology from [13]). In this case we have the following classification.

Theorem 2.3. *Let $n \geq 3$. A collineation θ of the symplectic polar space $\Delta = C_{n,1}(\mathbb{K})$ with $|\mathbb{K}| \geq 3$ has opposition diagram $\text{Diag}(\theta) = C_{n,2}^1$ if and only if θ is either*

- (1) *a product of two nontrivial perpendicular long root elations, or*
- (2) *a nontrivial member of the group generated by two opposite long root groups and not conjugate to a long root elation.*

In particular, if $\text{char}(\mathbb{K}) \neq 2$ then all nontrivial short root elations have opposition diagram $C_{n,2}^1$.

Proof. Suppose that θ has opposition diagram $C_{n,2}^1$. Since $|\mathbb{K}| \geq 3$ the polar space is large, and so by Theorem 1 the collineation θ pointwise fixes a subspace S of corank 2. We claim that in the ambient projective space $\text{PG}(2n - 1, \mathbb{K})$, S is a subspace, necessarily of dimension $2n - 3$. Indeed, since S is a subspace and is fixed pointwise, the projective subspace it generates in $\text{PG}(2n - 1, \mathbb{K})$ is fixed pointwise by θ and we may assume it coincides with S (otherwise we replace S by it). If $\dim S = 2n - 2$, then S is a hyperplane and θ is line-domestic (as every line contains a fixed point), a contradiction. Now assume for a contradiction that $\dim S \leq 2n - 4$. Since θ is not line-domestic, there exists a singular plane π of Δ intersecting S in just a point p . Let L be any line in π not through p . Then every singular plane through L intersects S , so $L^\perp \subseteq \langle L, S \rangle$. This yields $\dim S \in \{2n - 4, 2n - 5\}$. If $\dim S = 2n - 5$, then $L^\perp = \langle L, S \rangle$, and considering another line $M \subseteq \pi$ not through p , we obtain $M^\perp \subseteq \langle M, S \rangle = \langle L, S \rangle$ and hence comparing dimensions, we see $L^\perp = M^\perp$, a contradiction. This shows $\dim S = 2n - 4$, hence $\dim \langle L, S \rangle = 2n - 2$ and there is a unique point x on L such that $x^\perp = \langle L, S \rangle = \langle \pi, S \rangle$. Varying L in π appropriately, this implies $x^\perp = y^\perp$ for at least two points of π , a contradiction. Hence $\dim S = 2n - 3$.

So $S = p^\perp \cap q^\perp$, for two points p, q of Δ . Suppose first that p and q are not collinear. Then the matrix M_A of θ can be written as a block diagonal matrix with blocks the $(2n - 2) \times (2n - 2)$ -identity matrix and one 2×2 block A preserving a non-degenerate 2-dimensional symplectic form. Hence A has determinant 1. But then, A , and hence θ , is generated by the unipotent upper triangular and lower triangular matrices, hence by opposite long root elations.

Suppose now that p and q are collinear. We consider the symplectic form $x_1y_2 - x_2y_1 + x_3y_4 - x_4y_3 + \dots + x_{2n-1}y_{2n} - x_{2n}y_{2n-1}$ and we may assume without loss of generality that $p = (1, 0, 0, \dots, 0)$, $q = (0, 0, 1, 0, \dots, 0)$. Then one easily calculates that the matrix of θ is block diagonal, with all trivial blocks except the first 4×4 block, which equals

$$\begin{pmatrix} 1 & a & 0 & c \\ 0 & 1 & 0 & 0 \\ 0 & c & 1 & b \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

for some $a, b, c \in \mathbb{K}$. If we denote the corresponding collineation of Δ by $\theta(a, b, c)$, then we see that $\theta(a, b, c) = \theta(a, 0, 0)\theta(0, b, 0)\theta(0, 0, c)$, where the first two are long root elations and the third one is a short root elation. Also, one calculates that, if $a \neq 0$, then $\theta(a, c^2/a, c)$ is a long root elation with centre $(a, 0, c, 0, 0, \dots, 0)$, and $\theta(a, b, c) = \theta(a, c^2/a, c)\theta(0, b - c^2/a, 0)$ is the product of two perpendicular long root elations. Similarly if $b \neq 0$. Hence we may assume that $(a, b) = (0, 0)$. In this case $\theta = \theta(0, 0, c)$ is a short root elation. If $\text{char}(\mathbb{K}) = 2$, then θ is a long root elation in the corresponding building of type B_n and hence has opposition diagram $B_{n,1}^2 = C_{n,1}^2$ (by [13, Theorem 2.1]). Suppose now $\text{char}(\mathbb{K}) \neq 2$. Then $\theta(0, 0, c) = \theta(c/2, c/2, c/2)\theta(-c/2, -c/2, c/2)$, with $\theta(\pm d, \pm d, d)$ a central long root elation with centre $(1, 0, \pm 1, 0, \dots, 0)$ (using respective signs).

Thus we have shown that every collineation with diagram $C_{n;2}^1$ is either as in (1) or (2). The converse is a direct calculation (see the proof of Proposition 4.4, for example). \square

Quite similar to symplectic polar space is the result for thick non-embeddable polar spaces. We use $C_{3;1}^1$ for the opposition diagrams here since these polar spaces behave like symplectic ones (and their Dynkin type is morally C_3 in view of the underlying root system in the algebraic group sense, made apparent by the commutation relations of their root groups).

Theorem 2.4. *Every collineation θ of a thick non-embeddable polar space Δ with polar opposition diagram (that is, $\text{Diag}(\theta) = C_{3;1}^1$) is a central elation.*

Proof. According to [5], the only geometric hyperplanes are the singular ones, that is, all points collinear with a given fixed point p . Let x be an arbitrary point not collinear to p . Then $x^\perp \cap p^\perp$ is pointwise fixed, and so x^θ is contained in $\{p, x\}^{\perp\perp}$. But then θ coincides with the unique central elation with centre p mapping x to x^θ . For an explicit expression of such central elation (showing it really exists), see [7, Proposition 4.13]. \square

A similar argument shows:

Proposition 2.5. *Every collineation θ of an orthogonal polar space Δ over a field of characteristic not 2 with opposition diagram $\text{Diag}(\theta) = B_{n;1}^1$ pointwise fixes a non-degenerate geometric hyperplane. Also, θ is an involution.*

Proof. Since the characteristic of the field is not 2 the polar space is large, and so by Theorem 1 the collineation θ fixes a geometric hyperplane H pointwise. Suppose that H is singular, that is, $H = p^\perp$ for some point p of Δ . Since θ commutes with the non-degenerate polarity ρ arising from the symmetric bilinear form that defines Δ , and since θ fixes all points in $p^\rho = \langle p^\perp \rangle$ (generation in the ambient projective space), it has to stabilise all hyperplanes through p , hence also all lines through p . But each such line outside H intersects Δ in one further point x , hence $x^\theta = x$, for each point outside H . This shows the first assertion.

A similar argument shows that θ is a homology in the ambient projective space with axis $\langle H \rangle$ and centre $c := \langle H \rangle^\rho$. Since lines through c intersect Δ in at most two points, θ is an involution. \square

Next we consider split polar spaces of type B_n , $n \geq 3$, and D_n , $n \geq 4$. For type B_n , there is not anything we can say on top of Proposition 2.5. However, for type D_n , we can be slightly more specific. Indeed, the opposition diagram $D_{n;1}^1$ characterises the unique involution pointwise fixing a polar subspace of split type B_{n-1} , and it fixes no chamber.

Theorem 2.6. *Every collineation θ of a polar space Δ of type D_n , $n \geq 4$, with opposition diagram $D_{n;1}^1$ is an involution pointwise fixing a polar subspace of type B_{n-1} . It follows that θ is automatically type-rotating.*

Proof. This follows from the fact that there are no non-degenerate quadrics in $\text{PG}(2n - 2, \mathbb{K})$ of Witt index n (since every pair of maximal subspaces would have to intersect nontrivially), and there is a unique quadric in $\text{PG}(2n + 1, \mathbb{K})$ with Witt index n , and it is a parabolic one. \square

Remark 2.7. Note that if $\text{char}(\mathbb{K}) = 2$ then $B_{n;1}(\mathbb{K})$ is a subspace of $C_{n;1}(\mathbb{K})$ (with equality if \mathbb{K} is perfect). In the non-perfect case there exist line-domestic collineations distinct from central collineations, whereas in the perfect case, we can rely on Theorem 2.2 to deduce that we always have a central collineation. A counterexample for the non-perfect case is given by the collineations of $\text{PG}(2n - 1, \mathbb{K})$ with action

$$(x_{-n}, x_{-n+1}, \dots, x_{-1}, x_1, \dots, x_n) \mapsto (x_{-n}, x_{-n+1}, \dots, x_{-2}, ax_1, a^{-1}x_{-1}, x_2, \dots, x_n),$$

with a a non-square, with respect to the (standard) form $x_{-n}x_n + \dots + x_{-1}x_1 \in \mathbb{K}^2$. (If this was a central collineation, then the point with coordinates all 0 except $(x_{-1}, x_1) = (a, 1)$ would be the centre, but this point does not belong to the polar space.) In fact one can choose the coordinates so that the fixed subspace is the above one, that is, has equation $X_{-1} = aX_1$, and then a generic collineation pointwise fixing that geometric hyperplane is given by

$$(x_{-n}, \dots, x_{-1}, x_1, \dots, x_n) \mapsto (x_{-n}, \dots, (ab + 1)x_{-1} + a^2bx_1, bx_{-1} + (ab + 1)x_1, \dots, x_n),$$

with $ab^2 + b \in \mathbb{K}^2$.

3 Point-domestic collineations of a polar space

In this section we prove Theorems 3–9, dealing with point-domestic collineations. We begin with preliminary observations, before dividing the analysis into collineations of class II and class III. Note that by [12, Theorem 1] all point-domestic collineations of a polar space are capped.

3.1 Preliminary observations

Here is the first basic observation.

Proposition 3.1. *Let Δ be a polar space of rank at least 2. Let θ be an automorphism of Δ and suppose that θ is point-domestic. Then for every point p with $p^\theta \neq p$, the line $\langle p, p^\theta \rangle$ is fixed under θ .*

Proof. Suppose for a contradiction that there exists a point p with $p^\theta \neq p$ and $\langle p, p^\theta \rangle$ not fixed under θ . Pick $x \in \langle p, p^\theta \rangle \setminus \{p, p^\theta\}$. Then $x^\theta \in \langle p^\theta, p^{\theta^2} \rangle \setminus \{p^\theta, p^{\theta^2}\}$. By point-domesticity, x is collinear to all points of $\langle p^\theta, p^{\theta^2} \rangle$, and so $p, p^\theta, p^{\theta^2}$ are contained in a plane π of Δ , which they span. Let α be a plane containing $\langle p, p^\theta \rangle$ and so that α and π are not contained in a common singular subspace of Δ . Pick $q \in \alpha \setminus \pi$ and $r \in \langle q, p^\theta \rangle \setminus \{q, p^\theta\}$ and note that q is not collinear to p^{θ^2} . We then have $r \perp r^\theta$ and $p^\theta \perp r^\theta$ (as $p \perp r$, both are in α). So r^θ is collinear to all points of $\langle r, p^\theta \rangle$. Likewise, q^θ is collinear to all points of $\langle q, p^\theta \rangle = \langle r, p^\theta \rangle$. Hence all points of $\langle q^\theta, r^\theta \rangle$ (in particular, p^{θ^2}) are collinear to all points of $\langle q, p^\theta \rangle$ (in particular, q). This contradiction concludes the proof. \square

We note an immediate consequence of Proposition 3.1.

Corollary 3.2. *If θ is a point-domestic collineation of some polar space Δ , then every member of $\langle \theta \rangle$ is point-domestic.*

Proof. Let x be any point of Δ . If $x = x^\theta$ then x is fixed by every member of $\langle \theta \rangle$; if $x \neq x^\theta$ then θ stabilises $\langle x, x^\theta \rangle$ and so does every member of $\langle \theta \rangle$. \square

The above proposition prompts the following question: Which collineations of a projective space have the property that each line joining a non-fixed point and its image is fixed? The answer is in the following proposition.

Proposition 3.3. *Let Σ be a finite-dimensional projective space with dimension at least 2 and θ a collineation with the property that each point of Σ is contained in at least one fixed line (or, equivalently, the line $\langle x, x^\theta \rangle$ is fixed for every point $x \neq x^\theta$). Then exactly one of the following holds.*

- (i) *There are no fixed points and the set of fixed lines forms a line spread of Σ . The projective dimension of Σ is odd, say $2n + 1$, and the fixed subspaces form a projective space of dimension n .*
- (ii) *The set of fixed points forms a Baer subspace \mathfrak{B} of Σ .*
- (iii) *The set of fixed points is the union of two complementary subspaces U, V of Σ and θ is a (U, V) -homology, i.e., θ fixes all subspaces in $U \cup V$ and all subspaces containing one of U, V .*
- (iv) *The set of fixed points is a subspace U of dimension ℓ , with $2\ell + 1 \geq \dim \Sigma$, and the intersection of all fixed hyperplanes is a subspace $V \subseteq U$ of dimension $\dim \Sigma - \ell - 1$. Then θ is a (U, V) -elation, i.e., θ fixes every subspace of U and every subspace containing V .*

Proof. If θ has no fixed points, then clearly (i) holds. So from now on we may assume that there is at least one fixed point. The proof is divided into the following steps.

Step 1. Suppose first $\dim \Sigma = 2$. If all lines are fixed, then θ is the identity and (iv) holds with $\ell = 2$. Hence we may assume that some line L is not fixed. We claim that $x = L \cap L^\theta$ is fixed. Indeed, if not then $x \neq x^\theta \in L^\theta$. Since x^θ also belongs to every line through x that is fixed under θ , we obtain the contradiction that L^θ is fixed. The claim is proved. We obtain a second fixed point y by intersecting the fixed lines through two distinct points of $L \setminus \{x\}$.

Suppose first that no point off xy is fixed. Let $z \in xy$ be arbitrary and choose a line $M \ni z$, with $M \neq L$. If $M = M^\theta$, then $z = M \cap xy$ is fixed; if $M \neq M^\theta$, then $M \cap M^\theta$ is fixed and it is necessarily z as otherwise it lies off xy , contrary to our assumption. We have proved that each point of xy is fixed and therefore θ is axial and hence central, so (iv) holds with $\ell = 1$.

Suppose now a point z off xy is fixed, but no quadrangle is fixed. Let K be a line disjoint from $\{x, y, z\}$. If K is fixed, then the intersection of the lines joining x and $yz \cap K$, and y and $xz \cap K$, respectively, is fixed and forms a quadrangle with x, y, z , a contradiction. Hence $K \neq K^\theta$ and since $K \cap K^\theta$ is fixed, it lies on one of the lines xy, yx, xz , say xy . Using a similar argument as above, one shows that for every line M disjoint from $\{x, y, z\}$, the image M^θ intersects M on xy . Hence we conclude that xy is fixed pointwise and (iii) holds.

Finally suppose θ fixes some quadrangle. Then the fixed point structure is a subplane π . Every point is contained in at least one line of π (by the assumption on θ), and every line N is either fixed, and hence contains many fixed points, or is not fixed and contains the fixed point $N \cap N^\theta$. So (ii) holds.

Step 2. We note that, if some subspace U is fixed, then the restriction $\theta|_U$ of θ to U satisfies the assumptions of the proposition for \mathcal{E} replaced by U .

Step 3. Suppose $\dim \mathcal{E} \geq 3$ and the fixed point set P does not generate \mathcal{E} . We first claim that P is a subspace. Indeed, let $x, y \in P$ and let $z \in \mathcal{E} \setminus \langle P \rangle$. Then the line $L = \langle z, z^\theta \rangle$ is fixed, and hence the plane $\alpha = \langle L, x \rangle$ is fixed. Hence steps 1 and 2, and the choice of z outside $\langle P \rangle$ implies that $\alpha \cap P$ is a full line M through x . We may assume $y \notin M$, as otherwise the claim follows. Then $u = L \cap M$ is fixed and so is the plane $\langle y, z, u \rangle$. Again steps 1 and 2 and the choice of z imply that all points of the line $\langle y, u \rangle$ are fixed. Then the fixed plane $\langle x, y, u \rangle$ contains two lines all of whose points are fixed, and the claim follows from steps 1 and 2.

Note that the previous argument also shows that, whenever $z \in \mathcal{E} \setminus P$, then $\langle z, z^\theta \rangle$ contains a unique fixed point. Let Q be the set of all points $q \in P$ such that some line through q not in P is fixed. We claim that Q is a subspace. Indeed, let $q, r \in Q$. Then the definition of Q implies that there exist points $x, y \in \mathcal{E} \setminus P$ with $q \in \langle x, x^\theta \rangle$ and $r \in \langle y, y^\theta \rangle$. Clearly $X = \langle x, y, q, r \rangle$ is 3-dimensional, and $X \cap P = \langle q, r \rangle$. Hence the line $L = \langle x, y \rangle$ does not intersect P . This implies that $L \cap L^\theta = \emptyset$. Let $s \in \langle q, r \rangle$ arbitrary. Let $z \in L$ and $z' \in L^\theta$ be such that $s \in \langle z, z' \rangle$. Then z' is the unique point on L^θ such that $\langle z, z' \rangle$ intersects P . We deduce that $z' = z^\theta$ and so $s \in Q$. The claim is proved.

Next we claim that Q is the intersection of all fixed hyperplanes. Indeed, first let H be a hyperplane containing Q . Pick $z \in H \setminus P$. Then $\langle z, z^\theta \rangle$ intersects $Q \subseteq H$, hence $z^\theta \in H$. Consequently H is fixed. Now let H be a hyperplane not containing Q . Select $x \in Q \setminus H$. Let $z \in \mathcal{E} \setminus P$ be such that $x \in \langle z, z^\theta \rangle$. Since $x \notin H$, we may assume $z \in H$ (a hyperplane intersects every line nontrivially). But then $z^\theta \notin H$ and so $H \neq H^\theta$. The claim is proved.

Now notice that in the dual of \mathcal{E} , the roles of P and Q are interchanged. Hence, up to duality, we may assume $k = \dim Q \leq \frac{1}{2}(n - 2)$ (if $k = \frac{1}{2}(n - 1) = \dim P$, then (iv) holds). Let W be a subspace complementary to Q . Then $W \cap W^\theta$ has dimension at least $n - 2k - 2 \geq 0$. Hence $W \cap W^\theta$ is nonempty. Let $w \in W \cap W^\theta$ be arbitrary. Then $\langle Q, w \rangle$ is fixed and so is $w = \langle Q, w \rangle \cap W = \langle Q, w \rangle \cap W^\theta$. Hence $W \cap W^\theta \subseteq P$ and so $\ell = \dim P \geq n - k - 1$. If $\ell \geq n - k$, then the dual argument could be applied and would yield $n - k - 1 \geq n - k$, a contradiction. Hence $k + \ell = n - 1$ and (iv) holds.

Step 4. Suppose $\dim \mathcal{E} \geq 3$, the fixed point set \mathfrak{B} generates \mathcal{E} and has the structure of a thick projective space. Then by steps 1 and 2, every fixed plane α intersects \mathfrak{B} in a Baer subplane of α and (ii) holds.

Step 5. Suppose $\dim \mathcal{E} \geq 3$, the fixed point set P generates \mathcal{E} and has the structure of a non-thick generalised projective space. We first claim that, if L is a thick line of P , then all points of \mathcal{E} incident with L belong to P . Indeed, suppose not and let $\{x, y\}$ be a thin line of P . The plane $\pi = \langle L, x \rangle$ is fixed by θ and steps 1 and 2 imply that $\pi \cap P$ is a Baer subplane of π . Let B be a line of π through x intersecting P in at least three points. Then B is a proper subline of its carrier in \mathcal{E} . The plane $\langle B, y \rangle$ is fixed and steps 1 and 2 imply that $\langle x, y \rangle$ contains at least three points of P , a contradiction. The claim is proved.

Now let $\{x, y\}$ again be a thin line (which exists by assumption). Let P_x and P_y be the sets of points of P lying on a thick line together with x and y , respectively. We claim that $\{P_x, P_y\}$ is a partition of P . Indeed, if $u \in P_x \cap P_y$, then the plane $\langle x, y, u \rangle$ contains two lines entirely consisting of fixed points, hence steps 1 and 2 imply that all points of that plane are fixed; in particular all points of $\langle x, y \rangle$, contradicting our assumption on $\langle x, y \rangle$. Also, suppose there exists $v \in P \setminus (P_x \cup P_y)$. Then the plane $\langle x, y, v \rangle$ is fixed and $\langle x, y, v \rangle \cap P = \{x, y, v\}$.

This contradicts steps 1 and 2 once again. The claim follows. Consider now two arbitrary points z, w of P_x with $z \neq w \neq x \neq z$. Then the plane $\langle x, z, w \rangle$ contains two lines all of whose points are in P . Steps 1 and 2 imply that also all points of $\langle z, w \rangle$ are fixed. Hence P_x , and similarly P_y , is a subspace of Σ . Now (iii) holds. \square

Examples of each kind are easy to find. One remark concerning Case (ii): if the underlying structure is a field (the projective space is Pappian then), then such an automorphism θ is induced by the identity matrix and a companion field automorphism pointwise fixing a subfield \mathbb{F} of a field \mathbb{K} such that \mathbb{K} is 2-dimensional over \mathbb{F} . Hence θ is an involution. But if the underlying division ring is not commutative, then examples exist where θ is not an involution (note that this corrects [17, Theorem 7.2] where the statement “ideal Baer involution” should be replaced with “ideal Baer collineation”). Consider for instance a field \mathbb{F} with a quadratic Galois extension \mathbb{K} , and denote the Galois involution by $x \mapsto \bar{x}$. Then define the division ring $\mathbb{K}_{\mathbb{F}}((t))$ of Laurent series $f(t)$ over \mathbb{K} with multiplication induced by $xt = t\bar{x}$, for all $x \in \mathbb{K}$. Let $a \in \mathbb{K}$ be such that $a\bar{a} = 1$, but $a \notin \{1, -1\}$ (this always exists). Then the mapping $\theta : f(t) \mapsto f(at)$ defines an automorphism pointwise fixing the field $\mathbb{K}((t^2))$ of Laurent series $f(t^2)$ over \mathbb{K} (and $\mathbb{K}_{\mathbb{F}}((t))$ is 2-dimensional over $\mathbb{K}((t^2))$ and the induced automorphism on any projective space of dimension at least 2 over $\mathbb{K}_{\mathbb{F}}((t))$ satisfies (ii)). However, the order of θ is the multiplicative order of a in \mathbb{K} , which is distinct from 2 as $a \neq -1$.

This has now the following interesting consequence.

Corollary 3.4. *A point-domestic collineation θ of a polar space $\Delta = (X, \Omega)$ of odd rank r fixes a chamber.*

Proof. We first claim that θ fixes a maximal singular subspace, i.e., a singular subspace of dimension $r - 1$. Note that [18, Theorem 4.2] proves the claim for polar spaces that are not symplectic, however here we provide an independent proof that includes the symplectic case.

Suppose θ does not fix any maximal singular subspace. Let $d \leq r - 2$ be the maximal dimension of a fixed singular subspace and let U be fixed under θ with $\dim U = d$. Then $d \geq 1$ by Proposition 3.1. Since $d < r - 1$, there exists a point x collinear to all points of U . If x is fixed, then the singular subspace generated by U and x is fixed under θ and has dimension $d + 1$, a contradiction to the maximality of d . Hence $x \neq x^\theta$ and θ fixes the line xx^θ . Since U is fixed, x^θ is collinear to all points of U and U and xx^θ generate a subspace U' of dimension $d + 1$ or $d + 2$, fixed under θ , again a contradiction to the maximality of d . We conclude that the claim holds.

Now let U be a maximal singular subspace fixed by θ . By Proposition 3.3, we always fix a chamber of U , and hence a chamber of Δ , except in Case (i). But that case does not occur here since $\dim U = r - 1$ is even. \square

Lemma 3.5. *A point-domestic collineation θ of a polar space Δ of rank at least 3 with non-Pappian planes (hence defined over a non-commutative division ring) fixes a chamber.*

Proof. If Δ has rank 3, then this follows from Corollary 3.4. Suppose now that the rank of Δ is at least 4. Then Δ is thick. If the opposition diagram of θ is $B_{n,1}^2$, then Proposition 3.11, proved independently, implies that θ is an axial collineation, hence it fixes any chamber containing the axis. Consequently, by the classification of opposition diagrams, θ has opposition diagram $B_{n,i}^2$ with $i > 1$. But that means that there is some non-domestic 3-space U . The duality θ_U , however, must be domestic, implying it is a symplectic polarity, contradicting the fact that U is non-Pappian. \square

Finally, we observe that we automatically have a class II collineation if it is $(n - 1)$ -domestic, as follows immediately from Lemma 2.1.

Corollary 3.6. *Let Δ be a thick building of type X_n with $X \in \{B, C, D\}$ and let $\Pi = (\mathcal{P}, \Omega)$ be the associated polar space. If a collineation θ has opposition diagram $X_{n,i}^2$ with $2i < n$, then θ pointwise fixes a geometric subspace of corank at most $n - 1$.*

3.2 Class II automorphisms

If θ is a class II automorphism of a polar space of rank n , then $\text{Diag}(\theta) = X_{n,i}^2$ for some i with $0 \leq i \leq n/2$ (where $X \in \{B, C, D\}$).

3.2.1 The opposition diagrams $X_{n,1}^2$

The point-domestic collineations with opposition diagram $X_{n,1}^2$ (with $X \in \{B, C, D\}$) can be classified. This is the content of Theorem 3, which we prove in this subsection. We begin with the symplectic case, that is, Theorem 3(1).

Theorem 3.7. *Let $n \geq 2$. A collineation θ of the symplectic polar space $\Delta := C_{n,1}(\mathbb{K})$, assumed to have at least one fixed point if $n = 2$, having copolar opposition diagram (that is, $C_{n,1}^2$) is unique. It is a involutive homology if $\text{char}(\mathbb{K}) \neq 2$, and if $\text{char}(\mathbb{K}) = 2$ then θ is an axial elation.*

Proof. First assume $n \geq 3$. Then θ is j -domestic, for all $j \geq 3, j \leq n$. So it follows from [18, Theorem 6.1] that the fixed point structure of θ is a subspace of corank 2. Hence the same cases as in the proof of Theorem 2.3 apply. Clearly, with the notation of that proof, the case where p and q are non-collinear only occurs if all points of the hyperbolic line $\langle p, q \rangle$ are fixed. This gives rise to a unique involutive homology if $\text{char}(\mathbb{K}) \neq 2$ and to the identity if $\text{char}(\mathbb{K}) = 2$. If we have a collineation $\theta(a, b, c), a, b, c \in \mathbb{K}$ then one easily calculates that $\theta(a, b, c)$ maps the point $(x_i)_{1 \leq i \leq 2n}$ to a collinear one if and only if

$$ax_2^2 + 2cx_2x_4 + bx_4^2 = 0,$$

which implies the theorem.

Next assume that $n = 2$. Then we have a point-domestic symplectic generalised quadrangle. It follows from [19] that θ either linewise fixes a spread, or pointwise fixes an ideal subquadrangle, or is an axial elation. Since it is assumed that θ fixes at least one point, it is easily seen that the first case cannot occur (it leads to the identity).

Suppose now that θ pointwise fixes an ideal subquadrangle. Taking the standard symplectic form $x_0y_1 - x_1y_0 + x_2y_3 - x_3y_2$, we may assume that the ideal subquadrangle contains all lines meeting both hyperbolic lines $(*, *, 0, 0)$ and $(0, 0, *, *)$, with self-explaining notation. Then θ has the form $(x_0, x_1, x_2, x_3) \mapsto (x_0, x_1, ax_2, ax_3)$, for some $a \in \mathbb{K}$. This preserves the given symplectic form if and only if $a^2 = 1$, showing that we have an involutive homology if $\text{char}(\mathbb{K}) \neq 2$, and the identity if $\text{char}(\mathbb{K}) = 2$. Since in characteristic $\neq 2$ a symplectic quadrangle does not admit axial elations, the theorem is proved. \square

Next we consider all other polar spaces. Rank 3 will be an exception, but the domestic collineations in this case are already classified in [18, Section 6]. This includes the more intricate non-embeddable polar spaces (which do not admit axial elations).

In order to prove Theorem 3(2), we need some preparations. We start with gathering some properties of geometric hyperplanes and subhyperplanes in the following proposition. The *radical* of a subspace S is the set of points of S collinear to all points of S . It is always a singular subspace.

Proposition 3.8. *Let $\Delta = (X, \Omega)$ be a non-degenerate polar space of rank at least 3. Then the following properties hold.*

- (i) *Each geometric hyperplane and subhyperplane is a possibly degenerate polar space (and every point of the radical is called a deep point).*
- (ii) *The radical of a geometric hyperplane or subhyperplane is at most 0-dimensional, or 1-dimensional, respectively.*
- (iii) *Let S be a geometric subhyperplane and let $p \in X$ be a point off S . Set $S' = \{x \in X \mid x \in L, p \in L, S \cap L \neq \emptyset\}$. If $S \subseteq p^\perp$, then $S' = p^\perp$ is a geometric hyperplane. If some point of S is not collinear to p , then S' is a geometric subhyperplane.*
- (iv) *Every plane containing a deep point of a geometric subhyperplane contains a line of that subhyperplane.*

Proof. (i) This follows immediately from the definition of subspace.

- (ii) Suppose that the radical of a geometric hyperplane H contains a line L . Then a line L' opposite L has no point in common with H , contradicting the definition of a geometric hyperplane. The other statement is proved in a similar fashion.

(iii) Suppose first that $S \subseteq p^\perp$. Assume for a contradiction that some line L through p does not intersect S . Select any plane π containing L . Since S is a geometric subhyperplane, it intersects π in a point x (it cannot intersect in a line since this would contradict the fact that L does not meet S). Let K be a line in π not containing x , and let α be a plane through K not collinear to p . Then α and S should have a point y in common, and by construction $y \notin K$. Then $y \perp p$, contradicting the choice of α . This shows that $p^\perp = S'$. Clearly, this is a geometric hyperplane.

Now suppose that some point of S is not collinear to p . Assume for a contradiction that some plane β is disjoint from S' . Let β_p be a plane through p intersecting β in a line M . Then β_p contains some point s of S ; hence the line ps is contained in S' . Consequently the point $ps \cap M$ is contained in S' . So β contains a point of S' after all.

(iv) Suppose for a contradiction that some plane α intersects a geometric subhyperplane S (only) in a deep point x . Let α' be a plane through a line K of α not through x such that α' is not collinear to x . Then α' has some point y in common with S and xy belongs to S , hence $xy \cap K$ belongs to S , which is a contradiction.

The proposition is proved. □

Next, we need some more properties of subspaces (of certain corank).

Proposition 3.9. *Let S be a subspace of a polar space and assume that S is not a singular subspace (but it can be a degenerate polar space).*

- (i) *If some singular subspace D of dimension d is a geometric hyperplane of S , then it contains the radical and maximal singular subspaces of S have dimension either d or $d + 1$.*
- (ii) *If a set D of points of S is collinear to its complement $S \setminus D$, then either it is contained in the radical or the complement of D is contained in the radical.*
- (iii) *The complement in S of a proper geometric hyperplane of S can never be contained in the radical.*

Proof. For $x \in S$ we briefly write x^\perp for $x^\perp \cap S$.

- (i) Suppose first that there exists a point x in the radical R of S , which is not contained in D . Let y, z be two non-collinear points of S . The lines xy and xz both contain some point of D , which, by assumption, are collinear, contradicting the choice of y and z . So D contains the radical of S . Then D/R is a singular subspace which is a hyperplane in the residue of R , which is either a (non-degenerate) polar space (a contradiction since for every subspace there exists a disjoint line) or just a set of points. Then D/R is either empty or a single point, which proves (i).
- (ii) Suppose D is not contained in R and let $x \in D \setminus R$. Then x^\perp is a (proper) geometric hyperplane of S and contains $S \setminus D$ by assumption. Suppose $x^\perp \setminus (D \cup R) \neq \emptyset$ and let $y \in x^\perp \setminus (D \cup R)$. Select $z \in S \setminus (x^\perp \cup y^\perp)$. Then $z \in D$ as $z \notin x^\perp$. But then $y \in D$ as $y \notin z^\perp$, a contradiction.
- (iii) Suppose that H is a proper geometric hyperplane of S containing the complement of the radical R of S . Let $x \in R \setminus H$ and $y \in H \setminus R$. Then every point $z \in \langle x, y \rangle \setminus \{x, y\}$ belongs to $S \setminus (R \cup H) = \emptyset$, which is a contradiction.

The proposition is proved. □

Finally we need to recognise symplectic polar spaces, which we already treated in Theorem 3.7.

Proposition 3.10. *A polar space Δ of rank at least 3 is symplectic if and only if for some pair of noncollinear points $\{x, y\}$ (and then for all such pairs) every point z is contained in a line of the polar space containing a point of $\{x, y\}^\perp$ and a point of $\{x, y\}^{\perp\perp}$.*

Proof. The “only if” part is clear. We now show the “if” part.

Replacing Δ with the intersection of the perps of two appropriate opposite singular subspaces of codimension 2, we may assume that Δ has rank 2 (but is classical, implying that its automorphism group acts transitively on opposite pairs of points, and hence we may assume the given condition holds for each pair of non-collinear points). We show that each point is projective i.e., for each point x and every pair $\{y, z\}$ of points opposite x , the sets $\{x, y\}^\perp$ and $\{x, z\}^\perp$ either have exactly one point in common, or coincide.

Indeed, we may assume that $z \notin \{x, y\}^{\perp\perp}$ (otherwise $\{x, y\}^{\perp} = \{x, z\}^{\perp}$). Now, the given condition implies that z is on a line L containing $y' \in \{x, y\}^{\perp\perp}$ and $z' \in \{x, y\}^{\perp}$. Then $z' \in \{x, y, z\}^{\perp}$. If also $z' \neq z'' \in \{x, y, z\}^{\perp}$, then $z \in \{z', z''\}^{\perp} \supseteq \{x, y\}^{\perp\perp}$. Hence $y' \in \{z', z''\}^{\perp}$, implying the whole line L is collinear with z'' , hence $z'' \in L$ and consequently $z' = z''$.

The result now follows from [15]. □

We now prove Theorem 3(2).

Proposition 3.11. *Let $\Delta = (X, \Omega)$ be a polar space of rank at least 3, not of symplectic type, and let θ be an automorphism of Δ with opposition diagram $B_{n;1}^2$ or $D_{n;1}^2$. Then θ is an axial elation (and so Δ is an orthogonal polar space), or the rank is 3 and θ is an ideal Baer collineation.*

Proof. We first show that the fixed point structure of θ is precisely a subspace of corank 2, whenever θ is not an ideal Baer collineation in rank 3.

Theorems 6.1 and Theorem 7.2 of [18] already imply that θ pointwise fixes a subspace S of corank 2. Note that S is not a singular subspace as otherwise we can find a plane disjoint from S . We show that θ does not fix any point outside S . Indeed, suppose for a contradiction that θ fixes some point $x \in X \setminus S$.

We first claim that θ fixes pointwise every line incident with x and with a point of S . Indeed, we distinguish between two situations.

Suppose first that not all points of S are collinear to x . Then $x^{\perp} \cap S$ is a geometric hyperplane H of S . Suppose for a contradiction that all points of $S \setminus H$ are collinear to all points of H . Then by Proposition 3.9(ii), (iii), H is contained in the radical of S . Hence the union of all lines through x intersecting S is a singular subspace, but also a geometric subhyperplane by Proposition 3.8(iii), clearly a contradiction (an opposite subspace is disjoint). We conclude that there exist points $a \in H$ and $b \in S \setminus H$ which are not collinear. Hence the projection of b onto the line ax is also fixed. Consider any line L in S through a ; then the restriction of θ to the singular plane spanned by L and x is the identity as it fixes a line pointwise and two additional points. So θ pointwise fixes all lines through x intersecting S in a point collinear to a . By connectivity of H , θ fixes all lines through x intersecting S pointwise.

Now suppose that all points of S are collinear to x . Assume first that there is a point $y \neq x$ with $y \perp S$. Note that $\{x, y\}^{\perp} = S$ in this case (this follows from Proposition 3.8(iii)). By Proposition 3.10 we can select a point z not contained in a line joining a point of S and $\{x, y\}^{\perp\perp}$. Note that z is not fixed under θ as, by the choice of z , we can find a point w of $x^{\perp} \setminus S$ collinear to z , which must be fixed by the one-or-all axiom as otherwise ww^{θ} contains x . Then the line zz^{θ} intersects x^{\perp} in some point u , which is fixed by θ , since every line through x is fixed and since zz^{θ} is fixed by Proposition 3.1. If $u \notin S$, then a similar argument as in the previous paragraph implies that θ fixes x^{\perp} pointwise, implying line-domesticity, a contradiction. Hence we may assume that $u \in S$. Since S is a section of x^{\perp} , it is a non-degenerate polar space, so there is a point $v \in S \setminus u^{\perp}$. The projection x' of v onto zz^{θ} is a fixed point of θ and S is not contained in x'^{\perp} . Hence, by the first part of the proof, the line zz^{θ} is fixed pointwise. This contradiction implies that z is fixed after all, and hence also the point w above, and hence, as before, this implies that x^{\perp} is fixed pointwise. This contradiction shows the claim.

The previous proof clearly showed that S is not contained in x^{\perp} , as otherwise x^{\perp} is fixed pointwise and so θ is line-domestic. Let H_x be the union of all lines through x intersecting S . Then by Proposition 3.8(iii) H_x is a geometric subhyperplane of Δ .

Finally we claim that every line K contains a fixed point. Indeed, let K be any line and let β be any plane containing L . Since S is a subspace of corank 2, β contains a point $s \in S$, and since H_x is a subspace of corank 2, β contains a point $h \in H_x$. If $s \neq h$, then interchanging the roles of h and x yields a pointwise fixed line in β (namely, hs), and hence a fixed point on K . So we may assume that $s = h \in H$. Suppose first that there exists a point $r \in S$ not collinear to s . Then by the first part of the proof of the present claim, the plane containing r and the projection of r onto β contains a pointwise fixed line, which intersects K , proving the claim. So we may assume that s is collinear with all points of S . Then s is a deep point of S and by Proposition 3.8(iv) every plane through s has a line in common with S , again yielding a fixed point on K .

Hence θ is line-domestic, contradicting our hypothesis. This shows that indeed, the set of fixed points for θ is the geometric subhyperplane S . Let x be a point off S . We claim that the line xx^{θ} intersects S . Indeed, suppose

not. Then $x^\perp \cap S = (x^\theta)^\perp \cap S$. Since the line xx^θ does not intersect S , Proposition 3.8(iii) implies that S is not contained in x^\perp . But then the projection from a point of $S \setminus x^\perp$ onto the line xx^θ is a fixed point outside S , a contradiction. The claim follows.

Set $xx^\theta \cap S = c$. If some point z opposite c existed in S , then the projection of z onto xx^θ is a point off S that would be fixed, a contradiction. Hence c is a deep point for S . We can now choose $y \in X \setminus S$ opposite c and obtain a second deep point $c' = yy^\theta \cap S$ of S . Hence cc' is the radical of S (since the radical cannot be larger than a line by Proposition 3.8(ii)) and every point z which is not fixed satisfies $zz^\theta \cap cc' \neq \emptyset$. So θ is an axial elation with axis cc' . \square

3.2.2 Split polar spaces

Our attention now goes to the split case. Indeed, we can classify all point-domestic collineations of split polar spaces which are also domestic in the maximal singular subspaces. First a lemma and some notation: For three pairwise disjoint lines L, M, N of $\text{PG}(3, \mathbb{K})$, denote by $\text{proj}_M^L(N)$ the projection of N with centre L onto M , that is, $\text{proj}_M^L(N)$ takes a point $x \in N$ onto the intersection $\langle L, x \rangle \cap M$.

Lemma 3.12. *Let L_1, L_2 and L_3 be three pairwise disjoint lines in $\text{PG}(3, \mathbb{K})$ and let M be a transversal, that is, M is a line which intersects L_i in a point x_i for all $i \in \{1, 2, 3\}$. Let L be a line through x_1 in the plane $\langle L_1, M \rangle$ with $L_1 \neq L \neq M$, and, for $|\mathbb{K}| > 2$, let L' be a line intersecting both L_1 and M , but disjoint from the triple $\{x_1, x_2, x_3\}$. Then $\text{proj}_{L_2}^{L_1}(L_3) \text{proj}_{L_3}^L(L_2)$ has a unique fixed point, and $\text{proj}_{L_2}^{L_1}(L_3) \text{proj}_{L_3}^{L'}(L_2)$ has exactly two fixed points.*

Proof. This follows easily as $\{L_1, L_2, L_3\}$ and $\{L, L_2, L_3\}$ have exactly one common transversal, namely M , and $\{L_1, L_2, L_3\}$ and $\{L', L_2, L_3\}$ have exactly two common transversals, namely M and the unique transversal of $\{L_1, L_2, L_3\}$ through $L' \cap L_1$. \square

The next theorem is the bulk of Theorem 4. Recall the definition of θ_a for a collineation θ that maps an object a to opposite, just after Theorem 1.3 in Section 1. Recall also that a proper symplectic polar space is one in characteristic different from 2.

Theorem 3.13. *Let θ be a point-domestic collineation of a large parabolic, proper symplectic, or hyperbolic polar space Δ with opposition diagram $B_{n,i}^2, 2 \leq 2i \leq n-1$, $C_{n,i}^2, 2 \leq 2i \leq n-1$, or $D_{n,i}^2, 2 \leq 2i \leq n-2$, respectively. Then θ is the product of i pairwise orthogonal long root elations, an $(I_{2n-2i}, -I_{2i})$ -homology, or the product of i pairwise orthogonal long root elations, respectively.*

Proof. We show this theorem by induction on i , the case $i = 1$ being contained in Proposition 3.11 and Theorem 3.7. Let now $i \geq 2$.

We first observe that, under the product of at most $n/2$ perpendicular root elations in a parabolic polar space of rank n , or a hyperbolic polar space of rank $n+2$, every fixed line is either pointwise fixed or a unique point on it is fixed. Moreover, every point is in a fixed line. Also, under an $(I_{2n-2i}, -I_{2i})$ -homology of a symplectic polar space of rank n , every fixed line is either pointwise fixed or exactly two points on it are fixed. Moreover, every point is on a fixed line.

The opposition diagram tells us that there is a line L mapped onto an opposite. By largeness, we can choose L such that θ_L is by induction either the product of $i-1$ pairwise orthogonal long root elations, an $(I_{2n-2i}, -I_{2i-2})$ -homology, or the product of $i-1$ pairwise orthogonal long root elations, respectively. Now let Σ be a singular 3-space through L fixed under θ_L and such that all planes through L in Σ are also fixed under θ_L . Then each plane π of Σ containing L has the property $\pi \cap \pi^\theta \neq \emptyset$. Set $K = \Sigma \cap \Sigma^\theta$ and let $K' \subseteq \Sigma$ be such that $K'^\theta = K$. First assume that K and K' are disjoint. Then the point $x \in K'$ is mapped onto the point $\langle L, x \rangle \cap K$. Hence the restriction to K' of the action of θ is given by $\text{proj}_K^L(K')$. Let M be a line in Σ disjoint from $K \cup K'$. We claim that M is opposite M^θ . Suppose not and let $u \in M^\theta$ be a point collinear to all points of M . Note that $u \notin K$ and $\langle M, K \rangle = \Sigma$, which now contradicts the fact that u is collinear to all points of $M \cup K$. Hence the claim. Now choosing M appropriately in Σ , Lemma 3.12 leads to a contradiction to our observation in the second paragraph of this proof. If $K' \cap K$ is some point z , then the restriction to K' of θ is given by the perspectivity with centre

$L \cap \langle K, K' \rangle$. Considering a line M intersecting K, K' in an appropriate different point (on the line $\langle z, L \cap \langle K, K' \rangle \rangle$ for the symplectic case, not on that line for the other cases), we again obtain a contradiction to our observation in the second paragraph of this proof.

So we have shown that K is fixed pointwise. Now let π be any plane through L not fixed under θ_L . Then by our observation again, π is contained in a singular 3-space Σ fixed by θ_L . We claim that $K := \Sigma \cap \Sigma^\theta$ is stabilised by θ (with exactly one or two fixed points, depending on the orthogonal or symplectic case). There are again two cases to rule out: K' (defined as before as $K'^\theta = K$) disjoint from K , and $K' \cap K$ a singleton. Let's explain the first one; the second one is much easier and left to the reader.

First the orthogonal case. Let π be the unique plane containing L in Σ fixed by θ_L . Let M be the line in π intersecting all of L, K and K' in points. We can now choose a unique line T through $M \cap L$ in the plane $\langle L, M \rangle$ such that $\text{proj}_K^T(K') = \theta/K'$. It follows that θ_T fixes all planes through T inside Σ , and the previous part of the proof now implies that $K = \Sigma \cap \Sigma^\theta$ is fixed pointwise under θ , a contradiction (as θ_L does not fix all planes through L in Σ).

Now the symplectic case goes similarly, noting that there are two planes π_1, π_2 through L fixed under θ_L . Let M_i be the unique transversal of L, K, K' in the plane π_i , then we can again appropriately choose a point z on M_1 such that, setting $T := \langle z, M_2 \cap L \rangle$, the map $\text{proj}_K^T(K')$ coincides with θ/K' . We obtain the same contradiction.

Hence we have shown that θ stabilises $W := L^\perp \cap L^{\theta^\perp}$, in which θ induces by induction, in the orthogonal case, the product σ of $i - 1$ pairwise orthogonal long root elations, say with axis the $(2i - 3)$ -dimensional singular subspace A_W , and in the proper symplectic case, an $(I_{2n-2-2i}, -I_{2i-2})$ -homology σ , say with axes the $(2n - 2i - 3)$ -dimensional singular subspace A_W and the $(2i - 3)$ -dimensional singular subspace A'_W . In any case, W contains a line L^* which is mapped onto an opposite. Playing the same game with L^* as with L above, and noting that, in the orthogonal case, the hyperbolic quadric H defined by L and L^θ is stabilised, and in the (proper) symplectic case the union $L \cup L^\theta$ is stabilised (as these are the unique lines intersecting all of the stabilised lines $\langle x, x^\theta \rangle$ with $x \in L$) we see that, in the orthogonal case, θ induces an axial elation σ' on H , say with axis the line A , or a unique $(I_2, -I_2)$ -homology σ' in the non-degenerate symplectic space induced in the span of L and L^θ , say with axes the lines A and A' .

Now we first treat the orthogonal case. Let σ' be the natural extension of σ to Δ (that is, σ' is the product of the $(i - 1)$ perpendicular long root elations generating σ , naturally extended to Δ ; note that every long root elation of W extends uniquely to a long root elation of Δ by considering the same axis and the same pair of corresponding lines). Let σ'' be the long root elation with axis A mapping L to L^θ . We claim that $\sigma'\sigma''$ restricted to $W^* := L^* \cap L^{*\perp}$ (denote it by σ^*) coincides with θ restricted to W^* (denote it by θ^*). Indeed, we know that θ^* is the product of at most $(i - 1)$ perpendicular long root elations. Consequently the set of fixed points is a singular subspace, which must hence contain A and $A_W \cap W^*$. But these generate a singular subspace A^* of dimension $2i - 3$, hence σ^* is the product of precisely $(i - 1)$ perpendicular long root elations and has axis B . We can multiply $\sigma'\sigma''$ with the root elation ρ with axis $\langle L^*, L^{*\theta} \rangle \cap A_W$, which is a line, mapping $L^{*\theta}$ to L^* . Since ρ induces the identity on $L^* \cap L^{*\perp}$, we deduce that also σ^* is the product of at most $(i - 1)$ perpendicular long root elations. But its fixed point set also contains B and hence B is the axis of σ^* . Since σ^* and θ^* agree over $W \cap W^*$ and both map L to L^θ , the claim is proved. Hence $\sigma'\sigma''$ coincides with θ over $W \cup W^*$. By [13, Proposition 5.4] we obtain $\sigma'\sigma'' = \theta$ and θ is the product of i perpendicular long root elations, as claimed.

Now we treat the proper symplectic case. Here, it is more convenient to work algebraically. We can choose the standard alternating bilinear form so that it also has standard form in W and in H . Then it follows from the foregoing that there exists $k \in \mathbb{K}$ such that the matrix of θ is diagonal, containing on the diagonal $2n - 2i - 2$ times 1, $2i - 2$ times -1 , two times k and two times $-k$. But clearly, in order θ to be an automorphism of Δ , we should have $k \in \{1, -1\}$ and the assertion follows. \square

We now extend Theorem 3.13 in the symplectic case to diagrams $C_{2n,n}^2$, thereby completing the proof of Theorem 4. Note that domestic collineations with that opposition diagram exist which do not fix any chamber. But if we require at least one fixed point, then we have a well-defined involutive homology.

Theorem 3.14. *Let θ be a point-domestic collineation of a large proper symplectic polar space Δ with opposition diagram $C_{n,i}^2$ where $2 \leq 2i \leq n$, fixing at least one point if $n = 2i$. Then θ is an $(I_{2n-2i}, -I_{2i})$ -homology.*

Proof. If $2i < n$, then this is proved in Theorem 3.13, so assume from now on that $n = 2i$. We proceed by induction on n . For $n = 2$, the theorem follows from Theorem 3.7. Now assume $n > 1$.

Let x be a fixed point. It is easy to see that point-domesticity implies that the map $\theta_{(x)}$ induced by θ in $\text{Res}_S(x)$ is also point-domestic. We claim that some plane π through x is such that π, π^θ is an opposite pair in $\text{Res}_\Delta(x)$. Indeed, suppose not. Then $\theta_{(x)}$ is point- and line-domestic and hence the identity. Let X be a non-domestic 3-space (which exists by the opposition diagram). Then it is easy to see that $x^\perp \cap X$ intersects $x^\perp \cap X^\theta$ as x and $x^\perp \cap X$ span a singular subspace of dimension $1 + \dim(x^\perp \cap X)$ which is moreover fixed under the action of θ . This is a contradiction, which proves the claim.

Now let L be a line in π not through x . Then L^θ is opposite L in Δ . The map θ_L fixes the plane spanned by L and x and therefore induction implies that it is either the identity, or an $(I_{n-2}, -I_{n-2})$ -homology. Either way, similarly as in the proof of Theorem 3.13 one now shows that $\Delta_L := L^\perp \cap (L^\theta)^\perp$ is stabilised by θ . Suppose Δ_L is fixed pointwise (in other words, θ_L is the identity). Then every singular subspace of dimension at least 4 is domestic, a contradiction if $n \geq 6$. So if $n \geq 6$, then θ induces an $(I_{n-2}, -I_{n-2})$ -homology in Δ_L . Pick, in this case, a non-domestic line M in Δ_L . Considering an embedding of Δ in $\text{PG}(2n-1, \mathbb{K})$, we see that $M^\perp \cap \Delta_L$ has dimension $2n-7$ and hence intersects the pointwise fixed subspaces in Δ_L , which have dimension $n-3$, nontrivially if $(2n-7) + (n-3) - (2n-4) \geq 0$, hence for $n \geq 6$. Consequently, if $n \geq 6$, then θ_M has a fixed point and we can again apply induction and show that $\Delta_M := M^\perp \cap (M^\theta)^\perp$ is fixed; moreover θ induces an $(I_{n-2}, -I_{n-2})$ -homology in Δ_M . The homologies induced in Δ_L and Δ_M agree on the non-empty intersection (which has dimension at least 3), and since Δ_L and Δ_M span the whole space, θ is itself an $(I_n, -I_n)$ -homology.

We are left with the case $n = 4$. Assume for a contradiction that Δ_L is fixed pointwise. Then Δ_L^\perp is a symplectic quadrangle stabilised by θ and Theorem 3.7 implies that θ either fixes a spread \mathcal{S} or is an $(I_2, -I_2)$ -homology. In the former case, let $S \in \mathcal{S}$ be a line of the spread and p any point of Δ_L . Then the singular plane π spanned by S and p is fixed by θ whereas no point on L is fixed. But for an arbitrary point $q \in \pi \setminus (S \cup \{p\})$ we have $q \neq q^\theta$ and qq^θ is fixed, implying $S \cap qq^\theta$ is fixed, a contradiction. Hence θ induces an $(I_2, -I_2)$ -homology in Δ_L^\perp , say with axes A and A' . Now Propositions 3.1 and 3.3(iii) imply that either $\langle A, \Delta_L \rangle$ or $\langle A', \Delta_L \rangle$ are fixed pointwise. This means that θ is an $(I_6, -I_2)$ -homology and hence has opposition diagram $\mathcal{C}_{4,1}^2$, which is a contradiction. The claim is proved.

Hence θ induces an $(I_2, -I_2)$ -homology in Δ_L , say with corresponding axes A, A' . Let again M be a line of Δ_L mapped onto an opposite. Then $\Delta_M = \Delta_L^\perp$ and interchanging the roles of L and M we see that θ induces an $(I_2, -I_2)$ -homology in Δ_M , say with axes B, B' . Then Propositions 3.1 and 3.3(iii) imply that either $\langle A, B \rangle$ and $\langle A', B' \rangle$ are fixed pointwise, or $\langle A, B' \rangle$ and $\langle A', B \rangle$ are. Either way, we obtain an $(I_4, -I_4)$ -homology. \square

3.2.3 General case of class II collineations

Since the rank 3 situation is completely known, we may restrict to the rank ≥ 4 case. We have the following starting points.

Proposition 3.15. *Let $\Delta = (X, \Omega)$ be a polar space of rank $n \geq 4$. Let θ be a point-domestic collineation of Δ . Then all planes of Δ are Pappian, i.e., Δ is defined over a field, and θ has as companion field automorphism the one related to the corresponding Hermitian form (hence trivial for symplectic and orthogonal polar spaces, and an involution for Hermitian polar spaces). If θ is point-domestic, $(n-1)$ -domestic and $(n-2)$ -domestic, then Δ is always a symplectic or orthogonal polar space.*

Proof. The opposition diagram tells us that there is some singular subspace Σ of dimension 3 mapped to an opposite. Then θ_Σ is a domestic duality, which by largeness is a symplectic polarity (anyway, if the polar space would be small, then it is also defined over a field). These only exist in Pappian projective spaces. Now pick an arbitrary line L in Σ that is not fixed by the symplectic polarity. Then L and L^θ are opposite, and the 3-dimensional subspace W in the ambient projective space of Δ generated by L and L^θ intersects Δ in a generalised quadrangle. It follows directly from the proof of Theorem 8.5.11 in [21] that the map $\kappa : L^\theta \rightarrow L : x \mapsto x^\perp \cap L$ is the restriction of a semi-linear mapping with companion field automorphism the involution σ corresponding to the Hermitian form defining Δ (and hence trivial for symplectic and orthogonal polar spaces). Let σ' be the

companion field automorphism of θ . Let L' be the projection of L^θ onto Σ . Then, for each point $x \in L$, we have $x^{\theta\kappa} = L \cap (x^\theta)^\perp = L \cap x^{\theta\varepsilon} = x$, since θ_ε is symplectic and hence $x \in x^{\theta\varepsilon}$. Since the companion field automorphism of $\theta\kappa$ is $\sigma'\sigma$, which must be the identity in view of $x^{\theta\kappa} = x$, for all $x \in L$, we deduce that $\sigma = \sigma'$.

Left to show is the last assertion. Suppose, to that end, that θ is also $(n - 1)$ -domestic and $(n - 2)$ -domestic. Theorem 6.1 of [18] implies that θ pointwise fixes a subspace of corank at most $n - 2$. Since such a subspace always contains a line, the companion field automorphism of θ is the identity, and so Δ is not Hermitian. \square

Proposition 3.16. *A point-domestic collineation θ of a polar space of rank $n \geq 3$ with a fixed point fixes a chamber (through each fixed point).*

Proof. Let x be a fixed point of θ . We proceed by induction on n . If $n = 2$, then it follows immediately from the fact that θ is point-domestic that every line through x is stabilised. Now let $n > 2$. Then by induction, it suffices to show that some line through x is fixed by θ . Let L be an arbitrary line through x and assume that L is not fixed. Then by Proposition 3.1 the plane π spanned by L and L^θ is fixed. Combined with Proposition 3.3 this yields a fixed line through x in π . The proposition follows. \square

From Proposition 3.15 emerge three cases: the symplectic case, the orthogonal case and the Hermitian case. The symplectic case is treated in Theorem 3.14. We now treat the *proper* orthogonal case, that is, orthogonal polar spaces for which the minimal embedding has no secants of size at least 3. We will show, slightly stronger than Proposition 3.16, that θ belongs to a conjugate of the group of collineations pointwise fixing the set $E_1(C)$ (in Tits' notation [20]) of all chambers adjacent to C .

Lemma 3.17. *A point-domestic collineation θ of a (strictly) orthogonal polar space Δ of rank $n \geq 3$ with a fixed point fixes a chamber C and all adjacent chambers. In particular, this happens if θ is both point-domestic and $(n - 1)$ -domestic, or if θ is point-domestic and n is odd.*

Proof. Let x be a fixed point of θ . By Proposition 3.16, θ fixes a maximal singular subspace U containing x . We can again apply Proposition 3.3. Since we know that θ is linear (see Proposition 3.15), and we have at least one fixed point (by assumption, namely x), there are only two possibilities for the fixed point structure: (1) Either θ pointwise fixes exactly two complementary subspaces W_1 and W_2 , or (2) θ induces an elation in U (possibly a trivial one).

Suppose first (1) that θ pointwise fixes two complementary subspaces W_1 and W_2 . Pick hyperplanes $V_1 \subseteq W_1$ and $V_2 \subseteq W_2$ and consider the action of θ in the residue R of the subspace V of dimension $n - 3$ spanned by V_1 and V_2 . Note that R is an orthogonal generalised quadrangle. Then θ fixes $\langle V_1, W_1 \rangle$ and $\langle V_2, W_1 \rangle$, which are points on the line U of R , and no other points on that line are fixed. But θ induces a point-domestic collineation in R . Since such collineations pointwise fix a dual geometric hyperplane, the fixed point structure in R is an ideal subquadrangle. But proper orthogonal quadrangles are line-minimal (for terminology and this result, see Section 5.9 of [21]). This contradiction shows we always have Case (2).

Hence θ induces an elation in U . So we can select a chamber C' in U such that θ pointwise fixes $E_1(C')$. Set $C = C' \cup \{U\}$. Then θ fixes C and all i -adjacent chambers of C with $1 \leq i \leq n - 1$. There remains to show that θ also fixes all n -adjacent chambers. Consider the singular subspace Σ of dimension $n - 3$ in C' . Then the induced action of θ on the residue of Σ , which is a generalised quadrangle, fixes the point corresponding to the singular subspace Σ' of dimension $n - 2$ in C' . By point-domesticity, it fixes all lines through that point, i.e., θ fixes all maximal singular subspaces containing Σ' . Hence θ fixes all chambers n -adjacent to C .

If θ is $(n - 1)$ -domestic, then it fixes a point by Lemma 2.1. If n is odd, then the opposition diagram of θ shows that θ is $(n - 1)$ -domestic. \square

In particular the previous lemma holds for all hyperbolic polar spaces.

Now we investigate the Hermitian case. A Baer subspace of a projective space Σ has the following two properties: (1) every hyperplane of Σ is incident with at least one point of Σ and (2) every point of Σ is on some line of Σ' . Let Δ' be a rank n sub polar space of the polar space Δ of rank n with the following two properties:

- (Pr1) Every maximal singular subspace of Δ' is a Baer subspace of some maximal singular subspace of Δ .
- (Pr2) Every maximal singular subspace of Δ containing a submaximal singular subspace of Δ' belongs to Δ' .

Then we call Δ' an *ideal Baer sub polar space* of Δ . This generalises the definition of an ideal Baer sub polar space of rank 3 in [18]. An involution that fixes an ideal Baer sub polar space of Δ and nothing more is called an *ideal Baer involution*.

Proposition 3.18. *A point-domestic collineation θ of a Hermitian polar space Δ of rank $n \geq 3$ with a fixed point is an ideal Baer involution and has opposition diagram $B_{n,m}^2$ with $2m \in \{n, n + 1\}$.*

Proof. Let x be any fixed point. Again by Proposition 3.16 there exists a maximal singular subspace U through x , fixed under the action of θ . Now Propositions 3.3 and 3.15 imply that the fixed point structure of θ in U is a Baer subspace Σ' , and the restriction to U of θ is an involution. Let W be a hyperplane of U spanned by a hyperplane of Σ' . Then, by point-domesticity, all maximal singular subspaces of Δ containing W are fixed by θ , and so they all contain further points fixed under θ . This shows that for each fixed point x there exist many opposite fixed points. So the fixed point structure is a non-degenerate polar space, which is easily seen to be an ideal Baer sub polar space.

Clearly θ^2 is linear and point-domestic (by Corollary 3.2) and so the identity by Proposition 3.15. Hence θ is an involution and the lemma is proved (the opposition diagrams follow from the last assertion of Proposition 3.15). □

Example 3.19. Let Δ be the rank n polar space corresponding to the Hermitian variety with the equation $\sum_{i=1}^n X_{-i}\bar{X}_i - X_i\bar{X}_{-i} = 0$ in $\text{PG}(2n - 1, \mathbb{K})$, where $x \mapsto \bar{x}$ is the corresponding field involution and \mathbb{K} is commutative. Let θ be the collineation defined by mapping a point with coordinates $(x_{-n}, x_{-n+1}, \dots, x_{-1}, x_1, \dots, x_n)$ to $(\bar{x}_{-n}, \bar{x}_{-n+1}, \dots, \bar{x}_{-1}, \bar{x}_1, \dots, \bar{x}_n)$. If we denote by \mathbb{F} the fixed field of the field involution, then the fixed points in Δ are precisely all points of a Baer subspace $\text{PG}(2n - 1, \mathbb{F})$, hence θ pointwise fixes a symplectic polar space of rank n . The collineation is point-domestic as one easily calculates.

We can now turn our attention to collineations of class III.

3.3 Class III automorphisms

If θ is a class III automorphism of a polar space of rank n then, by Corollaries 3.4 and 3.6, the rank n is even and $\text{Diag}(\theta) = X_{n;n/2}^2$ (where $X \in \{B, C, D\}$).

First we need to be a little more precise in Case (i) of Proposition 3.3.

Proposition 3.20. *Let Σ be a finite-dimensional projective space over a field \mathbb{K} with dimension at least 2 and let θ be a fixed point free collineation with the property that the line $\langle x, x^\theta \rangle$ is fixed for every point x . Then the projective dimension of Σ is odd, say $2n + 1$, and the fixed subspaces form a projective space of dimension n either over a quadratic (not necessarily separable) extension \mathbb{L} of \mathbb{K} , or over a quaternion division algebra \mathbb{H} which naturally contains \mathbb{K} as a 2-dimensional subalgebra over its centre.*

Proof. It is clear that the fixed point structure is a projective space \mathfrak{P} of dimension n . In order to determine the coordinatizing division ring, we consider a fixed 3-dimensional subspace, i.e., we consider the case $n = 1$. Applying the Klein correspondence, θ acts on a hyperbolic quadric Q in $\text{PG}(5, \mathbb{K})$ pointwise fixing an ovoid O .

If θ is linear, then any triple of points of O determines a conic on Q which is fixed pointwise (by linearity). Considering four fixed points no three on a conic we see that θ pointwise fixes a 3-dimensional subspace Σ of $\text{PG}(5, \mathbb{K})$ intersecting Q in O . Hence O is an elliptic quadric in Σ (that is, a non-degenerate nonruled quadric). Since the stabiliser of O induces a group containing $\text{PGL}_2(\mathbb{L})$ for some quadratic extension \mathbb{L} of \mathbb{K} , and since the group of projectivities of a line of \mathfrak{P} is inherited from the collineation group of a 3-dimensional subspace of Σ fixed by θ , we deduce that \mathfrak{P} is defined over \mathbb{L} .

Now suppose that θ is not linear. Note that θ does not map any point to a collinear one. We show that θ is an involution. Let x be an arbitrary point. If x is fixed by θ , then x is also fixed by θ^2 . Now suppose that x^θ is opposite to x . Let π be any singular plane through x . Since θ preserves types, π^θ intersects π in a unique point p . Clearly $p^\theta \in \pi^\theta \ni p$, hence by our remark above $p = p^\theta$. Hence the set of fixed points in $p^\perp \cap (p^\theta)^\perp$ forms an ovoid C ,

which cannot be a plane conic as θ is not linear. Hence C spans $p^\perp \cap (p^\theta)^\perp$ and so θ preserves $p^\perp \cap (p^\theta)^\perp$. Hence θ also preserves the perp of that set, which is $\{p, p^\theta\}$, implying that $p^{\theta^2} = p$. This shows that θ is an involution.

So back to \mathfrak{P} , θ acts as a Galois group and hence \mathfrak{P} arises from Galois descent, implying that the coordinatising structure is a quaternion division algebra. This can also be seen in an elementary way: denote by $\mathbb{K} \rightarrow \mathbb{K} : k \mapsto \bar{k}$ the companion field involution; then coordinates in the case $n = 1$ can be chosen such that θ maps (x_1, x_2, x_3, x_4) to $(\bar{x}_2, a\bar{x}_1, \bar{x}_4, a\bar{x}_3)$, with $a = a^\theta \notin \{x\bar{x} : x \in \mathbb{K}\}$. This determines the quaternion division ring where multiplication is given by $(x, y) \cdot (u, v) = (xu + a\bar{y}v, yu + \bar{x}v)$. \square

An immediate corollary is Theorem 5:

Corollary 3.21. *Let $\Delta = (X, \Omega)$ be a polar space of rank $n \geq 4$ defined over the field \mathbb{K} . Let θ be a point-domestic collineation of Δ without any fixed points. Let Y be the set of fixed lines, and let Υ be the set of fixed singular subspaces. Then $\Gamma = (Y, \Upsilon)$ is a polar space defined over either a quadratic extension of \mathbb{K} , or over a quaternion division algebra \mathbb{H} which is 2-dimensional over \mathbb{K} (and \mathbb{K} is 2-dimensional over the centre of \mathbb{H}).*

Now we first handle the split case (Theorem 6). Recall the definition of a *minimal* Hermitian polar space of rank n : one with ambient projective space of dimension $2n - 1$. A *mixed* polar space (of rank n) is a polar space which is a proper subspace of a symplectic polar space of rank n . Here, the characteristic is necessarily equal to 2.

Theorem 3.22. *Let $\Delta = (X, \Omega)$ be a split polar space of rank $n \geq 4$ defined over the field \mathbb{K} . Let θ be a point-domestic collineation of Δ without any fixed point. Then n is even, say $n = 2m$, and the opposition diagram is either $C_{n,m}^2$ or $D_{n,m}^2$. Moreover, Δ is either*

- *a symplectic polar space, and the fixed point structure of θ is a symplectic polar space over a quadratic extension of \mathbb{K} , or*
- *a hyperbolic orthogonal polar space, and the fixed point structure of θ is either a minimal Hermitian polar space, or a mixed polar space.*

Proof. The symplectic case immediately follows from Proposition 3.20 and the fact that symplectic polar spaces are the only embedded polar spaces with the property that all points of the ambient projective space belong to the polar space.

In the hyperbolic case it suffices to consider the rank $n = 4$. Applying triality, we may assume that θ has opposition diagram $D_{4,2}^1$, in which case the fixed point structure is a subspace of corank 2, hence an orthogonal generalised quadrangle naturally embedded in projective 5-space and, by Remark 3.4.10 of [21], either a mixed quadrangle, or the dual of a minimal Hermitian quadrangle.

In the parabolic case, we consider two opposite maximal singular subspaces fixed by θ . Then θ stabilises the hyperbolic polar subspace H generated by those. A fixed line not contained in H intersects H in a point (as H is a geometric hyperplane), which must subsequently be fixed, yielding a contradiction. \square

Note that the previous theorem says that no parabolic polar space admits a point-domestic collineation without fixed points. Hence in the symplectic case, the defining field cannot be perfect when it has characteristic 2. In fact, we can even be more specific concerning the collineations and the said quadratic extensions. The next proposition is Theorem 7.

Proposition 3.23. *Let $\Delta = (X, \Omega)$ be a symplectic polar space of rank $n \geq 4$ defined over the field \mathbb{K} . Let θ be a point-domestic collineation of Δ without any fixed point. Let \mathbb{L} be the quadratic extension of \mathbb{K} over which the fixed points structure of θ is defined according to Theorem 3.22. Then θ is an involution and either*

- *the characteristic of \mathbb{K} is not 2 and \mathbb{L} is a separable extension of \mathbb{K} , or*
- *the characteristic of \mathbb{K} is 2 and \mathbb{L} is an inseparable extension of \mathbb{K} .*

Conversely, if \mathbb{K} admits a quadratic extension \mathbb{L}/\mathbb{K} , inseparable if the characteristic of \mathbb{K} is 2, then Δ admits a point-domestic collineation without fixed points as above.

Proof. Clearly it suffices to show that, if in a symplectic quadrangle over \mathbb{K} a spread is fixed by a fixed point free collineation θ , then θ is an involution and the spread corresponds to an inseparable extension if the characteristic of \mathbb{K} is 2.

Let the symplectic quadrangle Δ be defined by the standard symplectic form $x_0y_1 - x_1y_0 + x_2y_3 - x_3y_2$. Without loss of generality we may assume that θ (which has trivial companion field automorphism by Proposition 3.15) maps the point $(1, 0, 0, 0)$ to $(0, 0, 0, 1)$ and $(0, 0, 1, 0)$ to $(0, 1, 0, 0)$. Hence $(0, 1, 0, 0)$ is mapped to $(0, a, b, 0)$, and $(0, 0, 0, 1)$ to $(c, 0, 0, d)$, yielding the matrix

$$\begin{pmatrix} 0 & 0 & 0 & c \\ 0 & a & e & 0 \\ 0 & b & 0 & 0 \\ 1 & 0 & 0 & d \end{pmatrix},$$

for some $a, b, c, d, e \in \mathbb{K}$. Expressing that θ maps each point (x_0, x_1, x_2, x_3) to a collinear one, we obtain

$$cx_3x_1 - (ax_1 + ex_2)x_0 + bx_1x_3 - (x_0 + dx_3)x_2 \equiv 0.$$

This yields $e = -1, c = -b$ and $a = d = 0$.

Since θ has no fixed points, b is a nonsquare and we see that $\mathbb{L} = \mathbb{K}(\sqrt{b})$, which proves the assertions (the converse is obvious in view of our explicit form of θ). \square

The hyperbolic case is even more canonical. This is Theorem 8 from the introduction.

Proposition 3.24. *Let $\Delta = (X, \Omega)$ be a hyperbolic polar space of rank $n \geq 4$ defined over the field \mathbb{K} . Let θ be a point-domestic collineation of Δ without any fixed point. Let $\text{PG}(2n - 1, \mathbb{K})$ be the ambient projective space. Then θ naturally extends to $\text{PG}(2n - 1, \mathbb{K})$ pointwise fixing a spread S which defines a projective space $\text{PG}(n - 1, \mathbb{L})$ over a quadratic extension \mathbb{L} of \mathbb{K} . Every collineation of $\text{PG}(2n - 1, \mathbb{K})$ with pointwise fixes S is a point-domestic collineation of Δ without fixed points. If \mathbb{L}/\mathbb{K} is separable, then the fixed polar space is minimal Hermitian; if \mathbb{L}/\mathbb{K} is inseparable, then the fixed polar space is the subspace of the symplectic polar space over \mathbb{L} obtained by restricting the long root elations to \mathbb{K} .*

Conversely, if \mathbb{K} admits a quadratic extension \mathbb{L}/\mathbb{K} , then there exists a point-domestic collineation of Δ without fixed points as above.

Proof. Again it suffices to consider the case $n = 4$. We order coordinates as in

$$(x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_1, x_2, x_3, x_4),$$

take X to be the set of points with equation $x_{-4}x_4 - x_{-3}x_3 + x_{-2}x_2 - x_{-1}x_1 = 0$, and assume that θ stabilises the lines $\langle p_{-4}, p_{-3} \rangle, \langle p_4, p_3 \rangle, \langle p_{-2}, p_{-1} \rangle, \langle p_2, p_1 \rangle$, where p_i is the point with a 1 in position i and 0 elsewhere. Set

$$M_i = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix},$$

$a_i, b_i, c_i, d_i \in \mathbb{K}, i = -2, -1, 1, 2$. Then we can assume that θ has block matrix

$$\begin{pmatrix} M_{-2} & 0 & 0 & 0 \\ 0 & M_{-1} & 0 & 0 \\ 0 & 0 & M_1 & 0 \\ 0 & 0 & 0 & M_2 \end{pmatrix}.$$

(Recall from Proposition 3.15 that θ is linear; action of θ on the left)

Since θ has no fixed points, we have $b_i \neq 0 \neq c_i$ for all $i \in \{-2, -1, 1, 2\}$. Expressing that θ preserves the quadric (hence the quadratic form up to a scalar multiple), we deduce that $\det A_{-2} = \det A_{-1} = \det A_1 = \det A_2 := d$. Expressing that θ maps each point to a collinear one we deduce $\text{tr } A_{-2} = \text{tr } A_{-1} = \text{tr } A_1 = \text{tr } A_2 := t$. We also have $A_{-i} = A_i$ for $i = 1, 2$. Moreover, since there are no fixed points in Δ , the equation $x^2 - tx + d = 0$ has no solution in \mathbb{K} and determines a quadratic field extension \mathbb{L}/\mathbb{K} . It follows that θ pointwise fixes a spread

\mathcal{S} of $\text{PG}(7, \mathbb{K})$. Clearly, every collineation of $\text{PG}(7, \mathbb{K})$ fixing that spread pointwise restricts to a point-domestic collineation of Δ .

Let δ be a solution in \mathbb{L} of the equation $x^2 - tx + d = 0$ above. Then we can write every element of \mathbb{L} as $a + b\delta$ with $a, b \in \mathbb{K}$ and so with $\delta^2 = t\delta - d$. Now we identify each point $p = (x_{-4}, x_{-3}, \dots, x_3, x_4)$ of $\text{PG}(7, \mathbb{K})$ with the point $p^\beta = (c_2x_{-4} - a_2x_{-3} + x_{-3}\delta, c_1x_{-2} - a_1x_{-1} + x_{-1}\delta, c_1x_1 - a_1x_2 + x_2\delta, c_2x_3 - a_2x_4 + x_4\delta)$ of $\text{PG}(3, \mathbb{L})$. An elementary computation shows that the image of a point under the action of θ is identified with the same point of $\text{PG}(3, \mathbb{L})$ (the coordinates are equal up to the scalar δ). Hence we can identify the fixed lines of $\text{PG}(7, \mathbb{K})$ with the points of $\text{PG}(3, \mathbb{L})$. Denote the conjugate $u + tv - v\delta$ of the element $u + v\delta \in \mathbb{L}$ by $\bar{u} + v\bar{\delta}$, where $u, v \in \mathbb{K}$. Then one calculates that a point p of $\text{PG}(7, \mathbb{K})$ belongs to Δ if and only if the coordinates $(y_{-2}, y_{-1}, y_1, y_2)$ of p^β satisfy the relation $c_1Y_{-2}\bar{Y}_2 + c_2Y_{-1}\bar{Y}_1 \in \mathbb{K}$.

This proves the proposition (noting that the converse is again obvious by our explicit form of θ). □

We now consider non-split polar spaces. The neatest situation occurs with the minimal Hermitian polar spaces. The following is Theorem 9.

Proposition 3.25. *Let $\Delta = (X, \Omega)$ be a minimal Hermitian polar space of rank $n \geq 4$ defined over the field \mathbb{K} , with corresponding field involution σ and field extension \mathbb{K}/\mathbb{F} . Let θ be a point-domestic collineation of Δ without any fixed point. Let $\text{PG}(2n - 1, \mathbb{K})$ be the ambient projective space. Then θ is an involution and naturally extends to $\text{PG}(2n - 1, \mathbb{K})$ pointwise fixing a spread \mathcal{S} which defines a projective space $\text{PG}(n - 1, \mathbb{H})$ over a quaternion division algebra \mathbb{H} over its centre \mathbb{F} , with \mathbb{K} a 2-dimensional subalgebra of \mathbb{H} . The only nontrivial collineation of Δ fixing every member of \mathcal{S} in Δ is θ . The fixed polar space is a minimal quaternion polar space related to a nonstandard involution of \mathbb{H} pointwise fixing a 3-dimensional subalgebra over \mathbb{F} .*

Conversely, if \mathbb{K} admits a quaternion extension \mathbb{H} such that the centre \mathbb{F} of \mathbb{H} is a subfield of \mathbb{K} and \mathbb{K}/\mathbb{F} is a separable quadratic field extension, then there exists a point-domestic collineation of Δ without fixed points as above.

Proof. Note that by Proposition 3.15, θ is a involution. Also, if another involution θ' of Δ would fix exactly the same lines as θ , then $\theta\theta'$ is a linear point-domestic collineation of Δ , implying the stated uniqueness of θ . We may again only consider the case $n = 4$. The arguments run parallel to those of the proof of Proposition 3.24. We order coordinates as in

$$(x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_1, x_2, x_3, x_4),$$

take X to be the set of points with equation

$$X_{-4}X_4^\sigma + X_4X_{-4}^\sigma - X_{-3}X_3^\sigma - X_3X_{-3}^\sigma + X_{-2}X_2^\sigma + X_2X_{-2}^\sigma - X_{-1}X_1^\sigma - X_1X_{-1}^\sigma = 0,$$

and assume that θ stabilises the lines $\langle p_{-4}, p_{-3} \rangle, \langle p_4, p_3 \rangle, \langle p_{-2}, p_{-1} \rangle, \langle p_2, p_1 \rangle$, where p_i is the point with a 1 in position i and 0 elsewhere.

Since θ is an involution we may assume that it interchanges p_{-4} with p_{-3} , p_4 with p_3 , p_{-2} with p_{-1} , and p_2 with p_1 . Hence it has block matrix

$$\begin{pmatrix} A_{-2} & 0 & 0 & 0 \\ 0 & A_{-1} & 0 & 0 \\ 0 & 0 & A_1 & 0 \\ 0 & 0 & 0 & A_2 \end{pmatrix} \quad \text{with} \quad M_i = \begin{pmatrix} 0 & a_i \\ b_i & 0 \end{pmatrix},$$

$a_i, b_i \in \mathbb{K}$, $i = -2, -1, 1, 2$. Expressing that θ preserves X and that θ maps points of Δ to collinear points readily implies that $a_{-i} = a_i$ and $b_{-i} = b_i$ for $i = 1, 2$, and $a_1b_1^\sigma = a_2b_2^\sigma = b_1a_1^\sigma = b_2a_2^\sigma =: \ell \in \mathbb{F}$. Then θ has no fixed points in Δ if and only if $\ell \notin \{kk^\sigma : k \in \mathbb{K}\}$. It is easily checked that this now defines a fixed point free involution in $\text{PG}(7, \mathbb{K})$, pointwise fixing a spread \mathcal{S} .

We can define the quaternion division algebra $\mathbb{H} := \{u + v\delta : (u, v) \in \mathbb{K} \times \mathbb{K}\}$ with ordinary addition and multiplication determined by $\delta v = v^\sigma \delta$ and $\delta^2 = \ell$. Clearly \mathbb{F} is the centre of \mathbb{H} and \mathbb{K} is a 2-dimensional division subalgebra over \mathbb{F} . Now we identify each point $p = (x_{-4}, x_{-3}, \dots, x_3, x_4)$ of $\text{PG}(7, \mathbb{K})$ with the point $p^\beta = (a_2^\sigma x_{-4} + x_{-3}\delta, a_1^\sigma x_{-2} + x_{-1}\delta, a_1^\sigma x_1 + x_2\delta, a_2^\sigma x_3 + x_4\delta)$ of the left projective space $\text{PG}(3, \mathbb{H})$. An elementary

computation shows that the image of a point under the action of θ is identified with the same point of $\text{PG}(3, \mathbb{H})$ (the coordinates are equal up to the left scalar δ). Hence we can identify the fixed lines of $\text{PG}(7, \mathbb{K})$ with the points of $\text{PG}(3, \mathbb{H})$. Denote the non-standard involution $u - v^\sigma \delta$ of the element $u + v\delta \in \mathbb{H}$ by $\overline{u + v\delta}$, where $u, v \in \mathbb{K}$. Then one calculates that a point p of $\text{PG}(7, \mathbb{K})$ belongs to Δ if and only if the coordinates $(y_{-2}, y_{-1}, y_1, y_2)$ of p^β satisfy the relation $Y_{-2}a_2^{-\sigma}\overline{Y}_2 + Y_{-1}a_1^{-\sigma}\overline{Y}_1 = Y_2a_2^{-\sigma}\overline{Y}_{-2} + Y_1a_1^{-\sigma}\overline{Y}_{-1}$.

This proves the proposition (noting that the converse is again obvious by our explicit form of θ). □

Left to consider are the orthogonal and Hermitian polar spaces whose standard form has a nontrivial anisotropic part. These contain hyperbolic and minimal Hermitian polar subspaces fixed under the given point-domestic collineation, and so Propositions 3.24 and 3.25 provide necessary conditions for the existence. In general, if a polar subspace of hyperbolic or minimal Hermitian type of an arbitrary polar space admits a point-domestic collineation without fixed points, then it depends on the shape of the anisotropic form whether this collineation can be extended as a point-domestic collineation without fixed points, and the general rule is that the anisotropic form must, up to recoordination, be the sum of multiples of the norm form of the corresponding field extension or quaternion division algebra, respectively. Let us give a few examples for particular fields to illustrate this.

Some special fields

Quadrics over the reals. Consider a real quadric of Witt index $2n$ in a projective space of dimension $4n + 2k - 1$. Its standard equation is

$$X_{-2n}X_{2n} + X_{-2n+1}X_{2n-1} + \dots + X_{-1}X_1 = X_{0,-k}^2 + X_{0,-k+1}^2 + \dots + X_{0,-1}^2 + X_{0,1}^2 + \dots + X_{0,k}^2.$$

Let M be a real 2×2 matrix with non-real eigenvalues. By multiplying with an appropriate scalar we may assume that its determinant is 1. Then its trace is of the form $2 \cos \varphi$, for some $\varphi \in]0, \pi[$. Then the collineation defined by

$$\left\{ \begin{array}{l} \begin{pmatrix} X_{-2\ell} \\ X_{-2\ell+1} \end{pmatrix} \mapsto M \cdot \begin{pmatrix} X_{-2\ell} \\ X_{-2\ell+1} \end{pmatrix}, \quad \text{for } \ell = 1, 2, \dots, n, \\ \begin{pmatrix} X_{2\ell-1} \\ X_{2\ell} \end{pmatrix} \mapsto M \cdot \begin{pmatrix} X_{2\ell-1} \\ X_{2\ell} \end{pmatrix}, \quad \text{for } \ell = 1, 2, \dots, n, \\ \begin{pmatrix} X_{0,-\ell} \\ X_{0,\ell} \end{pmatrix} \mapsto \begin{pmatrix} \cos \varphi & \pm \sin \varphi \\ \mp \sin \varphi & \cos \varphi \end{pmatrix} \cdot \begin{pmatrix} X_{0,-\ell} \\ X_{0,\ell} \end{pmatrix}, \quad \text{for } \ell = 1, 2, \dots, k \end{array} \right.$$

is point-domestic and fixed point free. The fix point structure is a Hermitian polar space in $\text{PG}(2n + k - 1, \mathbb{C})$ with equation (and previous conventions)

$$Y_{-n}\overline{Y}_n - Y_n\overline{Y}_{-n} + \dots + Y_{-1}\overline{Y}_1 - Y_1\overline{Y}_{-1} = Y_{0,k}\overline{Y}_{0,k} + \dots + Y_{0,1}\overline{Y}_{0,1}.$$

Quadrics over the rationals. Consider a rational quadric of Witt index 4 in a projective space $\text{PG}(11, \mathbb{Q})$ with equation

$$X_{-4}X_4 + X_{-3}X_3 + X_{-2}X_2 + X_{-1}X_1 = X_{0,-2}^2 + X_{0,-1}^2 + X_{0,1}^2 + \ell X_{0,2}^2$$

with $\ell \in \mathbb{Q}$. If ℓ is a perfect square, then we can set $X'_{0,2} = \sqrt{\ell}X_{0,2}$ and similarly as before we can construct fixed point free point-domestic collineations. However, if ℓ is not a perfect square, then $X_{0,-2}^2 + \ell X_{0,2}^2$ and $X_{0,-1}^2 + X_{0,1}^2$ do not define the norm of the same quadratic extension of \mathbb{Q} , and indeed one can easily show that no fixed point free point-domestic collineation exists.

Complex Hermitian polar spaces. Here similar considerations hold as for the real quadrics. Let a Hermitian variety of Witt index $2n$ in the projective space $\text{PG}(4n + 2k - 1, \mathbb{C})$ be given by the (standard) equation

$$\sum_{\ell=1}^{-2n} X_{-2\ell}\overline{X}_{2\ell} + X_{2\ell}\overline{X}_{-2\ell} - X_{-2\ell+1}\overline{X}_{2\ell-1} - X_{2\ell-1}\overline{X}_{-2\ell+1} = \sum_{\ell=1}^k X_{0,-\ell}\overline{X}_{0,-\ell} + X_{0,\ell}\overline{X}_{0,\ell}.$$

Then, for each $r \in \mathbb{R}^+$, the semi-linear involution defined by

$$\begin{cases} \begin{pmatrix} x_{-2\ell} \\ x_{-2\ell+1} \end{pmatrix} \mapsto \begin{pmatrix} \bar{x}_{-2\ell+1} \\ -r \cdot \bar{x}_{-2\ell} \end{pmatrix}, & \text{for } \ell = 1, 2, \dots, n, \\ \begin{pmatrix} x_{2\ell-1} \\ x_{2\ell} \end{pmatrix} \mapsto \begin{pmatrix} \bar{x}_{2\ell} \\ -r \cdot \bar{x}_{2\ell-1} \end{pmatrix}, & \text{for } \ell = 1, 2, \dots, n, \\ \begin{pmatrix} x_{0,-\ell} \\ x_{0,\ell} \end{pmatrix} \mapsto \begin{pmatrix} \pm \sqrt{r} \cdot \bar{x}_{0,\ell} \\ \mp \sqrt{r} \cdot \bar{x}_{0,-\ell} \end{pmatrix}, & \text{for } \ell = 1, 2, \dots, k \end{cases}$$

is point-domestic and fixed point free.

Now a property similar to that of rational quadrics explained above holds for complex rational Hermitian varieties, and we will not repeat it.

Finite fields. The previous considerations allow us now to conclude that a finite polar space Δ of rank at least 3 admits a fixed point free point-domestic collineation if and only if it is a symplectic polar space in odd characteristic, or it is a hyperbolic or elliptic quadric. Moreover, all such collineations can be explicitly written down on appropriate standard forms.

Remark 3.26. The inverse procedure, namely, to start with a Hermitian form over a field extension \mathbb{L}/\mathbb{K} or over a quaternion division algebra \mathbb{H} with 2-dimensional subfield \mathbb{K} , and to consider it over the subfield \mathbb{K} to obtain a quadratic form or a Hermitian form, respectively, is well known (sometimes called *field reduction*). However, other situations turn up here as this is not always related to a collineation and its fixed point structure (e.g., when the 2-dimensional subfield \mathbb{K} of the quaternion division algebra \mathbb{H} is an inseparable extension of the ground field of the algebra). But when it is related to a collineation and its fixed point structure, then it is always a point-domestic fixed point free collineation.

4 Polar closed diagrams

In this section we prove Theorem 10, classifying those admissible Dynkin diagrams which arise as the opposition diagram of a unipotent element of a split spherical building. The result for the exceptional types E_n ($n = 6, 7, 8$), F_4 and G_2 is [13, Theorem 5], and so here we consider the classical types. We will also prove Theorem 11.

We shall need the following lemma. A *non-degenerate generalised polarity* ρ in a projective space $\text{PG}(V)$ is a symmetric relation in the point set such that for every point p the set p^ρ of points in relation with p is a hyperplane of $\text{PG}(V)$. We say that a polar space Δ is *embedded in a non-degenerate generalised polarity* ρ of some projective space $\text{PG}(V)$ if each point p of Δ is a point of $\text{PG}(V)$ and if the hyperplane spanned by p^\perp coincides precisely with p^ρ .

Lemma 4.1. *Let Γ be any polar space of rank n embedded in a non-degenerate generalised polarity ρ of a projective space $\text{PG}(V)$ over \mathbb{L} such that Γ spans $\text{PG}(V)$. Let $0 \leq i \leq n - 1$. Then no geometric subspace of corank i is contained in a subspace of codimension $i + 1$.*

Proof. Suppose for a contradiction that S is a subspace of $\text{PG}(V)$ of codimension $i + 1$ containing a geometric subspace of Γ of corank i . Since Γ is embedded in ρ , every point x of Γ outside S^ρ has the property that $x^\perp \cap S$ is a subspace of codimension $i + 2$. Let x_1 be such a point (which certainly exists as Γ is not contained in S^ρ by assumption). In $\Gamma_1 := \text{Res}(x_1)$ this means that we obtain a subspace S_1 with $\text{codim} S_1 = i + 2$ intersecting every singular subspace of dimension $(i - 1)$ nontrivially. Since $\text{Res}(x_1)$ is embedded in a non-degenerate polarity of $\text{PG}(V_2)$ with $\text{codim} V_2 = 2$ and spans $\text{PG}(V_2)$, we can apply the same argument and continue like this until we reach a polar space Γ_i of rank $n - i$ spanning $\text{PG}(V_{2i})$, with $\text{codim} V_{2i} = 2i$, and a subspace S_i of codimension $2i + 1$ in $\text{PG}(V)$, hence codimension 1 in $\text{PG}(V_{2i})$, intersecting every subspace of dimension 0 of Γ_i nontrivially. In other words, Γ_i is contained in S_i , which is a contradiction. \square

We can now show that the opposition diagram $B_{n;i}^1$ for odd i does not occur as the opposition diagram of any unipotent element in a split building of type B_n . In fact, we can prove a slightly more general result by considering all proper orthogonal polar spaces. Recall that a *proper orthogonal* polar space is a polar space for which the minimal embedding has no secants of size at least 3. These include all the quadrics of Witt index at least 2 in characteristic different from 2, and all hyperbolic ones.

Proposition 4.2. *Let Δ be a proper orthogonal polar space standard embedded in some projective space $PG(V)$ and let θ be a collineation of Δ with opposition diagram $B_{n;i}^1$ for some odd i with $1 \leq i \leq n - 1$. Then θ is not a unipotent element.*

Proof. Suppose for a contradiction that θ is a unipotent element of Δ with opposition diagram $B_{n;i}^1$ for some odd i with $1 \leq i \leq n - 1$. By [18, Theorem 6.1] and Lemma 2.1, θ pointwise fixes a geometric subspace of corank i , which, by Lemma 4.1 generates a subspace S of $PG(V)$ of codimension at most i . Since θ is unipotent, the set of fixed points of θ is a subspace T , and we claim it is S . Indeed, since S is spanned by the pointwise fixed subspace of corank i , it is pointwise fixed (this also follows from the proof of Theorem 6.1 in [18]). Suppose now that $\text{codim} T < i$. Then a random singular subspace of dimension $i - 1$ intersects T in at least a point and thus cannot be mapped onto an opposite, which is a contradiction. So $S = T$ and $\text{codim} S = i$.

Since S is an axis for θ , the subspace S^\perp is a centre for θ , that is, all subspaces of $PG(V)$ containing S^θ are fixed. Since unipotent elements have incident pairs (axis, centre), we have $S^\perp \subseteq S$. This now implies that S^\perp is a singular subspace of Δ with dimension $i - 1$. Since it acts as the centre of θ , the latter maps an $(i - 1)$ -dimensional singular subspace W of Δ disjoint from S onto a disjoint singular subspace of dimension $i - 1$ contained in $\langle S, W \rangle$. Hence W and W^θ are two disjoint maximal singular subspaces of the same type of the hyperbolic quadric $\langle S, W \rangle \cap \Delta$ of rank i , implying that i is even, which is a contradiction. \square

We can now prove one direction of Theorem 10.

Corollary 4.3. *Let θ be an automorphism of a split spherical building. If the opposition diagram of θ is not polar closed, then θ is not a unipotent element.*

Proof. Recall that all admissible diagrams of classical type are polar closed except for those of type $A_{n;(n-1)/2}^2$ (n odd), $A_{n;n}^1$, $B_{n;i}^1$ or $D_{n;i}^1$ (with i odd and $1 \leq i < n$), $C_{n;i}^2$ (with $1 \leq i \leq n/2$), and those in Table 2. The diagrams $A_{n;(n-1)/2}^2$ (n odd), $A_{n;n}^1$, $B_{n;i}^1$ (with i odd and $1 \leq i < n$), and those in Table 2 all correspond to non-type preserving automorphisms of Δ , and hence cannot arise as the opposition diagram of a conjugate of U . Proposition 4.2 and Theorem 3.14 show that the remaining non-polar closed diagrams $B_{n;i}^1$ (i odd) and $C_{n;i}^2$ cannot be obtained as the opposition diagrams of unipotent elements. \square

We must now prove the converse. To do so, we briefly recall explicit matrix realisations of the elements $x_\alpha(a)$ for split classical groups, following the conventions in [4] (note that this leads to different quadratic forms than those listed in Section 1.6). In each case we will give matrices for $x_\alpha(a)$ with $a \in \Phi^+$, and the corresponding elements $x_{-\alpha}(a)$ are obtained by taking the transpose.

For type A_n the root system has simple roots $\alpha_i = e_i - e_{i+1}$ for $1 \leq i \leq n$, and the highest root is $\varphi = e_1 - e_{n+1}$. The Chevalley generators are (for $a \in \mathbb{K}$) $x_{e_i - e_j}(a) = 1 + aE_{i,j}$ for $1 \leq i < j \leq n + 1$.

For type B_n the root system has simple roots $\alpha_i = e_i - e_{i+1}$ for $1 \leq i \leq n - 1$ and $\alpha_n = e_n$, and the highest root is $\varphi = e_1 + e_2$. Let (\cdot, \cdot) be the symmetric bilinear form defined on \mathbb{K}^{2n+1} by

$$(x, y) = 2x_0y_0 + x_1y_{n+1} + \cdots + x_ny_{2n} + x_{n+1}y_1 + \cdots + x_{2n}y_n$$

(indexing of rows and columns of vectors and matrices begins at 0 here). Then $\Delta = B_n(\mathbb{K})$ can be realised as the polar space associated to the quadric $f(x) = (x, x)$. The Chevalley generators are ([4, Section 11.2])

$$\begin{aligned} x_{e_i - e_j}(a) &= 1 + a(E_{i,j} - E_{n+j, n+i}) \\ x_{e_i + e_j}(a) &= 1 + a(E_{i, n+j} - E_{j, n+i}) \\ x_{e_k}(a) &= 1 + 2aE_{k,0} - aE_{0, n+k} - a^2E_{k, n+k}. \end{aligned}$$

For type C_n the root system has simple roots $\alpha_i = e_i - e_{i+1}$ for $1 \leq i \leq n - 1$ and $\alpha_n = 2e_n$, and the highest root is $\varphi = 2e_1$. Let (\cdot, \cdot) be the symplectic form on \mathbb{K}^{2n} given by

$$(x, y) = x_1y_{n+1} + \cdots + x_ny_{2n} - x_{n+1}y_1 - \cdots - x_{2n}y_n.$$

We write the standard basis of \mathbb{K}^{2n} as $\varepsilon_1, \dots, \varepsilon_n, f_1, \dots, f_n$, so that $(\varepsilon_i, \varepsilon_j) = (f_i, f_j) = 0$ for $1 \leq i, j \leq n$ and $(\varepsilon_i, f_j) = \delta_{i,j}$. The Chevalley generators are

$$\begin{aligned} x_{e_i - e_j}(a) &= 1 + a(E_{i,j} - E_{n+j, n+i}) \\ x_{e_i + e_j}(a) &= 1 + a(E_{i, n+j} + E_{j, n+i}) \\ x_{2e_k}(a) &= 1 + aE_{k, n+k}. \end{aligned}$$

For type D_n with $n \geq 4$ the root system has simple roots $\alpha_i = e_i - e_{i+1}$ for $1 \leq i \leq n - 1$ and $\alpha_n = e_{n-1} + e_n$, and the highest root is $\varphi = e_1 + e_2$. Let $f(x)$ be the quadratic form

$$f(x) = x_1x_{n+1} + \cdots + x_nx_{2n}$$

on \mathbb{K}^{2n} , and let (x, y) be the associated symmetric bilinear form $(x, y) = f(x + y) - f(x) - f(y)$. The Chevalley generators are

$$\begin{aligned} x_{e_i - e_j}(a) &= I + a(E_{i,j} - E_{n+j, n+i}) \\ x_{e_i + e_j}(a) &= I + a(E_{i, n+j} - E_{j, n+i}). \end{aligned}$$

Moreover, the diagram automorphism (interchanging nodes $n - 1$ and n) has the matrix

$$\sigma = 1 - E_{n,n} - E_{2n, 2n} + E_{n, 2n} + E_{2n, n}.$$

The following proposition is the other direction of Theorem 10. Recall the definition of $U(X)$ from Section 1.4.

Proposition 4.4. *Let $X = (\Gamma, J, \pi)$ be a polar closed admissible diagram of classical type. Every generic element of $U(X)$ has opposition diagram X .*

Proof. Let θ be a generic element of $U(X)$. By [13, Lemmas 1.1 and 3.5] the element θ maps the chamber w_0B to Weyl distance $w_{S \setminus J}w_0$, and hence the type J -simplex of this chamber is mapped onto an opposite simplex by θ . Thus $J \subseteq \text{Typ}(\theta)$. It remains to prove the reverse inclusion. We consider each nonempty polar closed diagram X , specifically, ${}^2A_{n,i}^1$ ($0 \leq i \leq n/2$), $B_{n,i}^1$ (with i even and $0 \leq i < n$), $B_{n,n}^1, B_{n,i}^2$ ($0 \leq i \leq n/2$), $C_{n,i}^1$ (for $0 \leq i \leq n/2$), $D_{n,i}^1$ (with i even), and $D_{n,i}^2$ (with $0 \leq i \leq n/2$).

In this proof (and only in this proof) “dimension” refers to the vector space dimension (rather than the projective dimension). Moreover, $\{i\}$ -domestic means domestic on type i vertices of the building.

(1) $X = A_{n,i}^1$ with $1 \leq i \leq n/2$. It is sufficient to show that θ is $\{j, n - j + 1\}$ -domestic for each $j > i$. The sequence of highest roots is $\varphi_k = e_k - e_{n-k+2}$ for $1 \leq k \leq i$. Thus

$$\theta = (I + a_1E_{1, n+1})(I + a_2E_{2, n}) \cdots (I + a_iE_{i, n-i+2}) = I + a_1E_{1, n+1} + \cdots + a_iE_{i, n-i+2}$$

with $a_1, \dots, a_i \neq 0$. Since θ pointwise fixes the $(n - i + 1)$ -dimensional space spanned by the vectors e_k with $1 \leq k \leq n - i + 1$, it follows that each space of dimension $j > i$ contains a fixed point. Thus if (V, V') is a type $\{j, n - j + 1\}$ -simplex (that is, $\dim(V) = j, \dim(V') = n - j + 1$, and $V \subseteq V'$) then V contains a fixed point. Thus $V^\theta \cap V' \neq \{0\}$, and so θ is $\{j, n - j + 1\}$ -domestic, hence the result.

(2) $X = B_{n, 2i}^1$ with $1 \leq i \leq n/2$. It is sufficient to show that θ is $\{j\}$ -domestic for $j > 2i$ and is not $\{1\}$ -domestic. The highest roots are φ_k with $k = 1, \dots, 2i$, where $\varphi_k = e_{2k-1} + e_{2k}$ and $\varphi_{i+k} = e_{2k-1} - e_{2k}$ for $1 \leq k \leq i$. Thus $\theta = \prod_{k=1}^i x_{e_{2k-1} + e_{2k}}(a_k)x_{e_{2k-1} - e_{2k}}(b_k)$ with $a_1, \dots, a_i, b_1, \dots, b_i \neq 0$, and so

$$\theta = I + \sum_{k=1}^i (a_k(E_{2k-1, n+2k} - E_{2k, n+2k-1}) + b_k(E_{2k-1, 2k} - E_{n+2k, n+2k-1}) + a_k b_k E_{2k-1, n+2k-1}).$$

The $(2n - 2i + 1)$ -dimensional space spanned by the vectors e_k with $k \notin \{2, 4, \dots, 2i\} \cup \{n + 1, n + 2, \dots, n + 2i\}$ and the vectors $e_{n+2k} - a_k b_k^{-1} e_{2k}$ with $k = 1, \dots, i$ is fixed pointwise by θ , and so θ is j -domestic for all $j > 2i$. Let $x = e_1 + e_2 + e_{n+1} - e_{n+2}$. Then $\langle x, x \rangle = 0$ (so $\langle x \rangle$ is a point of the polar space), we compute $\langle x, \theta x \rangle = -a_1 b_1 \neq 0$, and so this point is mapped onto an opposite point by θ .

(3) $X = B_{n,n}^1$. The first paragraph of the proof shows that θ maps a chamber to an opposite chamber, hence the result.

(4) $X = B_{n,i}^2$ with $1 \leq i \leq n/2$. It is sufficient to show that θ is $\{1\}$ -domestic, and $\{j\}$ -domestic for all $j > 2i$. The highest roots are $\varphi_k = e_{2k-1} + e_{2k}$ for $1 \leq k \leq i$, and so

$$\theta = \prod_{k=1}^i (I + a_k (E_{2k-1, n+2k} - E_{2k, n+2k-1})) = I + \sum_{k=1}^i a_k (E_{2k-1, n+2k} - E_{2k, n+2k-1})$$

with $a_1, \dots, a_i \neq 0$. The $(2n - 2i + 1)$ -dimensional space spanned by the vectors e_k with $k \notin \{n + 1, n + 2, \dots, n + 2i\}$ is fixed pointwise by θ , and it follows that every subspace of dimension $j > 2i$ contains a fixed point. Thus θ is $\{j\}$ -domestic for all $j > 2i$. Moreover, for each point $p = \langle x \rangle$ we compute $\langle x, \theta x \rangle = \langle x, x \rangle = 0$, and so θ is $\{1\}$ -domestic.

(5) $X = C_{n,i}^1$ with $1 \leq i \leq n$. It is sufficient to show that θ is $\{j\}$ -domestic for $j > i$ and that θ is not point domestic. The highest roots are $\varphi_k = 2e_k$ for $1 \leq k \leq i$. Thus

$$\theta = \prod_{k=1}^i (I + a_k E_{k, n+k}) = I + \sum_{k=1}^i a_k E_{k, n+k}.$$

Thus θ fixes the $(2n - i)$ -dimensional space spanned by e_k with $k \notin \{n + 1, \dots, n + i\}$. Thus θ is $\{j\}$ -domestic for all $j > i$. Since $\theta e_{n+1} = a_1 e_1 + e_{n+1}$ we have $\langle e_{n+1}, \theta e_{n+1} \rangle = -a_1 \neq 0$, and so $\langle e_{n+1} \rangle$ is mapped onto an opposite point, hence the result.

(6) $X = D_{n,2i}^1$ with $1 \leq i \leq n/2$. The highest roots are as in the $B_{n,2i}^1$ case, and the argument is identical to (2).

(7) $X = D_{n,i}^2$ with $1 \leq i \leq n/2$. The highest roots are as in the $B_{n,i}^2$ case, and the argument is identical to (3). \square

Finally we prove Theorem 11.

Proof of Theorem 11. The non-polar closed type preserving admissible diagrams are $B_{n,i}^1$ with i odd and $1 \leq i < n$ and $C_{n,i}^2$ for $1 \leq i \leq n/2$. If the underlying field is of characteristic 2 then, by assumption, the field is perfect and one can apply the previous proposition and duality. Thus suppose that the characteristic is not 2. We will show that these diagrams can be obtained by homologies of the associated buildings.

(1) $X = B_{n,i}^1$ with i odd and $1 \leq i < n$. More generally, we show that for $1 \leq i < n$ the homology

$$\theta = \text{diag}(\epsilon, \underbrace{-1, \dots, -1}_{i'}, \underbrace{1, \dots, 1}_{n-i'}, \underbrace{-1, \dots, -1}_{i'}, \underbrace{1, \dots, 1}_{n-i'})$$

where $\epsilon = (-1)^i$ and $i' = \lfloor i/2 \rfloor$, has opposition diagram $B_{n,i}^1$. Note that θ preserves the form (\cdot, \cdot) and hence $\theta \in B_n(\mathbb{F})$. Since θ fixes a subspace of dimension $2n - i + 1$ pointwise, every subspace of dimension $j \geq i + 1$ contains a fixed point, and so θ is j -domestic for each $j \geq i + 1$. If i is odd then the point $p = \langle e_0 + e_n - e_{2n} \rangle$ is mapped onto an opposite point, and if i is even then the point $p = \langle e_0 + e_1 - e_{n+1} \rangle$ is mapped onto an opposite point. Thus θ is not point domestic.

We now show that θ is not i -domestic. Suppose that i is even. Let $x_j = e_j + e_{2n-j+1}$ and $y_j = e_{n+j} - e_{n-j+1}$ for $1 \leq j \leq i' = n/2$. Then $V = \langle x_1, \dots, x_{i'}, y_1, \dots, y_{i'} \rangle$ is totally isotropic. If $V \ni p = \langle v \rangle$ with $v = a_1 x_1 + \dots + a_{i'} x_{i'} + b_1 y_1 + \dots + b_{i'} y_{i'}$ with $a_j \neq 0$ then p is opposite $\theta \langle y_j \rangle \in \theta_i V$, because $\langle v, \theta y_j \rangle = -2a_j$. Similarly, if $b_j \neq 0$ then p is opposite $\theta \langle x_j \rangle \in \theta V$. Thus V and θV are opposite. Suppose now that i is odd. There exists an integer k with $i' < k < n - i' + 1$, and thus both e_k and e_{n+k} are fixed by θ . Let $x_1, \dots, x_{i'}$ and $y_1, \dots, y_{i'}$ be as above, and let $z = e_0 + e_k - e_{n+k}$. Then $\langle z, z \rangle = \langle z, x_j \rangle = \langle z, y_j \rangle = 0$ for all $1 \leq j \leq i'$, and thus $V = \langle z, x_1, \dots, x_{i'}, y_1, \dots, y_{i'} \rangle$ is a totally isotropic space of dimension i . If $p = \langle v \rangle$ with $v = cz + a_1 x_1 + \dots + a_{i'} x_{i'} + b_1 y_1 + \dots + b_{i'} y_{i'}$ with $c \neq 0$

then p is opposite $\theta(z) \in \theta V$. If $c = 0$ then either $a_j \neq 0$ for some j (in which case p is opposite $\theta(y_j)$) or $b_j \neq 0$ for some j (in which case p is opposite $\theta(x_j)$). Thus V is opposite θV , hence the result.

(2) $X = C_{n,i}^2$ with $1 \leq i \leq n/2$. Let θ be the $(2n - 2i, 2i)$ -homology

$$\theta = \text{diag}(\underbrace{-1, \dots, -1}_i, \underbrace{1, \dots, 1}_{n-i}, \underbrace{-1, \dots, -1}_i, \underbrace{1, \dots, 1}_{n-i}).$$

Since θ fixes a space of dimension $2n - 2i$ pointwise, θ is j -domestic for all $j \geq 2i + 1$. Moreover, since $(x, \theta x) = 0$ for all $x \in \mathbb{F}^{2n}$ we see that θ is point-domestic. Let $x_j = e_j + f_{n-j+1}$ and $y_j = e_{n-j+1} + f_j$ for $1 \leq j \leq n/2$. Let $V = \langle x_1, \dots, x_i, y_1, \dots, y_i \rangle$ and note that V is totally isotropic. We claim that V and θV are opposite. If $p = \langle v \rangle$ with $v = a_1 x_1 + \dots + a_i x_i + b_1 y_1 + \dots + b_i y_i$ with $a_j \neq 0$ for some j then p and $\theta(y_j)$ are opposite, and if $b_j \neq 0$ for some j then p and $\theta(x_j)$ are opposite. Hence θ has opposition diagram $C_{n,i}^2$.

We now consider the non-type preserving diagrams.

(3) $X = A_{n;(n-1)/2}^2$ with $n \geq 3$ odd. Let (\cdot, \cdot) be any non-degenerate symplectic form on \mathbb{F}^{n+1} (note that $n + 1$ is even), and for subspaces U write $U^\circ = \{v \in \mathbb{F}^{n+1} \mid (u, v) = 0 \text{ for all } u \in U\}$. Then the map with $U^\theta = U^\circ$ is a symplectic polarity, and has the required diagram.

(4) $X = {}^2B_{2,1}^1$ or $X = {}^2C_{2,1}^1$, that is, we consider dualities of buildings of types B_2 and C_2 for perfect fields of characteristic 2. Every duality has this diagram.

(5) $X = D_{n,2i+1}^1$ with $0 \leq i < (n - 1)/2$. Let $0 \leq i \leq n/2$ and let

$$\begin{aligned} \theta_i &= I + \sum_{j=1}^i (E_{2j-1,2j} - E_{n+2j,n+2j-1}) \\ \eta &= I + E_{n-1,n} + E_{n-1,2n} - E_{n-1,2n-1} - E_{n,2n-1} - E_{2n,2n-1} \\ \sigma &= I - E_{n,n} - E_{2n,2n} + E_{n,2n} + E_{2n,n}, \end{aligned}$$

with $\theta_0 = I$. Similar calculations to the above show that for $0 \leq i \leq n/2$ the collineation θ_i has opposition diagram $D_{n,i}^2$, the collineation $\eta_i = \theta_i \eta$ has opposition diagram $D_{n,2i}^1$ and the duality $\sigma_i = \theta_i \sigma$ has opposition diagram $D_{n,2i+1}^1$.

Finally, trialities of buildings of type D_4 are treated in [22], and the proof is complete. \square

References

- [1] P. Abramenko, K. S. Brown, *Buildings*. Springer 2008. MR2439729 Zbl 1214.20033
- [2] P. Abramenko, K. S. Brown, Automorphisms of non-spherical buildings have unbounded displacement. *Innov. Incidence Geom.* **10** (2009), 1–13. MR2665192 Zbl 1276.51004
- [3] N. Bourbaki, *Lie groups and Lie algebras. Chapters 4–6*. Springer 2002. MR1890629 Zbl 0983.17001
- [4] R. W. Carter, *Simple groups of Lie type*. Wiley-Interscience 1989. MR1013112 Zbl 0723.20006
- [5] A. M. Cohen, E. E. Shult, Affine polar spaces. *Geom. Dedicata* **35** (1990), 43–76. MR1066558 Zbl 0755.51004
- [6] B. De Bruyn, *An introduction to incidence geometry*. Springer 2016. MR3585811 Zbl 1376.51001
- [7] B. De Bruyn, H. Van Maldeghem, Non-embeddable polar spaces. *Münster J. Math.* **7** (2014), 557–588. MR3426230 Zbl 1345.51004
- [8] B. Leeb, *A characterization of irreducible symmetric spaces and Euclidean buildings of higher rank by their asymptotic geometry*, volume 326 of *Bonner Mathematische Schriften*. Universität Bonn, Mathematisches Institut, Bonn 2000. MR1934160 Zbl 1005.53031
- [9] B. Mühlherr, H. P. Petersson, R. M. Weiss, *Descent in buildings*, volume 190 of *Annals of Mathematics Studies*. Princeton Univ. Press 2015. MR3364836 Zbl 1338.51002
- [10] J. Parkinson, B. Temmermans, H. Van Maldeghem, The combinatorics of automorphisms and opposition in generalised polygons. *Ann. Comb.* **19** (2015), 567–619. MR3395495 Zbl 1327.05048
- [11] J. Parkinson, H. Van Maldeghem, Opposition diagrams for automorphisms of large spherical buildings. *J. Combin. Theory Ser. A* **162** (2019), 118–166. MR3873873 Zbl 1411.51006
- [12] J. Parkinson, H. Van Maldeghem, Opposition diagrams for automorphisms of small spherical buildings. *Innov. Incidence Geom.* **17** (2019), 141–188. MR3956902 Zbl 1472.51005
- [13] J. Parkinson, H. Van Maldeghem, Automorphisms and opposition in spherical buildings of exceptional type, I. *Canad. J. Math.* **74** (2022), 1517–1578. MR4520661 Zbl 1510.20031

- [14] J. Parkinson, H. Van Maldeghem, Automorphisms and opposition in spherical buildings of exceptional type, II: Moufang hexagons. *Innov. Incidence Geom.* **20** (2023), 443–470. MR4641401 Zbl 07760467
- [15] A. E. Schroth, Characterizing symplectic quadrangles by their derivations. *Arch. Math. (Basel)* **58** (1992), 98–104. MR1139392 Zbl 0761.51004
- [16] R. Steinberg, *Lectures on Chevalley groups*, volume 66 of *University Lecture Series*. Amer. Math. Soc. 2016. MR3616493 Zbl 1361.20003
- [17] B. Temmermans, J. A. Thas, H. Van Maldeghem, Domesticity in projective spaces. *Innov. Incidence Geom.* **12** (2011), 141–149. MR2942721 Zbl 1305.51004
- [18] B. Temmermans, J. A. Thas, H. Van Maldeghem, Collineations of polar spaces with restricted displacements. *Des. Codes Cryptogr.* **64** (2012), 61–80. MR2914402 Zbl 1242.51002
- [19] B. Temmermans, J. A. Thas, H. Van Maldeghem, Domesticity in generalized quadrangles. *Ann. Comb.* **16** (2012), 905–916. MR3000452 Zbl 1259.51004
- [20] J. Tits, *Buildings of spherical type and finite BN-pairs*. Springer 1974. MR470099 Zbl 0295.20047
- [21] H. Van Maldeghem, *Generalized polygons*. Birkhäuser Verlag, Basel 1998. MR1725957 Zbl 0914.51005
- [22] H. Van Maldeghem, Characterizations of trialities of type I_{ii} in buildings of type D_4 . In: *Groups of exceptional type, Coxeter groups and related geometries*, volume 82 of *Springer Proc. Math. Stat.*, 205–216, Springer 2014. MR3207278 Zbl 1338.51012